

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problems Mailbox.**



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification⁵ : C12N 1/24, 15/00, C07H 21/00	A1	(11) International Publication Number: WO 92/06176 (43) International Publication Date: 16 April 1992 (16.04.92)
(21) International Application Number: PCT/US91/07141 (22) International Filing Date: 27 September 1991 (27.09.91) (30) Priority data: 590,664 28 September 1990 (28.09.90) US (71) Applicant: IXSYS, INC. [US/US]; 3550 General Atomics Court, Suite L103, San Diego, CA 92121 (US). (72) Inventor: HUSE, William, D. ; 471 Avenida Primavera, Del Mar, CA 92014 (US). (74) Agents: CAMPBELL, Cathryn et al.; Pretty, Schroeder, Brueggemann & Clark, 444 South Flower Street, Suite 2000, Los Angeles, CA 90071 (US).		(81) Designated States: AT (European patent), AU, BB, BE (European patent), BF (OAPI patent), BG, BJ (OAPI patent), BR, CA, CF (OAPI patent), CG (OAPI patent), CH (European patent), CI (OAPI patent), CM (OAPI patent), CS, DE (European patent), DK (European patent), ES (European patent), FI, FR (European patent), GA (OAPI patent), GB (European patent), GN (OAPI patent), GR (European patent), HU, IT (European patent), JP, KP, KR, LK, LU (European patent), MC, MG, ML (OAPI patent), MN, MR (OAPI patent), MW, NL (European patent), NO, PL, RO, SD, SE (European patent), SN (OAPI patent), SU ⁺ , TD (OAPI patent), TG (OAPI patent). Published <i>With international search report.</i>
(54) Title: SURFACE EXPRESSION LIBRARIES OF RANDOMIZED PEPTIDES (57) Abstract A composition of matter comprising a plurality of procaryotic cells containing a diverse population of expressible oligonucleotides operationally linked to expression elements, said expressible oligonucleotides having a desirable bias of random codon sequences.		

+ DESIGNATIONS OF "SU"

Any designation of "SU" has effect in the Russian Federation. It is not yet known whether any such designation has effect in other States of the former Soviet Union.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	ES	Spain	MG	Madagascar
AU	Australia	FI	Finland	ML	Mali
BB	Barbados	FR	France	MN	Mongolia
BE	Belgium	GA	Gabon	MR	Mauritania
BF	Burkina Faso	GB	United Kingdom	MW	Malawi
BG	Bulgaria	GN	Guinea	NL	Netherlands
BJ	Benin	GR	Greece	NO	Norway
BR	Brazil	HU	Hungary	PL	Poland
CA	Canada	IT	Italy	RO	Romania
CF	Central African Republic	JP	Japan	SD	Sudan
CG	Congo	KP	Democratic People's Republic of Korea	SE	Sweden
CH	Switzerland	KR	Republic of Korea	SN	Senegal
CI	Côte d'Ivoire	LI	Liechtenstein	SU ⁺	Soviet Union
CM	Cameroon	LK	Sri Lanka	TD	Chad
CS	Czechoslovakia	LU	Luxembourg	TG	Togo
DE	Germany	MC	Monaco	US	United States of America
DK	Denmark				

SURFACE EXPRESSION LIBRARIES
OF RANDOMIZED PEPTIDES

5

BACKGROUND OF THE INVENTION

This invention relates generally to methods for synthesizing and expressing oligonucleotides and, more particularly, to methods for expressing oligonucleotides having random codon sequences.

10 Oligonucleotide synthesis proceeds via linear coupling of individual monomers in a stepwise reaction. The reactions are generally performed on a solid phase support by first coupling the 3' end of the first monomer to the support. The second monomer is added to the 5' end of the
15 first monomer in a condensation reaction to yield a dinucleotide coupled to the solid support. At the end of each coupling reaction, the by-products and unreacted, free monomers are washed away so that the starting material for the next round of synthesis is the pure oligonucleotide
20 attached to the support. In this reaction scheme, the stepwise addition of individual monomers to a single, growing end of a oligonucleotide ensures accurate synthesis of the desired sequence. Moreover, unwanted side reactions are eliminated, such as the condensation of two
25 oligonucleotides, resulting in high product yields.

In some instances, it is desired that synthetic oligonucleotides have random nucleotide sequences. This result can be accomplished by adding equal proportions of all four nucleotides in the monomer coupling reactions,
30 leading to the random incorporation of all nucleotides and yielding a population of oligonucleotides with random sequences. Since all possible combinations of nucleotide sequences are represented within the population, all possible codon triplets will also be represented. If the

objective is ultimately to generate random peptide products, this approach has a severe limitation because the random codons synthesized will bias the amino acids incorporated during translation of the DNA by the cell into polypeptides.

The bias is due to the redundancy of the genetic code. There are four nucleotide monomers which leads to sixty-four possible triplet codons. With only twenty amino acids to specify, many of the amino acids are encoded by multiple codons. Therefore, a population of oligonucleotides synthesized by sequential addition of monomers from a random population will not encode peptides whose amino acid sequence represents all possible combinations of the twenty different amino acids in equal proportions. That is, the frequency of amino acids incorporated into polypeptides will be biased toward those amino acids which are specified by multiple codons.

To alleviate amino acid bias due to the redundancy of the genetic code, the oligonucleotides can be synthesized from nucleotide triplets. Here, a triplet coding for each of the twenty amino acids is synthesized from individual monomers. Once synthesized, the triplets are used in the coupling reactions instead of individual monomers. By mixing equal proportions of the triplets, synthesis of oligonucleotides with random codons can be accomplished. However, the cost of synthesis from such triplets far exceeds that of synthesis from individual monomers because triplets are not commercially available.

Amino acid bias can be reduced, however, by synthesizing the degenerate codon sequence NNK where N is a mixture of all four nucleotides and K is a mixture guanine and thymine nucleotides. Each position within an oligonucleotide having this codon sequence will contain a total of 32 codons (12 encoding amino acids being

r presented once, 5 represented twice, 3 represented three times and one codon being a stop c don). Oligonucleotides expressed with such degenerate codon sequences will produce peptide products whose sequences are biased toward those amino acids being represented more than once. Thus, populations of peptides whose sequences are completely random cannot be obtained from oligonucleotides synthesized from degenerate sequences.

There thus exists a need for a method to express oligonucleotides having a fully random or desirably biased sequence which alleviates genetic redundancy. The present invention satisfies these needs and provides additional advantages as well.

SUMMARY OF THE INVENTION

The invention provides a plurality of procaryotic cells containing a diverse population of expressible oligonucleotides operationally linked to expression elements, the expressible oligonucleotides having a desirable bias of random codon sequences.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 is a schematic drawing for synthesizing oligonucleotides from nucleotide monomers with random tuplets at each position using twenty reaction vessels.

Figure 2 is a schematic drawing for synthesizing oligonucleotides from nucleotide monomers with random tuplets at each position using ten reaction vessels.

Figure 3 is a schematic diagram of the two vectors used for sublibrary and library production from precursor oligonucleotide portions. M13IX22 (Figure 3A) is the vector used to clone the anti-sense precursor portions

(hatched box). The single-headed arrow represents the Lac p/o expression sequences and the double-headed arrow represents the portion of M13IX22 which is to be combined with M13IX42. The amber stop codon for biological selection and relevant restriction sites are also shown. M13IX42 (Figure 3B) is the vector used to clone the sense precursor portions (open box). Thick lines represent the pseudo-wild type (ψ gVIII) and wild type (gVIII) gene VIII sequences. The double-headed arrow represents the portion of M13IX42 which is to be combined with M13IX22. The two amber stop codons and relevant restriction sites are also shown. Figure 3C shows the joining of vector population from sublibraries to form the functional surface expression vector M13IX. Figure 3D shows the generation of a surface expression library in a non-suppressor strain and the production of phage. The phage are used to infect a suppressor strain (Figure 3E) for surface expression and screening of the library.

Figure 4 is a schematic diagram of the vector used for generation of surface expression libraries from random oligonucleotide populations (M13IX30). The symbols are as described for Figure 3.

Figure 5 is the nucleotide sequence of M13IX42 (SEQ ID NO: 1).

Figure 6 is the nucleotide sequence of M13IX22 (SEQ ID NO: 2).

Figure 7 is the nucleotide sequence of M13IX30 (SEQ ID NO: 3).

Figure 8 is the nucleotide sequence of M13ED03 (SEQ ID NO: 4).

Figure 9 is the nucleotide sequence of M13IX421 (SEQ

ID NO: 5).

Figure 10 is the nucleotide sequence of M13ED04 (SEQ ID NO: 6).

DETAILED DESCRIPTION OF THE INVENTION

5 This invention is directed to a simple and inexpensive method for synthesizing and expressing oligonucleotides having a desirable bias of random codons using individual monomers. The method is advantageous in that individual monomers are used instead of triplets and by synthesizing
10 only a non-degenerate subset of all triplets, codon redundancy is alleviated. Thus, the oligonucleotides synthesized represent a large proportion of possible random triplet sequences which can be obtained. The oligonucleotides can be expressed, for example, on the
15 surface of filamentous bacteriophage in a form which does not alter phage viability or impose biological selections against certain peptide sequences. The oligonucleotides produced are therefore useful for generating an unlimited number of pharmacological and research products.

20 In one embodiment, the invention entails the sequential coupling of monomers to produce oligonucleotides with a desirable bias of random codons. The coupling reactions for the randomization of twenty codons which specify the amino acids of the genetic code are performed
25 in ten different reaction vessels. Each reaction vessel contains a support on which the monomers for two different codons are coupled in three sequential reactions. One of the reactions couples an equal mixture of two monomers such that the final product has two different codon sequences.
30 The codons are randomized by removing the supports from the reaction vessels and mixing them to produce a single batch of supports containing all twenty codons at a particular position. Synthesis at the next codon position proceeds by

equally dividing the mixed batch of supports into ten reaction vessels as before and sequentially coupling the monomers for each pair of codons. The supports are again mixed to randomize the codons at the position just synthesized. The cycle of coupling, mixing and dividing continues until the desired number of codon positions have been randomized. After the last position has been randomized, the oligonucleotides with random codons are cleaved from the support. The random oligonucleotides can then be expressed, for example, on the surface of filamentous bacteriophage as gene VIII-peptide fusion proteins. Alternative genes can be used as well.

In its broadest form, the invention provides a diverse population of synthetic oligonucleotides contained in vectors so as to be expressible in cells. Such populations of diverse oligonucleotides can be fully random at one or more codon sites or can be fully defined at one or more site, so long as at least one site the codons are randomly variable. The populations of oligonucleotides can be expressed as fusion products in combination with surface proteins of filamentous bacteriophage, such as M13, as with gene VIII. The vectors can be transfected into a plurality of cells, such as the procaryote E. coli.

The diverse population of oligonucleotides can be formed by randomly combining first and second precursor populations, each precursor population having a desirable bias of random codon sequences. Methods of synthesizing and expressing the diverse population of expressible oligonucleotides are also provided.

In a preferred embodiment, two populations of random oligonucleotides are synthesized. The oligonucleotides within each population encode a portion of the final oligonucleotide which is to be expressed. Oligonucleotides within one population encode the carboxy terminal portion

of the expressed oligonucleotides. These oligonucleotides are cloned in frame with a gene VIII (gVIII) sequence so that translation of the sequence produces peptide fusion proteins. The second population of oligonucleotides are
5 cloned into a separate vector. Each oligonucleotide within this population encodes the anti-sense of the amino terminal portion of the expressed oligonucleotides. This vector also contains the elements necessary for expression. The two vectors containing the random oligonucleotides are
10 combined such that the two precursor oligonucleotide portions are joined together at random to form a population of larger oligonucleotides derived from two smaller portions. The vectors contain selectable markers to ensure maximum efficiency in joining together the two
15 oligonucleotide populations. A mechanism also exists to control the expression of gVIII-peptide fusion proteins during library construction and screening.

As used herein, the term "monomer" or "nucleotide monomer" refers to individual nucleotides used in the
20 chemical synthesis of oligonucleotides. Monomers that can be used include both the ribo- and deoxyribo- forms of each of the five standard nucleotides (derived from the bases adenine (A or dA, respectively), guanine (G or dG), cytosine (C or dC), thymine (T) and uracil (U)).
25 Derivatives and precursors of bases such as inosine which are capable of supporting polypeptide biosynthesis are also included as monomers. Also included are chemically modified nucleotides, for example, one having a reversible blocking agent attached to any of the positions on the
30 purine or pyrimidine bases, the ribose or deoxyribose sugar or the phosphate or hydroxyl moieties of the monomer. Such blocking groups include, for example, dimethoxytrityl, benzoyl, isobutyryl, beta-cyanoethyl and diisopropylamine groups, and are used to protect hydr xyls, exocyclic amin s
35 and phosphate moieties. Oth r blocking agents can also be used and are known to one skilled in the art.

As used herein, the term "tuplet" refers to a group of elements of a definable size. The elements of a tuplet as used herein are nucleotide monomers. For example, a tuplet can be a dinucleotide, a trinucleotide or can also be four
5 or more nucleotides.

As used herein, the term "codon" or "triplet" refers to a tuplet consisting of three adjacent nucleotide monomers which specify one of the twenty naturally occurring amino acids found in polypeptide biosynthesis.
10 The term also includes nonsense, or stop, codons which do not specify any amino acid.

"Random codons" or "randomized codons," as used herein, refers to more than one codon at a position within a collection of oligonucleotides. The number of different
15 codons can be from two to twenty at any particular position. "Randomized oligonucleotides," as used herein, refers to a collection of oligonucleotides with random codons at one or more positions. "Random codon sequences" as used herein means that more than one codon position
20 within a randomized oligonucleotide contains random codons. For example, if randomized oligonucleotides are six nucleotides in length (i.e., two codons) and both the first and second codon positions are randomized to encode all twenty amino acids, then a population of oligonucleotides
25 having random codon sequences with every possible combination of the twenty triplets in the first and second position makes up the above population of randomized oligonucleotides. The number of possible codon combinations is 20^2 . Likewise, if randomized
30 oligonucleotides of fifteen nucleotides in length are synthesized which have random codon sequences at all positions encoding all twenty amino acids, then all triplets coding for each of the twenty amino acids will be found in equal proportions at every position. The
35 population constituting the randomized oligonucleotides

will contain 20^{15} different possible species of oligonucleotides. "Random tuplets," or "randomized tuplets" are defined analogously.

As used herein, the term "bias" refers to a preference. It is understood that there can be degrees of preference or bias toward codon sequences which encode particular amino acids. For example, an oligonucleotide whose codon sequences do not preferably encode particular amino acids is unbiased and therefore completely random. The oligonucleotide codon sequences can also be biased toward predetermined codon sequences or codon frequencies and while still diverse and random, will exhibit codon sequences biased toward a defined, or preferred, sequence. "A desirable bias of random codon sequences" as used herein, refers to the predetermined degree of bias which can be selected from totally random to essentially, but not totally, defined (or preferred). There must be at least one codon position which is variable, however.

As used herein, the term "support" refers to a solid phase material for attaching monomers for chemical synthesis. Such support is usually composed of materials such as beads of control pore glass but can be other materials known to one skilled in the art. The term is also meant to include one or more monomers coupled to the support for additional oligonucleotide synthesis reactions.

As used herein, the terms "coupling" or "condensing" refers to the chemical reactions for attaching one monomer to a second monomer or to a solid support. Such reactions are known to one skilled in the art and are typically performed on an automated DNA synthesizer such as a MilliGen/Biosearch Cyclone Plus Synthesizer using procedures recommended by the manufacturer. "Sequentially coupling" as used herein, refers to the stepwise addition of monomers.

A method of synthesizing oligonucleotides having random tuplets using individual monomers is described. The method consists of several steps, the first being synthesis of a nucleotide tuplet for each tuplet to be randomized.

5 As described here and below, a nucleotide triplet (i.e., a codon) will be used as a specific example of a tuplet. Any size tuplet will work using the methods disclosed herein, and one skilled in the art would know how to use the methods to randomize tuplets of any size.

10 If the randomization of codons specifying all twenty amino acids is desired at a position, then twenty different codons are synthesized. Likewise, if randomization of only ten codons at a particular position is desired then those ten codons are synthesized. Randomization of codons from

15 two to sixty-four can be accomplished by synthesizing each desired triplet. Preferably, randomization of from two to twenty codons is used for any one position because of the redundancy of the genetic code. The codons selected at one position do not have to be the same codons selected at the

20 next position. Additionally, the sense or anti-sense sequence oligonucleotide can be synthesized. The process therefore provides for randomization of any desired codon position with any number of codons.

Codons to be randomized are synthesized sequentially

25 by coupling the first monomer of each codon to separate supports. The supports for the synthesis of each codon can, for example, be contained in different reaction vessels such that one reaction vessel corresponds to the monomer coupling reactions for one codon. As will be used

30 here and below, if twenty codons are to be randomized, then twenty reaction vessels can be used in independent coupling reactions for the first twenty monomers of each codon. Synthesis proceeds by sequentially coupling the second monomer of each codon to the first monomer to produce a

35 dimer, followed by coupling the third monomer for each

codon to each of the above-synthesized dimers to produce a trimer (Figure 1, step 1, where M_1 , M_2 and M_3 represent the first, second and third monomer, respectively, for each codon to be randomized).

5 Following synthesis of the first codons from individual monomers, the randomization is achieved by mixing the supports from all twenty reaction vessels which contain the individual codons to be randomized. The solid phase support can be removed from its vessel and mixed to
10 achieve a random distribution of all codon species within the population (Figure 1, step 2). The mixed population of supports, constituting all codon species, are then redistributed into twenty independent reaction vessels (Figure 1, step 3). The resultant vessels are all
15 identical and contain equal portions of all twenty codons coupled to a solid phase support.

For randomization of the second position codon, synthesis of twenty additional codons is performed in each of the twenty reaction vessels produced in step 3 as the
20 condensing substrates of step 1 (Figure 1, step 4). Steps 1 and 4 are therefore equivalent except that step 4 uses the supports produced by the previous synthesis cycle (steps 1 through 3) for codon synthesis whereas step 1 is the initial synthesis of the first codon in the
25 oligonucleotide. The supports resulting from step 4 will each have two codons attached to them (i.e., a hexanucleotide) with the codon at the first position being any one of twenty possible codons (i.e., random) and the codon at the second position being one of the twenty
30 possible codons.

For randomization of the codon at the second position and synthesis of the third position codon, steps 2 through 4 are again repeated. This process yields in each vessel a three codon oligonucleotide (i.e., 9 nucleotides) with

codon positions 1 and 2 randomized and position three containing one of the twenty possible codons. Steps 2 through 4 are repeated to randomize the third position codon and synthesize the codon at the next position. The process is continued until an oligonucleotide of the desired length is achieved. After the final randomization step, the oligonucleotide can be cleaved from the supports and isolated by methods known to one skilled in the art. Alternatively, the oligonucleotides can remain on the supports for use in methods employing probe hybridization.

The diversity of codon sequences, i.e., the number of different possible oligonucleotides, which can be obtained using the methods of the present invention, is extremely large and only limited by the physical characteristics of available materials. For example, a support composed of beads of about 100 μm in diameter will be limited to about 10,000 beads/reaction vessel using a 1 μM reaction vessel containing 25 mg of beads. This size bead can support about 1×10^7 oligonucleotides per bead. Synthesis using separate reaction vessels for each of the twenty amino acids will produce beads in which all the oligonucleotides attached to an individual bead are identical. The diversity which can be obtained under these conditions is approximately 10^7 copies of 10,000 x 20 or 200,000 different random oligonucleotides. The diversity can be increased, however, in several ways without departing from the basic methods disclosed herein. For example, the number of possible sequences can be increased by decreasing the size of the individual beads which make up the support. A bead of about 30 μm in diameter will increase the number of beads per reaction vessel and therefore the number of oligonucleotides synthesized. Another way to increase the diversity of oligonucleotides with random codons is to increase the volume of the reaction vessel. For example, using the same size bead, a larger volume can contain a greater number of beads than a smaller vessel and therefore

support the synthesis of a greater number of oligonucleotides. Increasing the number of codons coupled to a support in a single reaction vessel also increases the diversity of the random oligonucleotides. The total
5 diversity will be the number of codons coupled per vessel raised to the number of codon positions synthesized. For example, using ten reaction vessels, each synthesizing two codons to randomize a total of twenty codons, the number of different oligonucleotides of ten codons in length per 100
10 μm bead can be increased where each bead will contain about 2^{10} or 1×10^3 different sequences instead of one. One skilled in the art will know how to modify such parameters to increase the diversity of oligonucleotides with random codons.

15 A method of synthesizing oligonucleotides having random codons at each position using individual monomers wherein the number of reaction vessels is less than the number of codons to be randomized is also described. For example, if twenty codons are to be randomized at each
20 position within an oligonucleotide population, then ten reaction vessels can be used. The use of a smaller number of reaction vessels than the number of codons to be randomized at each position is preferred because the smaller number of reaction vessels is easier to manipulate
25 and results in a greater number of possible oligonucleotides synthesized.

The use of a smaller number of reaction vessels for random synthesis of twenty codons at a desired position within an oligonucleotide is similar to that described
30 above using twenty reaction vessels except that each reaction vessel can contain the synthesis products of more than one codon. For example, step one synthesis using ten reaction vessels proceeds by coupling about two different codons on supports contained in each of ten reaction
35 vessels. This is shown in Figur 2 where each of the two

codons coupled to a different support can consist of the following sequences: (1) (T/G)TT for Phe and Val; (2) (T/C)CT for Ser and Pro; (3) (T/C)AT for Tyr and His; (4) (T/C)GT for Cys and Arg; (5) (C/A)TG for Leu and Met; (6) 5 (C/G)AG for Gln and Glu; (7) (A/G)CT for Thr and Ala; (8) (A/G)AT for Asn and Asp; (9) (T/G)GG for Trp and Gly and (10) A(T/A)A for Ile and Cys. The slash (/) signifies that a mixture of the monomers indicated on each side of the slash are used as if they were a single monomer in the 10 indicated coupling step. The antisense sequence for each of the above codons can be generated by synthesizing the complementary sequence. For example, the antisense for Phe and Val can be AA(C/A). The amino acids encoded by each of the above pairs of sequences are given as the standard 15 three letter nomenclature.

Coupling of the monomers in this fashion will yield codons specifying all twenty of the naturally occurring amino acids attached to supports in ten reaction vessels. However, the number of individual reaction vessels to be 20 used will depend on the number of codons to be randomized at the desired position and can be determined by one skilled in the art. For example, if ten codons are to be randomized, then five reaction vessels can be used for coupling. The codon sequences given above can be used for 25 this synthesis as well. The sequences of the codons can also be changed to incorporate or be replaced by any of the additional forty-four codons which constitutes the genetic code.

The remaining steps of synthesis of oligonucleotides 30 with random codons using a smaller number of reaction vessels are as outlined above for synthesis with twenty reaction vessels except that the mixing and dividing steps are performed with supports from about half the number of reaction vessels. These remaining steps are shown in 35 Figure 2 (steps 2 through 4).

Oligonucleotides having at least one specified tuplet at a pred terminated position and the remaining positions having random tuplets can also be synthesized using the methods described herein. The synthesis steps are similar to those outlined above using twenty or less reaction vessels except that prior to synthesis of the specified codon position, the dividing of the supports into separate reaction vessels for synthesis of different codons is omitted. For example, if the codon at the second position of the oligonucleotide is to be specified, then following synthesis of random codons at the first position and mixing of the supports, the mixed supports are not divided into new reaction vessels but, instead, can be contained in a single reaction vessel to synthesize the specified codon. The specified codon is synthesized sequentially from individual monomers as described above. Thus, the number of reaction vessels can be increased or decreased at each step to allow for the synthesis of a specified codon or a desired number of random codons.

Following codon synthesis, the mixed supports are divided into individual reaction vessels for synthesis of the next codon to be randomized (Figure 1, step 3) or can be used without separation for synthesis of a consecutive specified codon. The rounds of synthesis can be repeated for each codon to be added until the desired number of positions with predetermined or randomized codons are obtained.

Synthesis of oligonucleotides with the first position codon being specified can also be synthesized using the above method. In this case, the first position codon is synthesized from the appropriate monomers. The supports are divided into the required number of reaction vessels needed for synthesis of random codons at the second position and the rounds of synthesis, mixing and dividing are performed as described above.

A method of synthesizing oligonucleotides having
tuplets which are diverse but biased toward a predetermined
sequence is also described herein. This method employs two
reaction vessels, one vessel for the synthesis of a
5 predetermined sequence and the second vessel for the
synthesis of a random sequence. This method is
advantageous to use when a significant number of codon
positions, for example, are to be of a specified sequence
since it alleviates the use of multiple reaction vessels.
10 Instead, a mixture of four different monomers such as
adenine, guanine, cytosine and thymine nucleotides are used
for the first and second monomers in the codon. The codon
is completed by coupling a mixture of a pair of monomers of
either guanine and thymine or cytosine and adenine
15 nucleotides at the third monomer position. In the second
vessel, nucleotide monomers are coupled sequentially to
yield the predetermined codon sequence. Mixing of the two
supports yields a population of oligonucleotides containing
both the predetermined codon and the random codons at the
20 desired position. Synthesis can proceed by using this
mixture of supports in a single reaction vessel, for
example, for coupling additional predetermined codons or,
further dividing the mixture into two reaction vessels for
synthesis of additional random codons.

25 The two reaction vessel method can be used for codon
synthesis within an oligonucleotide with a predetermined
tuplet sequence by dividing the support mixture into two
portions at the desired codon position to be randomized.
Additionally, this method allows for the extent of
30 randomization to be adjusted. For example, unequal mixing
or dividing of the two supports will change the fraction of
codons with predetermined sequences compared to those with
random codons at the desired position. Unequal mixing and
dividing of supports can be useful when there is a need to
35 synthesize random codons at a significant number of
positions within an oligonucleotide of a longer or shorter

length.

The extent of randomization can also be adjusted by using unequal mixtures of monomers in the first, second and third monomer coupling steps of the random codon position.

5 The unequal mixtures can be in any or all of the coupling steps to yield a population of codons enriched in sequences reflective of the monomer proportions.

Synthesis of randomized oligonucleotides is performed using methods well known to one skilled in the art. Linear

10 coupling of monomers can, for example, be accomplished using phosphoramidite chemistry with a MilliGen/Biosearch Cyclone Plus automated synthesizer as described by the manufacturer (Millipore, Burlington, MA). Other chemistries and automated synthesizers can be employed as

15 well and are known to one skilled in the art.

Synthesis of multiple codons can be performed without modification to the synthesizer by separately synthesizing the codons in individual sets of reactions. Alternatively, modification of an automated DNA synthesizer can be

20 performed for the simultaneous synthesis of codons in multiple reaction vessels.

In one embodiment, the invention provides a plurality of procaryotic cells containing a diverse population of expressible oligonucleotides operationally linked to

25 expression elements, the expressible oligonucleotides having a desirable bias of random codon sequences produced from diverse combinations of first and second oligonucleotides having a desirable bias of random sequences. The invention provides for a method for

30 constructing such a plurality of procaryotic cells as well.

The oligonucleotides synthesized by the above methods can be used to express a plurality of random peptides which

are unbiased, diverse but biased toward a predetermined sequence or which contain at least one specified codon at a predetermined position. The need will determine which type of oligonucleotide is to be expressed to give the
5 resultant population of random peptides and is known to one skilled in the art. Expression can be performed in any compatible vector/host system. Such systems include, for example, plasmids or phagemids in procaryotes such as E. coli, yeast systems, and other eucaryotic systems such as
10 mammalian cells, but will be described herein in context with its presently preferred embodiment, i.e. expression on the surface of filamentous bacteriophage. Filamentous bacteriophage can be, for example, M13, fl and fd. Such phage have circular single-stranded genomes and double
15 strand replicative DNA forms. Additionally, the peptides can also be expressed in soluble or secreted form depending on the need and the vector/host system employed.

Expression of random peptides on the surface of M13 can be accomplished, for example, using the vector system
20 shown in Figure 3. Construction of the vectors enabling one of ordinary skill to make them are explicitly set out in Examples I and II. The complete nucleotide sequences are given in Figures 5, 6 and 7 (SEQ ID NOS: 1, 2 and 3, respectively). This system produces random
25 oligonucleotides functionally linked to expression elements and to gVIII by combining two smaller oligonucleotide portions contained in separate vectors into a single vector. The diversity of oligonucleotide species obtained by this system or others described herein can be 5×10^7 or
30 greater. Diversity of less than 5×10^7 can also be obtained and will be determined by the need and type of random peptides to be expressed. The random combination of two precursor portions into a larger oligonucleotide increases the diversity of the population several fold and
35 has the added advantage of producing oligonucleotides larger than what can be synthesized by standard methods.

Additionally, although the correlation is not known, when the number of possible paths an oligonucleotide can take during synthesis such as described herein is greater than the number of beads, then there will be a correlation
5 between the synthesis path and the sequences obtained. By combining oligonucleotide populations which are synthesized separately, this correlation will be destroyed. Therefore, any bias which may be inherent in the synthesis procedures will be alleviated by joining two precursor portions into
10 a contiguous random oligonucleotide.

Populations of precursor oligonucleotides to be combined into an expressible form are each cloned into separate vectors. The two precursor portions which make up the combined oligonucleotide corresponds to the carboxy and
15 amino terminal portions of the expressed peptide. Each precursor oligonucleotide can encode either the sense or anti-sense and will depend on the orientation of the expression elements and the gene encoding the fusion portion of the protein as well as the mechanism used to
20 join the two precursor oligonucleotides. For the vectors shown in Figure 3, precursor oligonucleotides corresponding to the carboxy terminal portion of the peptide encode the sense strand. Those corresponding to the amino terminal portion encode the anti-sense strand. Oligonucleotide
25 populations are inserted between the Eco RI and Sac I restriction enzyme sites in M13IX22 and M13IX42 (Figure 3A and B). M13IX42 (SEQ ID NO: 1) is the vector used for sense strand precursor oligonucleotide portions and M13IX22 (SEQ ID NO: 2) is used for anti-sense precursor portions.

30 The populations of randomized oligonucleotides inserted into the vectors are synthesized with Eco RI and Sac I recognition sequences flanking opposite ends of the random codon sequences. The sites allow annealing and ligation of these single strand oligonucleotides into a
35 double stranded vector restricted with Eco RI and Sac I.

Alternatively, the oligonucleotides can be inserted into the vector by standard mutagenesis methods. In this latter method, single stranded vector DNA is isolated from the phage and annealed with random oligonucleotides having
5 known sequences complementary to vector sequences. The oligonucleotides are extended with DNA polymerase to produce double stranded vectors containing the randomized oligonucleotides.

The vector used for sense strand oligonucleotide
10 portions, M13IX42 (Figure 3B) contains down-stream and in frame with the Eco RI and Sac I restriction sites a sequence encoding the pseudo-wild type gVIII product. This gene encodes the wild type M13 gVIII amino acid sequence but has been changed at the nucleotide level to reduce
15 homologous recombination with the wild type gVIII contained on the same vector. The wild type gVIII is present to ensure that at least some functional, non-fusion coat protein will be produced. The inclusion of a wild type gVIII therefore reduces the possibility of non-viable phage
20 production and biological selection against certain peptide fusion proteins. Differential regulation of the two genes can also be used to control the relative ratio of the pseudo and wild type proteins.

Also contained downstream and in frame with the Eco RI
25 and Sac I restriction sites is an amber stop codon. The mutation is located six codons downstream from Sac I and therefore lies between the inserted oligonucleotides and the gVIII sequence. As was the function of the wild type gVIII, the amber stop codon also reduces biological
30 selection when combining precursor portions to produce expressible oligonucleotides. This is accomplished by using a non-suppressor (sup O) host strain because non-suppressor strains will terminate expression after the oligonucleotide sequences but before the pseudo gVIII
35 sequences. Therefore, the pseudo gVIII will never be

expressed on the phage surface under these circumstances. Instead, only soluble peptides will be produced. Expression in a non-suppressor strain can be advantageously utilized when one wishes to produce large populations of soluble peptides. Stop codons other than amber, such as opal and ochre, or molecular switches, such as inducible repressor elements, can also be used to unlink peptide expression from surface expression. Additional controls exist as well and are described below.

10 The vector used for anti-sense strand oligonucleotide portions, M13IX22, (Figure 3A), contains the expression elements for the peptide fusion proteins. Upstream and in frame with the Sac I and Eco RI sites in this vector is a leader sequence for surface expression. A ribosome binding
15 site and Lac Z promoter/operator elements are present for transcription and translation of the peptide fusion proteins.

Both vectors contain a pair of Fok I restriction enzyme sites (Figure 3 A and B) for joining together two precursor oligonucleotide portions and their vector
20 sequences. One site is located at the ends of each precursor oligonucleotide which is to be joined. The second Fok I site within the vectors is located at the end of the vector sequences which are to be joined. The 5' overhang of this second Fok I site has been altered to
25 encode a sequence which is not found in the overhangs produced at the first Fok I site within the oligonucleotide portions. The two sites allow the cleavage of each circular vector into two portions and subsequent ligation
30 of essential components within each vector into a single circular vector where the two oligonucleotide precursor portions form a contiguous sequence (Figure 3C). Non-compatible overhangs produced at the two Fok I sites allows optimal conditions to be selected for performing
35 concatamerization or circularization reactions for joining

the two vector portions. Such selection of conditions can be used to govern the reaction order and therefore increase the efficiency of joining.

Fok I is a restriction enzyme whose recognition
5 sequence is distal to the point of cleavage. Distal
placement of the recognition sequence in its location to
the cleavage point is important since if the two were
superimposed within the oligonucleotide portions to be
combined, it would lead to an invariant codon sequence at
10 the juncture. To alleviate the formation of invariant
codons at the juncture, Fok I recognition sequences can be
placed outside of the random codon sequence and still be
used to restrict within the random sequence. Subsequent
annealing of the single-strand overhangs produced by Fok I
15 and ligation of the two oligonucleotide precursor portions
allows the juncture to be formed. A variety of restriction
enzymes restrict DNA by this mechanism and can be used
instead of Fok I to join precursor oligonucleotides without
creating invariant codon sequences. Such enzymes include,
20 for example, Alw I, Bbu I, Bsp MI, Hga I, Hph I, Mbo II,
Mnl I, Ple I and Sfa NI. One skilled in the art knows how
to substitute Fok I recognition sequences for alternative
enzyme recognition sequences such as those above, and use
the appropriate enzyme for joining precursor
25 oligonucleotide portions.

Although the sequences of the precursor
oligonucleotides are random and will invariably have
oligonucleotides within the two precursor populations whose
sequences are sufficiently complementary to anneal after
30 cleavage, the efficiency of annealing can be increased by
insuring that the single-strand overhangs within one
precursor population will have a complementary sequence
within the second precursor population. This can be
accomplished by synthesizing a non-degenerate series of
35 known sequences at the Fok I cleavage site coding for each

of the twenty amino acids. Since the Fok I cleavage site contains a four base overhang, forty different sequences are needed to randomly encode all twenty amino acids. For example, if two precursor populations of ten codons in length are to be combined, then after the ninth codon position is synthesized, the mixed population of supports are divided into forty reaction vessels for each of the populations and complementary sequences for each of the corresponding reaction vessels between populations are independently synthesized. The sequences are shown in Tables III and VI of Example I where the oligonucleotides on columns 1R through 40R form complementary overhangs with the oligonucleotides on the corresponding columns 1L through 40L once cleaved. The degenerate X positions in Table VI are necessary to maintain the reading frame once the precursor oligonucleotide portions are joined. However, use of restriction enzymes which produce a blunt end, such as Mnl I can be alternatively used in place of Fok I to alleviate the degeneracy introduced in maintaining the reading frame.

The last feature exhibited by each of the vectors is an amber stop codon located in an essential coding sequence within the vector portion lost during combining (Figure 3C). The amber stop codon is present to select for viable phage produced from only the proper combination of precursor oligonucleotides and their vector sequences into a single vector species. Other non-sense mutations or selectable markers can work as well.

The combining step randomly brings together different precursor oligonucleotides within the two populations into a single vector (Figure 3C; M13IX). The vector sequences donated from each independent vector, M13IX22 and M13IX42, are necessary for production of viable phage. Also, since the expression elements are contained in M13IX22 and the gVIII sequences are contained in M13IX42, expression of

functional gVIII-peptide fusion proteins cannot be accomplished until the sequences are linked as shown in M13IX.

The combining step is performed by restricting each
5 population of vectors containing randomized
oligonucleotides with Fok I, mixing and ligating (Figure
3C). Any vectors generated which contain an amber stop
codon will not produce viable phage when introduced into a
non-suppressor strain (Figure 3D). Therefore, only the
10 sequences which do not contain an amber stop codon will
make up the final population of vectors contained in the
library. These vector sequences are the sequences required
for surface expression of randomized peptides. By
analogous methodology, more than two vector portions can be
15 combined into a single vector which expresses random
peptides.

The invention provides for a method of selecting
peptides capable of being bound by a ligand binding protein
from a population of random peptides by (a) operationally
20 linking a diverse population of first oligonucleotides
having a desirable bias of random codon sequences to a
first vector; (b) operationally linking a diverse
population of second oligonucleotides having a desirable
bias of random codon sequences to a second vector; (c)
25 combining the vector products of steps (a) and (b) under
conditions where said populations of first and second
oligonucleotides are joined together into a population of
combined vectors; (d) introducing said population of
combined vectors into a compatible host under conditions
30 sufficient for expressing said population of random
peptides; and (e) determining the peptides which bind to
said binding protein. The invention also provides for
determining the encoding nucleic acid sequence of such
peptides as well.

Surface expression of the random peptide library is performed in an amber suppressor strain. As described above, the amber stop codon between the random codon sequence and the gVIII sequence unlinks the two components in a non-suppressor strain. Isolating the phage produced from the non-suppressor strain and infecting a suppressor strain will link the random codon sequences to the gVIII sequence during expression (Figure 3E). Culturing the suppressor strain after infection allows the expression of all peptide species within the library as gVIII-peptide fusion proteins. Alternatively, the DNA can be isolated from the non-suppressor strain and then introduced into a suppressor strain to accomplish the same effect.

The level of expression of gVIII-peptide fusion proteins can additionally be controlled at the transcriptional level. The gVIII-peptide fusion proteins are under the inducible control of the Lac Z promoter/operator system. Other inducible promoters can work as well and are known by one skilled in the art. For high levels of surface expression, the suppressor library is cultured in an inducer of the Lac Z promoter such as isopropylthio- β -galactoside (IPTG). Inducible control is beneficial because biological selection against non-functional gVIII-peptide fusion proteins can be minimized by culturing the library under non-expressing conditions. Expression can then be induced only at the time of screening to ensure that the entire population of oligonucleotides within the library are accurately represented on the phage surface. Also this can be used to control the valency of the peptide on the phage surface.

The surface expression library is screened for specific peptides which bind ligand binding proteins by standard affinity isolation procedures. Such methods include, for example, panning, affinity chromatography and solid phase blotting procedures. Panning as described by

Parmley and Smith, Gene 73:305-318 (1988), which is incorporated herein by reference, is preferred because high titers of phage can be screened easily, quickly and in small volumes. Furthermore, this procedure can select
5 minor peptide species within the population, which otherwise would have been undetectable, and amplified to substantially homogenous populations. The selected peptide sequences can be determined by sequencing the nucleic acid encoding such peptides after amplification of the phage
10 population.

The invention provides a plurality of procaryotic cells containing a diverse population of oligonucleotides having a desirable bias of random codon sequences that are operationally linked to expression sequences. The
15 invention provides for methods of constructing such populations of cells as well.

Random oligonucleotides synthesized by any of the methods described previously can also be expressed on the surface of filamentous bacteriophage, such as M13, for
20 example, without the joining together of precursor oligonucleotides. A vector such as that shown in Figure 4, M13IX30, can be used. This vector exhibits all the functional features of the combined vector shown in Figure 3C for surface expression of gVIII-peptide fusion proteins.
25 The complete nucleotide sequence for M13IX30 (SEQ ID NO: 3) is shown in Figure 7.

M13IX30 contains a wild type gVIII for phage viability and a pseudo gVIII sequence for peptide fusions. The vector also contains in frame restriction sites for cloning
30 random peptides. The cloning sites in this vector are Xho I, Stu I and Spe I. Oligonucleotides should therefore be synthesized with the appropriate complementary ends for annealing and ligation or insertional mutagenesis. Alternatively, the appropriate termini can be generated by

PCR technology. Between the restriction sites and the pseudo gVIII sequence is an in-frame amber stop codon, again, ensuring complete viability of phage in constructing and manipulating the library. Expression and screening is performed as described above for the surface expression library of oligonucleotides generated from precursor portions.

Thus, the invention provides a method of selecting peptides capable of being bound by a ligand binding protein from a population of random peptides by (a) operationally linking a diverse population of oligonucleotides having a desirable bias of random codon sequences to expression elements; (b) introducing said population of vectors into a compatible host under conditions sufficient for expressing said population of random peptides; and (c) determining the peptides which bind to said binding protein. Also provided is a method for determining the encoding nucleic acid sequence of such selected peptides.

The following examples are intended to illustrate, but not limit the invention.

EXAMPLE I

Isolation and Characterization of Peptide Ligands Generated From Right and Left Half Random Oligonucleotides

This example shows the synthesis of random oligonucleotides and the construction and expression of surface expression libraries of the encoded randomized peptides. The random peptides of this example derive from the mixing and joining together of two random oligonucleotides. Also demonstrated is the isolation and characterization of peptide ligands and their corresponding nucleotide sequence for specific binding proteins.

Synthesis of Random Oligonucleotides

The synthesis of two randomized oligonucleotides which correspond to smaller portions of a larger randomized oligonucleotide is shown below. Each of the two smaller portions make up one-half of the larger oligonucleotide. The population of randomized oligonucleotides constituting each half are designated the right and left half. Each population of right and left halves are ten codons in length with twenty random codons at each position. The right half corresponds to the sense sequence of the randomized oligonucleotides and encode the carboxy terminal half of the expressed peptides. The left half corresponds to the anti-sense sequence of the randomized oligonucleotides and encode the amino terminal half of the expressed peptides. The right and left halves of the randomized oligonucleotide populations are cloned into separate vector species and then mixed and joined so that the right and left halves come together in random combination to produce a single expression vector species which contains a population of randomized oligonucleotides twenty codons in length. Electroporation of the vector population into an appropriate host produces filamentous phage which express the random peptides on their surface.

The reaction vessels for oligonucleotide synthesis were obtained from the manufacturer of the automated synthesizer (Millipore, Burlington, MA; supplier of MilliGen/Biosearch Cyclone Plus Synthesizer). The vessels were supplied as packages containing empty reaction columns (1 μ mole), frits, crimps and plugs (MilliGen/Biosearch catalog # GEN 860458). Derivatized and underivatized control pore glass, phosphoramidite nucleotides, and synthesis reagents were also obtained from MilliGen/Biosearch. Crimper and decrimp r tools were obtained from Fisher Scientific Co., Pittsburgh, PA (Catalog numbers 06-406-20 and 06-406-25A, respectively).

Ten reaction columns were used for right half synthesis of random oligonucleotides ten codons in length. The oligonucleotides have 5 monomers at their 3' end of the sequence 5'GAGCT3' and 8 monomers at their 5' end of the sequence 5'AATTCCAT3'. The synthesizer was fitted with a column derivatized with a thymine nucleotide (T-column, MilliGen/Biosearch # 0615.50) and was programmed to synthesize the sequences shown in Table I for each of ten columns in independent reaction sets. The sequence of the last three monomers (from right to left since synthesis proceeds 3' to 5') encode the indicated amino acids:

Table I

	<u>Column</u>	<u>Sequence (5' to 3')</u>	<u>Amino Acids</u>
15	column 1R	(T/G)TTGAGCT	Phe and Val
	column 2R	(T/C)CTGAGCT	Ser and Pro
	column 3R	(T/C)ATGAGCT	Tyr and His
	column 4R	(T/C)GTGAGCT	Cys and Arg
	column 5R	(C/A)TGGAGCT	Leu and Met
20	column 6R	(C/G)AGGAGCT	Gln and Glu
	column 7R	(A/G)CTGAGCT	Thr and Ala
	column 8R	(A/G)ATGAGCT	Asn and Asp
	column 9R	(T/G)GGGAGCT	Trp and Gly
	column 1R	A(T/A)AGAGCT	Ile and Cys

where the two monomers in parentheses denote a single monomer position within the codon and indicate that an equal mixture of each monomer was added to the reaction for coupling. The monomer coupling reactions for each of the columns were performed as recommended by the manufacturer (amidite version S1.06, # 8400-050990, scale 1 μ M). After the last coupling reaction, the columns were washed with acetonitrile and lyophilized to dryness.

Following synthesis, the plugs were removed from each

column using a decrimper and the reaction products were poured into a single weigh boat. Initially the bead mass increases, due to the weight of the monomers, however, at later rounds of synthesis material is lost. In either case, the material was equalized with underivatized control pore glass and mixed thoroughly to obtain a random distribution of all twenty codon species. The reaction products were then aliquotted into 10 new reaction columns by removing 25 mg of material at a time and placing it into separate reaction columns. Alternatively, the reaction products can be aliquotted by suspending the beads in a liquid that is dense enough for the beads to remain dispersed, preferably a liquid that is equal in density to the beads, and then aliquoting equal volumes of the suspension into separate reaction columns. The lip on the inside of the columns where the frits rest was cleared of material using vacuum suction with a syringe and 25 G needle. New frits were placed onto the lips, the plugs were fitted into the columns and were crimped into place using a crimper.

Synthesis of the second codon position was achieved using the above 10 columns containing the random mixture of reaction products from the first codon synthesis. The monomer coupling reactions for the second codon position are shown in Table II. An A in the first position means that any monomer can be programmed into the synthesizer. At that position, the first monomer position is not coupled by the synthesizer since the software assumes that the monomer is already attached to the column. An A also denotes that the columns from the previous codon synthesis should be placed on the synthesizer for use in the present synthesis round. Reactions were again sequentially repeated for each column as shown in Table II and the reaction products washed and dried as described above.

Table II

	<u>Column</u>	<u>Sequence (5' to 3')</u>	<u>Amino Acids</u>
	column 1R	(T/G)TTA	Phe and Val
5	column 2R	(T/C)CTA	Ser and Pro
	column 3R	(T/C)ATA	Tyr and His
	column 4R	(T/C)GTA	Cys and Arg
	column 5R	(C/A)TGA	Leu and Met
	column 6R	(C/G)AGA	Gln and Glu
10	column 7R	(A/G)CTA	Thr and Ala
	column 8R	(A/G)ATA	Asn and Asp
	column 9R	(T/G)GGA	Trp and Gly
	column 10R	A(T/A)AA	Ile and Cys

Randomization of the second codon position was achieved by removing the reaction products from each of the columns and thoroughly mixing the material. The material was again divided into new reaction columns and prepared for monomer coupling reactions as described above.

Random synthesis of the next seven codons (positions 3 through 9) proceeded identically to the cycle described above for the second codon position and again used the monomer sequences of Table II. Each of the newly repacked columns containing the random mixture of reaction products from synthesis of the previous codon position was used for the synthesis of the subsequent codon position. After synthesis of the codon at position nine and mixing of the reaction products, the material was divided and repacked into 40 different columns and the monomer sequences shown in Table III were coupled to each of the 40 columns in independent reactions. The oligonucleotides from each of the 40 columns were mixed once more and cleaved from the control pore glass as recommended by the manufacturer.

Table III

	<u>Column</u>	<u>Sequence (5' to 3')</u>
5	column 1R	AATTCCTTTTA
	column 2R	AATTCCTGTTA
	column 3R	AATTCGTTTA
	column 4R	AATTCGGTTA
	column 5R	AATTCCTTCTA
10	column 6R	AATTCCTCCTA
	column 7R	AATTCGTCTA
	column 8R	AATTCGCCTA
	column 9R	AATTCCTTATA
	column 10R	AATTCCTCATA
15	column 11R	AATTCGTATA
	column 12R	AATTCGCATA
	column 13R	AATTCCTTGTA
	column 14R	AATTCCTCGTA
	column 15R	AATTCGTGTA
20	column 16R	AATTCGCGTA
	column 17R	AATTCCTCTGA
	column 18R	AATTCCTATGA
	column 19R	AATTCGCTGA
	column 20R	AATTCGATGA
25	column 21R	AATTCCTCAGA
	column 22R	AATTCCTGAGA
	column 23R	AATTCGCAGA
	column 24R	AATTCGGAGA
	column 25R	AATTCCTACTA
30	column 26R	AATTCCTGCTA
	column 27R	AATTCGACTA
	column 28R	AATTCGGCTA
	column 29R	AATTCCTAATA
	column 30R	AATTCCTGATA
35	column 31R	AATTCGAATA
	column 32R	AATTCGGATA
	column 33R	AATTCCTGGA

	column 34R	<u>AATTCTGGGA</u>
	column 35R	<u>AATTCGTGGA</u>
	column 36R	<u>AATTCGGGGA</u>
	column 37R	<u>AATTCTATAA</u>
5	column 38R	<u>AATTCTAAAA</u>
	column 39R	<u>AATTCGATAA</u>
	column 40R	<u>AATTCGAAAA</u>

Left half synthesis of random oligonucleotides proceeded similarly to the right half synthesis. This half of the oligonucleotide corresponds to the anti-sense sequence of the encoded randomized peptides. Thus, the complementary sequence of the codons in Tables I through III are synthesized. The left half oligonucleotides also have 5 monomers at their 3' end of the sequence 5'GAGCT3' and 8 monomers at their 5' end of the sequence 5'AATTCCAT3'. The rounds of synthesis, washing, drying, mixing, and dividing are as described above.

For the first codon position, the synthesizer was fitted with a T-column and programmed to synthesize the sequences shown in Table IV for each of ten columns in independent reaction sets. As with right half synthesis, the sequence of the last three monomers (from right to left) encode the indicated amino acids:

Table IV

	<u>Column</u>	<u>Sequence (5' to 3')</u>	<u>Amino Acids</u>
5	column 1L	AA(A/C)GAGCT	Phe and Val
	column 2L	AG(A/G)GAGCT	Ser and Pro
	column 3L	AT(A/G)GAGCT	Tyr and His
	column 4L	AC(A/G)GAGCT	Cys and Arg
	column 5L	CA(G/T)GAGCT	Leu and Met
10	column 6L	CT(G/C)GAGCT	Gln and Glu
	column 7L	AG(T/C)GAGCT	Thr and Ala
	column 8L	AT(T/C)GAGCT	Asn and Asp
	column 9L	CC(A/C)GAGCT	Trp and Gly
	column 10L	T(A/T)TGAGCT	Ile and Cys

Following washing and drying, the plugs for each column
 15 were removed, mixed and aliquotted into ten new reaction
 columns as described above. Synthesis of the second codon
 position was achieved using these ten columns containing
 the random mixture of reaction products from the first
 codon synthesis. The monomer coupling reactions for the
 20 second codon position are shown in Table V.

Table V

	<u>Column</u>	<u>Sequence (5' to 3')</u>	<u>Amino Acids</u>
25	column 1L	AA(A/C) <u>A</u>	Phe and Val
	column 2L	AG(A/G) <u>A</u>	Ser and Pro
	column 3L	AT(A/G) <u>A</u>	Tyr and His
	column 4L	AC(A/G) <u>A</u>	Cys and Arg
	column 5L	CA(G/T) <u>A</u>	Leu and Met
30	column 6L	CT(G/C) <u>A</u>	Gln and Glu
	column 7L	AG(T/C) <u>A</u>	Thr and Ala
	column 8L	AT(T/C) <u>A</u>	Asn and Asp
	column 9L	CC(A/C) <u>A</u>	Trp and Gly
	column 10L	T(A/T)T <u>A</u>	Ile and Cys

Again, randomization of the second codon position was achieved by removing the reaction products from each of the columns and thoroughly mixing the beads. The beads were repacked into ten new reaction columns.

- 5 Random synthesis of the next seven codon positions proceeded identically to the cycle described above for the second codon position and again used the monomer sequences of Table V. After synthesis of the codon at position nine and mixing of the reaction products, the material was
 10 divided and repacked into 40 different columns and the monomer sequences shown in Table VI were coupled to each of the 40 columns in independent reactions.

Table VI

15	<u>Column</u>	<u>Sequence (5' to 3')</u>
	column 1L	AATTCCATAAAAXXA
	column 2L	AATTCCATAAACXXA
	column 3L	AATTCCATAACAXXA
	column 4L	AATTCCATAACCXXA
20	column 5L	AATTCCATAGAAXXA
	column 6L	AATTCCATAGACXXA
	column 7L	AATTCCATAGGAXXA
	column 8L	AATTCCATAGGCXXA
	column 9L	AATTCCATATAAXXA
25	column 10L	AATTCCATATACXXA
	column 11L	AATTCCATATGAXXA
	column 12L	AATTCCATATGCXXA
	column 13L	AATTCCATACAAXXA
	column 14L	AATTCCATACACXXA
30	column 15L	AATTCCATACGAXXA
	column 16L	AATTCCATACGCXXA
	column 17L	AATTCCATCAGAXXA
	column 18L	AATTCCATCAGCXXA
	column 19L	AATTCCATCATAXXA
35	column 20L	AATTCCATCATCXXA

36

	column 21L	<u>AATTCCATCTGAXXA</u>
	column 22L	<u>AATTCCATCTGCXXA</u>
	column 23L	<u>AATTCCATCTCAXXA</u>
	column 24L	<u>AATTCCATCTCCXXA</u>
5	column 25L	<u>AATTCCATAGTAXXA</u>
	column 26L	<u>AATTCCATAGTCXXA</u>
	column 27L	<u>AATTCCATAGCAXXA</u>
	column 28L	<u>AATTCCATAGCCXXA</u>
	column 29L	<u>AATTCCATATTAXXA</u>
10	column 30L	<u>AATTCCATATTCXXA</u>
	column 31L	<u>AATTCCATATCAXXA</u>
	column 32L	<u>AATTCCATATCCXXA</u>
	column 33L	<u>AATTCCATCCAAXXA</u>
	column 34L	<u>AATTCCATCCACXXA</u>
15	column 35L	<u>AATTCCATCCCAXXA</u>
	column 36L	<u>AATTCCATCCCCXXA</u>
	column 37L	<u>AATTCCATTATAXXA</u>
	column 38L	<u>AATTCCATTATCXXA</u>
	column 39L	<u>AATTCCATTTTAXXA</u>
20	column 40L	<u>AATTCCATTTTCXXA</u>

The first two monomers denoted by an "X" represent an equal mixture of all four nucleotides at that position. This is necessary to retain a relatively unbiased codon sequence at the junction between right and left half oligonucleotides.

25 The above right and left half random oligonucleotides were cleaved and purified from the supports and used in constructing the surface expression libraries below.

Vector Construction

Two M13-based vectors, M13IX42 (SEQ ID NO: 1) and

30 M13IX22 (SEQ ID NO: 2), were constructed for the cloning and propagation of right and left half populations of random oligonucleotides, respectively. The vectors were specially constructed to facilitate the random joining and subsequent expression of right and left half

oligonucleotide populations. Each vector within the population contains on right and one left half oligonucleotide from the population joined together to form a single contiguous oligonucleotide with random codons which is twenty-two codons in length. The resultant population of vectors are used to construct a surface expression library.

M13IX42, or the right-half vector, was constructed to harbor the right half populations of randomized oligonucleotides. M13mp18 (Pharmacia, Piscataway, NJ) was the starting vector. This vector was genetically modified to contain, in addition to the encoded wild type M13 gene VIII already present in the vector: (1) a pseudo-wild type M13 gene VIII sequence with a stop codon (amber) placed between it and an Eco RI-Sac I cloning site for randomized oligonucleotides; (2) a pair of Fok I sites to be used for joining with M13IX22, the left-half vector; (3) a second amber stop codon placed on the opposite side of the vector than the portion being combined with the left-half vector; and (4) various other mutations to remove redundant restriction sites and the amino terminal portion of Lac Z.

The pseudo-wild type M13 gene VIII was used for surface expression of random peptides. The pseudo-wild type gene encodes the identical amino acid sequence as that of the wild type gene; however, the nucleotide sequence has been altered so that only 63% identity exists between this gene and the encoded wild type gene VIII. Modification of the gene VIII nucleotide sequence used for surface expression reduces the possibility of homologous recombination with the wild type gene VIII contained on the same vector. Additionally, the wild type M13 gene VIII was retained in the vector system to ensure that at least some functional, non-fusion coat protein would be produced. The inclusion of wild type gene VIII therefore reduces the possibility of non-viable phage production from the random

peptide fusion genes.

The pseudo-wild type gene VIII was constructed by chemically synthesizing a series of oligonucleotides which encode both strands of the gene. The oligonucleotides are presented in Table VII (SEQ ID NOS: 7 through 16).

TABLE VII

Pseudo-Wild Type Gene VIII Oligonucleotide Series

	<u>Top Strand</u> <u>Oligonucleotides</u>	<u>Sequence (5' to 3')</u>
10	VIII 03	GATCC TAG GCT GAA GGC GAT GAC CCT GCT AAG GCT GC
	VIII 04	A TTC AAT AGT TTA CAG GCA AGT GCT ACT GAG TAC A
	VIII 05	TT GGC TAC GCT TGG GCT ATG GTA GTA GTT ATA GTT
15	VIII 06	GGT GCT ACC ATA GGG ATT AAA TTA TTC AAA AAG TT
	VIII 07	T ACG AGC AAG GCT TCT TA
20	<u>Bottom Strand</u> <u>Oligonucleotides</u>	
	VIII 08	AGC TTA AGA AGC CTT GCT CGT AAA CTT TTT GAA TAA TTT
	VIII 09	AAT CCC TAT GGT AGC ACC AAC TAT AAC TAC TAC CAT
25	VIII 10	AGC CCA AGC GTA GCC AAT GTA CTC AGT AGC ACT TG
	VIII 11	C CTG TAA ACT ATT GAA TGC AGC CTT AGC AGG GTC
	VIII 12	ATC GCC TTC AGC CTA G

30 Except for the terminal oligonucleotides VIII 03 (SEQ

ID NO: 7) and VIII 08 (SEQ ID NO: 12), the above oligonucleotides (oligonucleotides VIII 04-VIII 07 and 09-12 (SEQ ID NOS: 8 through 11 and 13 through 16)) were mixed at 200 ng each in 10 μ l final volume and phosphorylated with T4 polynucleotide Kinase (Pharmacia, Piscataway, NJ) with 1 mM ATP at 37°C for 1 hour. The reaction was stopped at 65°C for 5 minutes. Terminal oligonucleotides were added to the mixture and annealed into double-stranded form by heating to 65°C for 5 minutes, followed by cooling to room temperature over a period of 30 minutes. The annealed oligonucleotides were ligated together with 1.0 U of T4 DNA ligase (BRL). The annealed and ligated oligonucleotides yield a double-stranded DNA flanked by a Bam HI site at its 5' end and by a Hind III site at its 3' end. A translational stop codon (amber) immediately follows the Bam HI site. The gene VIII sequence begins with the codon GAA (Glu) two codons 3' to the stop codon. The double-stranded insert was phosphorylated using T4 DNA Kinase (Pharmacia, Piscataway, NJ) and ATP (10 mM Tris-HCl, pH 7.5, 10 mM MgCl₂) and cloned in frame with the Eco RI and Sac I sites within the M13 polylinker. To do so, M13mp18 was digested with Bam HI (New England Biolabs, Beverly, MA) and Hind III (New England Biolabs) and combined at a molar ratio of 1:10 with the double-stranded insert. The ligations were performed at 16°C overnight in 1X ligase buffer (50 mM Tris-HCl, pH 7.8, 10 mM MgCl₂, 20 mM DTT, 1 mM ATP, 50 μ g/ml BSA) containing 1.0 U of T4 DNA ligase (New England Biolabs). The ligation mixture was transformed into a host and screened for positive clones using standard procedures in the art.

Several mutations were generated within the right-half vector to yield functional M13IX42. The mutations were generated using the method of Kunkel et al., Meth. Enzymol. 154:367-382 (1987), which is incorporated herein by reference, for site-directed mutagenesis. The reagents, strains and protocols were obtained from a Bio Rad

Mutagenesis kit (Bio Rad, Richmond, CA) and mutagenesis was performed as recommended by the manufacturer.

A Fok I site used for joining the right and left halves was generated 8 nucleotides 5' to the unique Eco RI site using the oligonucleotide 5'-CTCGAATTCGTACATCCT GGTTCATAGC-3' (SEQ ID NO: 17). The second Fok I site retained in the vector is naturally encoded at position 3547; however, the sequence within the overhang was changed to encode CTTC. Two Fok I sites were removed from the vector at positions 239 and 7244 of M13mp18 as well as the Hind III site at the end of the pseudo gene VIII sequence using the mutant oligonucleotides 5'-CATTTTTGCAGATGGCTTAGA-3' (SEQ ID NO: 18) and 5'-TAGCATTAACGTCCAATA-3' (SEQ ID NO: 19), respectively. New Hind III and Mlu I sites were also introduced at position 3919 and 3951 of M13IX42. The oligonucleotides used for this mutagenesis had the sequences 5'-ATATATTTTAGTAAGCTTCATCTTCT-3' (SEQ ID NO: 20) and 5'-GACAAAGAACGCGTGAAACTTT-3' (SEQ ID NO: 21), respectively. The amino terminal portion of Lac Z was deleted by oligonucleotide-directed mutagenesis using the mutant oligonucleotide 5'-GCGGGCCTCTTCGCTATTGCTTAAGAAGCCTTGCT-3' (SEQ ID NO: 22). This deletion also removed a third M13mp18 derived Fok I site. The distance between the Eco RI and Sac I sites was increased to ensure complete double digestion by inserting a spacer sequence. The spacer sequence was inserted using the oligonucleotide 5'-TTCAGCCTAGGATCCGCCGAGCTCTCCTACCTGCGAATTCGTACATCC-3' (SEQ ID NO: 23). Finally, an amber stop codon was placed at position 4492 using the mutant oligonucleotide 5'-TGGATTATACTTCTA AATAATGGA-3' (SEQ ID NO: 24). The amber stop codon is used as a biological selection to ensure the proper recombination of vector sequences to bring together right and left halves of the randomized oligonucleotides. In constructing the above mutations, all changes made in a M13 coding region were performed such that the amino acid

sequence remained unaltered. It should be noted that several mutations within M13mp18 were found which differed from the published sequence. Where known, these sequence differences are recorded herein as found and therefore may
5 not correspond exactly to the published sequence of M13mp18.

The sequence of the resultant vector, M13IX42, is shown in Figure 5 (SEQ ID NO: 1). Figure 3A also shows M13IX42 where each of the elements necessary for producing
10 a surface expression library between right and left half randomized oligonucleotides is marked. The sequence between the two Fok I sites shown by the arrow is the portion of M13IX42 which is to be combined with a portion of the left-half vector to produce random oligonucleotides
15 as fusion proteins of gene VIII.

M13IX22, or the left-half vector, was constructed to harbor the left half populations of randomized oligonucleotides. This vector was constructed from M13mp19 (Pharmacia, Piscataway, NJ) and contains: (1) Two Fok I
20 sites for mixing with M13IX42 to bring together the left and right halves of the randomized oligonucleotides; (2) sequences necessary for expression such as a promoter and signal sequence and translation initiation signals; (3) an Eco RI-Sac I cloning site for the randomized
25 oligonucleotides; and (4) an amber stop codon for biological selection in bringing together right and left half oligonucleotides.

Of the two Fok I sites used for mixing M13IX22 with M13IX42, one is naturally encoded in M13mp18 and M13mp19
30 (at position 3547). As with M13IX42, the overhang within this naturally occurring Fok I site was changed to CTTC. The other Fok I site was introduced after construction of the translation initiation signals by site-directed mutagenesis using the oligonucleotide 5'-

TAACACTCATTCCGGATGGAATTCTGGAGTCTGGGT-3' (SEQ ID NO: 25).

The translation initiation signals were constructed by annealing of overlapping oligonucleotides as described above to produce a double-stranded insert containing a 5' Eco RI site and a 3' Hind III site. The overlapping oligonucleotides are shown in Table VIII (SEQ ID NOS: 26 through 34) and were ligated as a double-stranded insert between the Eco RI and Hind III sites of M13mp18 as described for the pseudo gene VIII insert. The ribosome binding site (AGGAGAC) is located in oligonucleotide 015 (SEQ ID NO: 26) and the translation initiation codon (ATG) is the first three nucleotides of oligonucleotide 016 (SEQ ID NO: 27).

TABLE VIII

Oligonucleotide Series for Construction of
Translation Signals in M13IX22

<u>Oligonucleotide</u>	<u>Sequence (5' to 3')</u>
015	AATT C GCC AAG GAG ACA GTC AT
016	AATG AAA TAC CTA TTG CCT ACG GCA
017	ATTA CTC GCT GCC CAA CCA GCC ATG
018	GACC CAG ACT CCA GATATC CAA CAG
019	GAA TGA GTG TTA AT
020	TCT AGA ACG CGT C
021	ACGT G ACG CGT TCT AGA AT TAA
022	CACTCA TTC CTG T
023	TG GAT ATC TGG AGT CTG GGT CAT
024	CAC GAG CTC GGC CAT G
025	GC TGG TTG GGC AGC GAG TAA TAA
026	CAA TCC AGC GGC TGC C
027	GT AGG CAA TAG GTA TTT CAT TAT
028	GAC TGT CCT TGG CG

Oligonucleotide 017 (SEQ ID NO: 27) contained a Sac I restriction site 67 nucleotides downstream from the ATG codon. The naturally occurring Eco RI site was removed and a new site introduced 25 nucleotides downstream from the
5 Sac I. Oligonucleotides 5'-TGACTGTCTCCTTGGCGTGTGAAATTGTTA-3' (SEQ ID NO: 35) and 5'-TAACACTCATTCCGGATGGAATTCTGGAGTCTGGGT-3' (SEQ ID NO: 36) were used to generate each of the mutations, respectively. An amber stop codon was also introduced at position 3263 of M13mp18 using the
10 oligonucleotide 5'-CAATTTTATCCTAAATCTTACCAAC-3' (SEQ ID NO: 37).

In addition to the above mutations, a variety of other modifications were made to remove certain sequences and redundant restriction sites. The LAC Z ribosome binding
15 site was removed when the original Eco RI site in M13mp18 was mutated. Also, the Fok I sites at positions 239, 6361 and 7244 of M13mp18 were likewise removed with mutant oligonucleotides 5'-CATTTTTGCAGATGGCTTAGA-3' (SEQ ID NO: 38), 5'-CGAAAGGGGGGTGTGCTGCAA-3' (SEQ ID NO: 39) and 5'-
20 TAGCATTAACGTCCAATA-3' (SEQ ID NO: 40), respectively. Again, mutations within the coding region did not alter the amino acid sequence.

The resultant vector, M13IX22, is 7320 base pairs in length, the sequence of which is shown in Figure 6 (SEQ ID
25 NO: 2). The Sac I and Eco RI cloning sites are at positions 6290 and 6314, respectively. Figure 3A also shows M13IX22 where each of the elements necessary for producing a surface expression library between right and left half randomized oligonucleotides is marked.

30 Library Construction

Each population of right and left half randomized oligonucleotides from columns 1R through 40R and columns 1L through 40L are cloned separately into M13IX42 and M13IX22,

respectively, to create sublibraries of right and left half randomized oligonucleotides. Therefore, a total of eighty sublibraries are generated. Separately maintaining each population of randomized oligonucleotides until the final
5 screening step is performed to ensure maximum efficiency of annealing of right and left half oligonucleotides. The greater efficiency increases the total number of randomized oligonucleotides which can be obtained. Alternatively, one can combine all forty populations of right half
10 oligonucleotides (columns 1R-40R) into one population and of left half oligonucleotides (columns 1L-40L) into a second population to generate just one sublibrary for each.

For the generation of sublibraries, each of the above populations of randomized oligonucleotides are cloned
15 separately into the appropriate vector. The right half oligonucleotides are cloned into M13IX42 to generate sublibraries M13IX42.1R through M13IX42.40R. The left half oligonucleotides are similarly cloned into M13IX22 to generate sublibraries M13IX22.1L through M13IX22.40L. Each
20 vector contains unique Eco RI and Sac I restriction enzyme sites which produce 5' and 3' single-stranded overhangs, respectively, when digested. The single strand overhangs are used for the annealing and ligation of the complementary single-stranded random oligonucleotides.

25 The randomized oligonucleotide populations are cloned between the Eco RI and Sac I sites by sequential digestion and ligation steps. Each vector is treated with an excess of Eco RI (New England Biolabs) at 37°C for 2 hours followed by addition of 4-24 units of calf intestinal
30 alkaline phosphatase (Boehringer Mannheim, Indianapolis, IN). Reactions are stopped by phenol/chloroform extraction and ethanol precipitation. The pellets are resuspended in an appropriate amount of distilled or deionized water (dH₂O). About 10 pmol of vector is mixed with a 5000-fold
35 molar excess of each population of randomized

oligonucleotides in 10 μ l of 1X ligase buffer (50 mM Tris-HCl, pH 7.8, 10 mM $MgCl_2$, 20 mM DTT, 1 mM ATP, 50 μ g/ml BSA) containing 1.0 U of T4 DNA ligase (BRL, Gaithersburg, MD). The ligation is incubated at 16°C for 16 hours. Reactions
5 are stopped by heating at 75°C for 15 minutes and the DNA is digested with an excess of Sac I (New England Biolabs) for 2 hours. Sac I is inactivated by heating at 75°C for 15 minutes and the volume of the reaction mixture is adjusted to 300 μ l with an appropriate amount of 10X ligase
10 buffer and dH_2O . One unit of T4 DNA ligase (BRL) is added and the mixture is incubated overnight at 16°C. The DNA is ethanol precipitated and resuspended in TE (10 mM Tris-HCl, pH 8.0, 1 mM EDTA). DNA from each ligation is electroporated into XL1 BlueTM cells (Stratagene, La Jolla,
15 CA), as described below, to generate the sublibraries.

E. coli XL1 BlueTM is electroporated as described by Smith et al., Focus 12:38-40 (1990) which is incorporated herein by reference. The cells are prepared by inoculating a fresh colony of XL1s into 5 mls of SOB without magnesium
20 (20 g bacto-tryptone, 5 g bacto-yeast extract, 0.584 g NaCl, 0.186 g KCl, dH_2O to 1,000 mls) and grown with vigorous aeration overnight at 37°C. SOB without magnesium (500 ml) is inoculated at 1:1000 with the overnight culture and grown with vigorous aeration at 37°C until the OD_{550} is
25 0.8 (about 2 to 3 h). The cells are harvested by centrifugation at 5,000 rpm (2,600 x g) in a GS3 rotor (Sorvall, Newtown, CT) at 4°C for 10 minutes, resuspended in 500 ml of ice-cold 10% (v/v) sterile glycerol and centrifuged and resuspended a second time in the same
30 manner. After a third centrifugation, the cells are resuspended in 10% sterile glycerol at a final volume of about 2 ml, such that the OD_{550} of the suspension is 200 to 300. Usually, resuspension is achieved in the 10% glycerol that remains in the bottle after pouring off the supernate.
35 Cells are frozen in 40 μ l aliquots in microcentrifuge tubes using a dry ice-ethanol bath and stored frozen at -70°C.

Frozen cells are electroporated by thawing slowly on ice before use and mixing with about 10 pg to 500 ng of vector per 40 μ l of cell suspension. A 40 μ l aliquot is placed in an 0.1 cm electroporation chamber (Bio-Rad, Richmond, CA) and pulsed once at 0°C using 200 Ω parallel resistor, 25 μ F, 1.88 kV, which gives a pulse length (τ) of ~4 ms. A 10 μ l aliquot of the pulsed cells are diluted into 1 ml SOC (98 mls SOB plus 1 ml of 2 M $MgCl_2$ and 1 ml of 2 M glucose) in a 12- x 75-mm culture tube, and the culture is shaken at 37°C for 1 hour prior to culturing in selective media, (see below).

Each of the eighty sublibraries are cultured using methods known to one skilled in the art. Such methods can be found in Sanbrook et al., Molecular Cloning: A Laboratory Manual, Cold Spring Harbor Laboratory, Cold Spring Harbor, 1989, and in Ausubel et al., Current Protocols in Molecular Biology, John Wiley and Sons, New York, 1989, both of which are incorporated herein by reference. Briefly, the above 1 ml sublibrary cultures were grown up by diluting 50-fold into 2XYT media (16 g tryptone, 10 g yeast extract, 5 g NaCl) and culturing at 37°C for 5-8 hours. The bacteria were pelleted by centrifugation at 10,000 xg. The supernatant containing phage was transferred to a sterile tube and stored at 4°C.

Double strand vector DNA containing right and left half randomized oligonucleotide inserts is isolated from the cell pellet of each sublibrary. Briefly, the pellet is washed in TE (10 mM Tris, pH 8.0, 1 mM EDTA) and recollected by centrifugation at 7,000 rpm for 5' in a Sorval centrifuge (Newtown, CT). Pellets are resuspended in 6 mls of 10% Sucrose, 50 mM Tris, pH 8.0. 3.0 ml of 10 mg/ μ l lysozyme is added and incubated on ice for 20 minutes. 12 mls of 0.2 M NaOH, 1% SDS is added followed by 10 minutes on ice. The suspensions are then incubated on ice for 20 minutes after addition of 7.5 mls of 3 M NaOAc,

pH 4.6. The samples are centrifuged at 15,000 rpm for 15 minutes at 4°C, RNased and extracted with phenol/chloroform, followed by ethanol precipitation. The pellets are resuspended, weighed and an equal weight of
5 CsCl₂ is dissolved into each tube until a density of 1.60 g/ml is achieved. EtBr is added to 600 µg/ml and the double-stranded DNA is isolated by equilibrium centrifugation in a TV-1665 rotor (Sorval) at 50,000 rpm for 6 hours. These DNAs from each right and left half
10 sublibrary are used to generate forty libraries in which the right and left halves of the randomized oligonucleotides have been randomly joined together.

Each of the forty libraries are produced by joining together one right half and one left half sublibrary. The
15 two sublibraries joined together corresponded to the same column number for right and left half random oligonucleotide synthesis. For example, sublibrary M13IX42.1R is joined with M13IX22.1L to produce the surface expression library M13IX.1RL. In the alternative situation
20 where only two sublibraries are generated from the combined populations of all right half synthesis and all left half synthesis, only one surface expression library would be produced.

For the random joining of each right and left half
25 oligonucleotide populations into a single surface expression vector species, the DNAs isolated from each sublibrary are digested an excess of Fok I (New England Biolabs). The reactions are stopped by phenol/chloroform extraction, followed by ethanol precipitation. Pellets are
30 resuspended in dH₂O. Each surface expression library is generated by ligating equal molar amounts (5-10 pmol) of Fok I digested DNA isolated from corresponding right and left half sublibraries in 10 µl of 1X ligase buffer containing 1.0 U of T4 DNA ligase (Bethesda Research
35 Laboratories, Gaithersburg, MD). The ligations proceed

overnight at 16°C and are electroporated into the sup 0 strain MK30-3 (Boehringer Mannheim Biochemical, (BMB), Indianapolis, IN) as previously described for XL1 cells. Because MK30-3 is sup 0, only the vector portions encoding
5 the randomized oligonucleotides which come together will produce viable phage.

Screening of Surface Expression Libraries

Purified phage are prepared from 50 ml liquid cultures of XL1 BlueTM cells (Stratagene) which are infected at a
10 m.o.i. of 10 from the phage stocks stored at 4°C. The cultures are induced with 2 mM IPTG. Supernatants from all cultures are combined and cleared by two centrifugations, and the phage are precipitated by adding 1/7.5 volumes of PEG solution (25% PEG-8000, 2.5 M NaCl), followed by
15 incubation at 4°C overnight. The precipitate is recovered by centrifugation for 90 minutes at 10,000 x g. Phage pellets are resuspended in 25 ml of 0.01 M Tris-HCl, pH 7.6, 1.0 mM EDTA, and 0.1% Sarkosyl and then shaken slowly at room temperature for 30 minutes. The solutions are
20 adjusted to 0.5 M NaCl and to a final concentration of 5% polyethylene glycol. After 2 hours at 4°C, the precipitates containing the phage are recovered by centrifugation for 1 hour at 15,000 X g. The precipitates are resuspended in 10 ml of NET buffer (0.1 M NaCl, 1.0 mM
25 EDTA, and 0.01 M Tris-HCl, pH 7.6), mixed well, and the phage repelleted by centrifugation at 170,000 X g for 3 hours. The phage pellets are subsequently resuspended overnight in 2 ml of NET buffer and subjected to cesium chloride centrifugation for 18 hours at 110,000 X g (3.86
30 g of cesium chloride in 10 ml of buffer). Phage bands are collected, diluted 7-fold with NET buffer, recentrifuged at 170,000 X g for 3 hours, resuspended, and stored at 4°C in 0.3 ml of NET buffer containing 0.1 mM sodium azide.

Ligand binding proteins used for panning on

streptavidin coated dishes are first biotinylated and then absorbed against UV-inactivated blocking phage (see below). The biotinylating reagents are dissolved in dimethylformamide at a ratio of 2.4 mg solid NHS-SS-Biotin
5 (sulfosuccinimidyl 2-(biotinamido)ethyl-1,3'-dithiopropionate; Pierce, Rockford, IL) to 1 ml solvent and used as recommended by the manufacturer. Small-scale reactions are accomplished by mixing 1 μ l dissolved reagent with 43 μ l of 1 mg/ml ligand binding protein diluted in
10 sterile bicarbonate buffer (0.1 M NaHCO_3 , pH 8.6). After 2 hours at 25°C, residual biotinylating reagent is reacted with 500 μ l 1 M ethanolamine (pH adjusted to 9 with HCl) for an additional 2 hours. The entire sample is diluted with 1 ml TBS containing 1 mg/ml BSA, concentrated to about
15 50 μ l on a Centricon 30 ultra-filter (Amicon), and washed on the same filter three times with 2 ml TBS and once with 1 ml TBS containing 0.02% NaN_3 and 7×10^{12} UV-inactivated blocking phage (see below); the final retentate (60-80 μ l) is stored at 4°C. Ligand binding proteins biotinylated
20 with the NHS-SS-Biotin reagent are linked to biotin via a disulfide-containing chain.

UV-irradiated M13 phage were used for blocking binding proteins which fortuitously bound filamentous phage in general. M13mp8 (Messing and Vieira, Gene 19: 262-276
25 (1982), which is incorporated herein by reference) was chosen because it carries two amber stop codons, which ensure that the few phage surviving irradiation will not grow in the sup O strains used to titer the surface expression libraries. A 5 ml sample containing 5×10^{13}
30 M13mp8 phage, purified as described above, was placed in a small petri plate and irradiated with a germicidal lamp at a distance of two feet for 7 minutes (flux 150 $\mu\text{W}/\text{cm}^2$). NaN_3 was added to 0.02% and phage particles concentrated to 10^{14} particles/ml on a Centricon 30-kDa ultrafilter
35 (Amicon).

For panning, polystyrene petri plates (60 x 15 mm, Falcon; Becton Dickinson, Lincoln Park, NJ) are incubated with 1 ml of 1 mg/ml of streptavidin (BMB) in 0.1 M NaHCO₃ pH 8.6-0.02% NaN₃ in a small, air-tight plastic box overnight in a cold room. The next day streptavidin is removed and replaced with at least 10 ml blocking solution (29 mg/ml of BSA; 3 µg/ml of streptavidin; 0.1 M NaHCO₃ pH 8.6-0.02% NaN₃) and incubated at least 1 hour at room temperature. The blocking solution is removed and plates are washed rapidly three times with Tris buffered saline containing 0.5% Tween 20 (TBS-0.5% Tween 20).

Selection of phage expressing peptides bound by the ligand binding proteins is performed with 5 µl (2.7 µg ligand binding protein) of blocked biotinylated ligand binding proteins reacted with a 50 µl portion of each library. Each mixture is incubated overnight at 4°C, diluted with 1 ml TBS-0.5% Tween 20, and transferred to a streptavidin-coated petri plate prepared as described above. After rocking 10 minutes at room temperature, unbound phage are removed and plates washed ten times with TBS-0.5% Tween 20 over a period of 30-90 minutes. Bound phage are eluted from plates with 800 µl sterile elution buffer (1 mg/ml BSA, 0.1 M HCl, pH adjusted to 2.2 with glycerol) for 15 minutes and eluates neutralized with 48 µl 2 M Tris (pH unadjusted). A 20 µl portion of each eluate is titered on MK30-3 concentrated cells with dilutions of input phage.

A second round of panning is performed by treating 750 µl of first eluate from each library with 5 mM DTT for 10 minutes to break disulfide bonds linking biotin groups to residual biotinylated binding proteins. The treated eluate is concentrated on a Centricon 30 ultrafilter (Amicon), washed three times with TBS-0.5% Tween 20, and concentrated to a final volume of about 50 µl. Final retentate is transferred to a tube containing 5.0 µl (2.7 µg ligand

binding protein) blocked biotinylated ligand binding proteins and incubated overnight. The solution is diluted with 1 ml TBS-0.5% Tween 20, panned, and eluted as described above on fresh streptavidin-coated petri plates.

- 5 The entire second eluate (800 μ l) is neutralized with 48 μ l 2 M Tris, and 20 μ l is titered simultaneously with the first eluate and dilutions of the input phage.

Individual phage populations are purified through 2 to 3 rounds of plaque purification. Briefly, the second
10 eluate titer plates are lifted with nitrocellulose filters (Schleicher & Schuell, Inc., Keene, NH) and processed by washing for 15 minutes in TBS (10 mM Tris-HCl, pH 7.2, 150 mM NaCl), followed by an incubation with shaking for an additional 1 hour at 37°C with TBS containing 5% nonfat dry
15 milk (TBS-5% NDM) at 0.5 ml/cm². The wash is discarded and fresh TBS-5% NDM is added (0.1 ml/cm²) containing the ligand binding protein between 1 nM to 100 mM, preferably between 1 to 100 μ M. All incubations are carried out in heat-sealable pouches (Sears). Incubation with the ligand
20 binding protein proceeds for 12-16 hours at 4°C with shaking. The filters are removed from the bags and washed 3 times for 30 minutes at room temperature with 150 mls of TBS containing 0.1% NDM and 0.2% NP-40 (Sigma, St. Louis, MO). The filters are then incubated for 2 hours at room
25 temperature in antiserum against the ligand binding protein at an appropriate dilution in TBS-0.5% NDM, washed in 3 changes of TBS containing 0.1% NDM and 0.2% NP-40 as described above and incubated in TBS containing 0.1% NDM and 0.2% NP-40 with 1 x 10⁶ cpm of ¹²⁵I-labeled Protein A
30 (specific activity = 2.1 x 10⁷ cpm/ μ g). After a washing with TBS containing 0.1% NDM and 0.2% NP-40 as described above, the filters are wrapped in Saran Wrap and exposed to Kodak X-Omat x-ray film (Kodak, Rochester, NY) for 1-12 hours at -70°C using Dupont Cronex Lightning Plus
35 Intensifying Screens (Dupont, Willmington, DE).

Positive plaques identified are cored with the large end of a pasteur pipet and placed into 1 ml of SM (5.8 g NaCl, 2 g $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 50 ml 1 M Tris-HCl, pH 7.5, 5 mls 2% gelatin, to 1000 mls with dH_2O) plus 1-3 drops of CHCl_3 and incubated at 37°C 2-3 hours or overnight at 4°C. The phage are diluted 1:500 in SM and 2 μl are added to 300 μl of XL1 cells plus 3 mls of soft agar per 100 mm^2 plate. The XL1 cells are prepared for plating by growing a colony overnight in 10 ml LB (10 g bacto-tryptone, 5 g bacto-yeast extract, 10 g NaCl, 1000 ml dH_2O) containing 100 μl of 20% maltose and 100 μl of 1 M MgSO_4 . The bacteria are pelleted by centrifugation at 2000 xg for 10 minutes and the pellet is resuspended gently in 10 mls of 10 mM MgSO_4 . The suspension is diluted 4-fold by adding 30 mls of 10 mM MgSO_4 to give an OD_{600} of approximately 0.5. The second and third round screens are identical to that described above except that the plaques are cored with the small end of a pasteur pipet and placed into 0.5 mls SM plus a drop of CHCl_3 and 1-5 μl of the phage following incubation are used for plating without dilution. At the end of the third round of purification, an individual plaque is picked and the templates prepared for sequencing.

Template Preparation and Sequencing

Templates are prepared for sequencing by inoculating a 1 ml culture of 2XYT containing a 1:100 dilution of an overnight culture of XL1 with an individual plaque. The plaques are picked using a sterile toothpick. The culture is incubated at 37°C for 5-6 hours with shaking and then transferred to a 1.5 ml microfuge tube. 200 μl of PEG solution is added, followed by vortexing and placed on ice for 10 minutes. The phage precipitate is recovered by centrifugation in a microfuge at 12,000 x g for 5 minutes. The supernatant is discarded and the pellet is resuspended in 230 μl of TE (10 mM Tris-HCl, pH 7.5, 1 mM EDTA) by gently pipeting with a yellow pipet tip. Phenol (200 μl)

is added, followed by a brief vortex and microfuged to separate the phases. The aqueous phase is transferred to a separate tube and extracted with 200 μ l of phenol/chloroform (1:1) as described above for the phenol
5 extraction. A 0.1 volume of 3 M NaOAc is added, followed by addition of 2.5 volumes of ethanol and precipitated at -20°C for 20 minutes. The precipitated templates are recovered by centrifugation in a microfuge at 12,000 x g for 8 minutes. The pellet is washed in 70% ethanol, dried
10 and resuspended in 25 μ l TE. Sequencing was performed using a SequenaseTM sequencing kit following the protocol supplied by the manufacturer (U.S. Biochemical, Cleveland, OH).

EXAMPLE II

15 Isolation and Characterization of Peptide Ligands Generated From Oligonucleotides Having Random Codons at Two Predetermined Positions

This example shows the generation of a surface expression library from a population of oligonucleotides
20 having randomized codons. The oligonucleotides are ten codons in length and are cloned into a single vector species for the generation of a M13 gene VIII-based surface expression library. The example also shows the selection of peptides for a ligand binding protein and
25 characterization of their encoded nucleic acid sequences.

Oligonucleotide Synthesis

Oligonucleotides were synthesized as described in Example I. The synthesizer was programmed to synthesize the sequences shown in Table IX. These sequences
30 correspond to the first random codon position synthesized and 3' flanking sequences of the oligonucleotide which hybridizes to the leader sequence in the vector. The

complementary sequences are used for insertional mutagenesis of the synthesized population of oligonucleotides.

Table IX

5	<u>Column</u>	<u>Sequence (5' to 3')</u>
	column 1	AA (A/C) GGT TGG TCG GTAC CGG
	column 2	AG (A/G) GGT TGG TCG GTAC CGG
	column 3	AT (A/G) GGT TGG TCG GTAC CGG
	column 4	AC (A/G) GGT TGG TCG GTAC CGG
10	column 5	CA (G/T) GGT TGG TCG GTAC CGG
	column 6	CT (G/C) GGT TGG TCG GTAC CGG
	column 7	AG (T/C) GGT TGG TCG GTAC CGG
	column 8	AT (T/C) GGT TGG TCG GTAC CGG
	column 9	CC (A/C) GGT TGG TCG GTAC CGG
15	column 10	T (A/T) TGG TGG TCG GTAC CGG

The next eight random codon positions were synthesized as described for Table V in Example I. Following the ninth position synthesis, the reaction products were once more combined, mixed and redistributed into 10 new reaction
20 columns. Synthesis of the last random codon position and 5' flanking sequences are shown in Table X.

Table X

	<u>Column</u>	<u>Sequence (5' to 3')</u>
25	column 1	AGGATCCGCCGAGCTCAA (A/C) <u>A</u>
	column 2	AGGATCCGCCGAGCTCAG (A/G) <u>A</u>
	column 3	AGGATCCGCCGAGCTCAT (A/G) <u>A</u>
	column 4	AGGATCCGCCGAGCTCAC (A/G) <u>A</u>
	column 5	AGGATCCGCCGAGCTCCA (G/T) <u>A</u>
	column 6	AGGATCCGCCGAGCTCCT (G/C) <u>A</u>
30	column 7	AGGATCCGCCGAGCTCAG (T/C) <u>A</u>
	column 8	AGGATCCGCCGAGCTCAT (T/C) <u>A</u>
	column 9	AGGATCCGCCGAGCTCCC (A/C) <u>A</u>
	column 10	AGGATCCGCCGAGCTCT (A/T) <u>TA</u>

The reaction products were mixed once more and the oligonucleotides cleaved and purified as recommended by the manufacturer. The purified population of oligonucleotides were used to generate a surface expression library as
5 described below.

Vector Construction

The vector used for generating surface expression libraries from a single oligonucleotide population (i.e., without joining together of right and left half
10 oligonucleotides) is described below. The vector is a M13-based expression vector which directs the synthesis of gene VIII-peptide fusion proteins (Figure 4). This vector exhibits all the functions that the combined right and left half vectors of Example I exhibit.

15 An M13-based vector was constructed for the cloning and surface expression of populations of random oligonucleotides (Figure 4, M13IX30), M13mp19 (Pharmacia) was the starting vector. This vector was modified to contain, in addition to the encoded wild type M13 gene
20 VIII: (1) a pseudo-wild type gene, gene VIII sequence with an amber stop codon placed between it and the restriction sites for cloning oligonucleotides; (2) Stu I, Spe I and Xho I restriction sites in frame with the pseudo-wild type gVIII for cloning oligonucleotides; (3) sequences necessary
25 for expression, such as a promoter, signal sequence and translation initiation signals; (4) various other mutations to remove redundant restriction sites and the amino terminal portion of Lac Z.

Construction of M13IX30 was performed in four steps.
30 In the first step, a precursor vector containing the pseudo gene VIII and various other mutations was constructed, M13IX01F. The second step involved the construction of a small cloning site in a separate M13mp18 vector to yield

M13IX03. In the third step, expression sequences and cloning sites were constructed in M13IX03 to generate the intermediate vector M13IX04B. The fourth step involved the incorporation of the newly constructed sequences from the intermediate vector into M13IX01F to yield M13IX30. Incorporation of these sequences linked them with the pseudo gene VIII.

Construction of the precursor vector M13IX01F was similar to that of M13IX42 described in Example I except for the following features: (1) M13mp19 was used as the starting vector; (2) the Fok I site 5' to the unique Eco RI site was not incorporated and the overhang at the naturally occurring Fok I site at position 3547 was not changed to 5'-CTTC-3'; (3) the spacer sequence was not incorporated between the Eco RI and Sac I sites; and (4) the amber codon at position 4492 was not incorporated.

In the second step, M13mp18 was mutated to remove the 5' end of Lac Z up to the Lac i binding site and including the Lac Z ribosome binding site and start codon. Additionally, the polylinker was removed and a Mlu I site was introduced in the coding region of Lac Z. A single oligonucleotide was used for these mutagenesis and had the sequence "5'-AAACGACGGCCAGTGCCAAGTGACGCGTGTGAAATTGTTATCC-3'" (SEQ ID NO: 41). Restriction enzyme sites for Hind III and Eco RI were introduced downstream of the MluI site using the oligonucleotide "5'-GGCGAAAGGGAATTCTGCAAGGCGATTAAGCTTGGGTAACGCC-3'" (SEQ ID NO: 42). These modifications of M13mp18 yielded the vector M13IX03.

The expression sequences and cloning sites were introduced into M13IX03 by chemically synthesizing a series of oligonucleotides which encode both strands of the desired sequence. The oligonucleotides are presented in Table XI (SEQ ID NOS: 43 through 50).

TABLE XI
M13IX30 Oligonucleotide Series

<u>Top Strand</u> <u>Oligonucleotides</u>		<u>Sequence (5' to 3')</u>
5	084	GGCGTTACCCAAGCTTTGTACATGGAGAAAATAAAG
	027	TGAAACAAAGCACTATTGCACTGGCACTCTTACCGT TACCGT
	028	TACTGTTTACCCCTGTGACAAAAGCCGCCCAGGTCC AGCTGC
10	029	TCGAGTCAGGCCTATTGTGCCCAGGGATTGTACTAG TGGATCCG
<u>Bottom</u> <u>Oligonucleotides</u>		<u>Sequence (5' to 3')</u>
	085	TGGCGAAAGGGAATTCGGATCCACTAGTACAATCCCTG
15	031	GGCACAATAGGCCTGACTCGAGCAGCTGGACCAGGGCG GCTT
	032	TTGTCACAGGGGTAAACAGTAACGGTAACGGTAAGTGT GCCA
20	033	GTGCAATAGTGCTTTGTTTCACTTTATTTTCTCCATGT ACAA

The above oligonucleotides except for the terminal oligonucleotides 084 (SEQ ID NO: 43) and 085 (SEQ ID NO: 47) of Table XI were mixed, phosphorylated, annealed and ligated to form a double stranded insert as described in Example I. However, instead of cloning directly into the intermediate vector the insert was first amplified by PCR using the terminal oligonucleotides 084 (SEQ ID NO: 43) and 085 (SEQ ID NO: 47) as primers. The terminal oligonucleotide 084 (SEQ ID NO: 43) contains a Hind III site 10 nucleotides internal to its 5' end. Oligonucleotide 085 (SEQ ID NO: 47) has an Eco RI site at its 5' end. Following amplification, the products were restricted with Hind III and Eco RI and ligated as described in Example I into the polylinker of M13mp18

digested with the same two enzymes. The resultant double stranded insert contained a ribosome binding site, a translation initiation codon followed by a leader sequence and three restriction enzyme sites for cloning random oligonucleotides (Xho I, Stu I, Spe I). The vector was named M13IX04.

During cloning of the double-stranded insert, it was found that one of the GCC codons in oligonucleotides 028 and its complement in 031 was deleted. Since this deletion did not affect function, the final construct is missing one of the two GCC codons. Additionally, oligonucleotide 032 contained a GTG codon where a GAG codon was needed. Mutagenesis was performed using the oligonucleotide 5'-TAACGGTAAGAGTGCCAGTGC-3' (SEQ ID NO: 51) to convert the codon to the desired sequence. The resultant intermediate vector was named M13IX04B.

The fourth step in constructing M13IX30 involved inserting the expression and cloning sequences from M13IX04B upstream of the pseudo-wild type gVIII in M13IX01F. This was accomplished by digesting M13IX04B with Dra III and Ban HI and gel isolating the 700 base pair insert containing the sequences of interest. M13IX01F was likewise digested with Dra III and Bam HI. The insert was combined with the double digested vector at a molar ratio of 3:1 and ligated as described in Example I. It should be noted that all modifications in the vectors described herein were confirmed by sequence analysis. The sequence of the final construct, M13IX30, is shown in Figure 7 (SEQ ID NO: 3). Figure 4 also shows M13IX30 where each of the elements necessary for surface expression of randomized oligonucleotides is marked.

Library Construction, Screening and Characterization of
Encoded Oligonucleotides

Construction of an M13IX30 surface expression library is accomplished identically to that described in Example I for sublibrary construction except the oligonucleotides described above are inserted into M13IX30 by mutagenesis instead of by ligation. The library is constructed and propagated on MK30-3 (BMB) and phage stocks are prepared for infection of XLI cells and screening. The surface expression library is screened and encoding oligonucleotides characterized as described in Example I.

EXAMPLE III

Isolation and Characterization of Peptide Ligands
Generated from Right and Left Half
Degenerate Oligonucleotides

This example shows the construction and expression of a surface expression library of degenerate oligonucleotides. The encoded peptides of this example derive from the mixing and joining together of two separate oligonucleotide populations. Also demonstrated is the isolation and characterization of peptide ligands and their corresponding nucleotide sequence for specific binding proteins.

Synthesis of Oligonucleotide Populations

A population of left half degenerate oligonucleotides and a population of right half degenerate oligonucleotides was synthesized using standard automated procedures as described in Example I.

The degenerate codon sequences for each population of oligonucleotides were generated by sequentially

synthesizing the triplet NNG/T where N is an equal mixture of all four nucleotides. The antisense sequence for each population of oligonucleotides was synthesized and each population contained 5' and 3' flanking sequences complementary to the vector sequence. The complementary termini was used to incorporate each population of oligonucleotides into their respective vectors by standard mutagenesis procedures. Such procedures have been described previously in Example I and in the Detailed Description. Synthesis of the antisense sequence of each population was necessary since the single-stranded form of the vectors are obtained only as the sense strand.

The left half oligonucleotide population was synthesized having the following sequence: 5'-AGCTCCCGGATGCCTCAGAAGATG(A/CNN)₉GGCTTTTGCCACAGGGG-3' (SEQ ID NO: 52). The right half oligonucleotide population was synthesized having the following sequence: 5'-CAGCCTCGGATCCGCC(A/CNN)₁₀ATG(A/C)GAAT-3' (SEQ ID NO. 53). These two oligonucleotide populations when incorporated into their respective vectors and joined together encode a 20 codon oligonucleotide having 19 degenerate positions and an internal predetermined codon sequence.

Vector Construction

Modified forms of the previously described vectors were used for the construction of right and left half sublibraries. The construction of left half sublibraries was performed in an M13-based vector termed M13ED03. This vector is a modified form of the previously described M13IX30 vector and contains all the essential features of both M13IX30 and M13IX22. M13ED03 contains, in addition to a wild type and a pseudo-wild type gene VIII, sequences necessary for expression and two Fok I sites for joining with a right half oligonucleotide

sublibrary. Therefore, this vector combines the advantages of both previous vectors in that it can be used for the generation and expression of surface expression libraries from a single oligonucleotide population or it can be joined with a sublibrary to bring together right and left half oligonucleotide populations into a surface expression library.

M13ED03 was constructed in two steps from M13IX30. The first step involved the modification of M13IX30 to remove a redundant sequence and to incorporate a sequence encoding the eight amino-terminal residues of human β -endorphin. The leader sequence was also mutated to increase secretion of the product.

During construction of M13IX04 (an intermediate vector to M13IX30 which is described in Example II), a six nucleotide sequence was duplicated in oligonucleotide 027 (SEQ ID NO: 44) and its complement 032 (SEQ ID NO: 49). This sequence, 5'-TTACCG-3', was deleted by mutagenesis in the construction of M13ED01. The oligonucleotide used for the mutagenesis was 5'-GGTAAACAGTAACGGTAAGAGTGCCAG-3' (SEQ ID NO: 54). The mutation in the leader sequence was generated using the oligonucleotide 5'-GGGCTTTTGCCACAGGGGT-3' (SEQ ID NO: 55). This mutagenesis resulted in the A residue at position 6353 of M13IX30 being changed to a G residue. The resultant vector was designated M13IX32.

To generate M13ED01, the nucleotide sequence encoding β -endorphin (8 amino acid residues of β -endorphin plus 3 extra amino acid residues) was incorporated after the leader sequence by mutagenesis. The oligonucleotide used had the following sequence: 5'-AGGGTCATCGCCTTCAGCTCCGGATCCCTCAGAAGTCATAAACCCCCATAGGCTTTTGCCAC-3' (SEQ ID NO: 56). This mutagenesis also removed some of the downstream sequences through the Spe

I site.

The second step in the construction of M13ED03 involved vector changes which put the β -endorphin sequence in frame with the downstream pseudo-gene VIII sequence and incorporated a Fok I site for joining with a sublibrary of right half oligonucleotides. This vector was designed to incorporate oligonucleotide populations by mutagenesis using sequences complementary to those flanking or overlapping with the encoded β -endorphin sequence. The absence of β -endorphin expression after mutagenesis can therefore be used to measure the mutagenesis frequency. In addition to the above vector changes, M13ED03 was also modified to contain an amber codon at position 3262 for biological selection during joining of right and left half sublibraries.

The mutations were incorporated using standard mutagenesis procedures as described in Example I. The frame shift changes and Fok I site were generated using the oligonucleotide 5'-

TCGCCTTCAGCTCCCGGATGCCTCAGAAGCATGAACCCCCCATAGGC-3' (SEQ ID NO: 57). The amber codon was generated using the oligonucleotide 5'-CAATTTTATCCTAAATCTTACCAAC-3' (SEQ ID NO: 58). The full sequence of the resultant vector, M13ED03, is provided in Figure 8 (SEQ ID NO: 4).

The construction of right half oligonucleotide sublibraries was performed in a modified form of the M13IX42 vector. The new vector, M13IX421, is identical to M13IX42 except that the amber codon between the Eco RI-SacI cloning site and the pseudo-gene VIII sequence was removed. This change ensures that all expression off of the Lac Z promoter produces a peptide-gene VIII fusion protein. Removal of the amber codon was performed by mutagenesis using the following oligonucleotide: 5'-GCCTTCAGCCTCGGATCCGCC-3' (SEQ ID NO: 59). The full

sequence of M13IX421 is shown in Figure 9 (SEQ ID NO: 5).

Library Construction, Screening and Characterization of
Encoded Oligonucleotides

A sublibrary was constructed for each of the
5 previously described degenerate populations of
oligonucleotides. The left half population of
oligonucleotides was incorporated into M13ED03 to
generate the sublibrary M13ED03.L and the right half
10 population of oligonucleotides was incorporated into
M13IX421 to generate the sublibrary M13IX421.R. Each of
the oligonucleotide populations were incorporated into
their respective vectors using site-directed mutagenesis
as described in Example I. Briefly, the nucleotide
15 sequences flanking the degenerate codon sequences were
complementary to the vector at the site of incorporation.
The populations of nucleotides were hybridized to single-
stranded M13ED03 or M13IX421 vectors and extended with T4
DNA polymerase to generate a double-stranded circular
vector. Mutant templates were obtained by uridine
20 selection in vivo, as described by Kunkel et al., supra.
Each of the vector populations were electroporated into
host cells and propagated as described in Example I.

The random joining of right and left half
sublibraries into a single surface expression library was
25 accomplished as described in Example I except that prior
to digesting each vector population with Fok I they were
first digested with an enzyme that cuts in the unwanted
portion of each vector. Briefly, M13ED03.L was digested
with Bgl II (cuts at 7094) and M13IX421.R was digested
30 with Hind III (cuts at 3919). Each of the digested
populations were further treated with alkaline
phosphatase to ensure that the ends would not religate
and then digested with an excess of Fok I. Ligations,
electroporation and propagation of the resultant library

was performed as described in Example I.

The surface expression library was screened for ligand binding proteins using a modified panning procedure. Briefly, 1 ml of the library, about 10^{12} phage particles, was added to 1-5 μ g of the ligand binding protein. The ligand binding protein was either an antibody or receptor globulin (Rg) molecule, Aruffo et al., Cell 61:1303-1313 (1990), which is incorporated herein by reference. Phage were incubated shaking with affinity ligand at room temperature for 1 to 3 hours followed by the addition of 200 μ l of latex beads (Biosite, San Diego, CA) which were coated with goat-antimouse IgG. This mixture was incubated shaking for an additional 1-2 hours at room temperature. Beads were pelleted for 2 minutes by centrifugation in a microfuge and washed with TBS which can contain 0.1% Tween 20. Three additional washes were performed where the last wash did not contain any Tween 20. The bound phage were then eluted with 200 μ l 0.1 M Glycine-HCl, pH 2.2 for 15 minutes and the beads were spun down by centrifugation. The supernatant-containing phage (eluate) was removed and phage exhibiting binding to the ligand binding protein were further enriched by one-to-two more cycles of panning. Typical yields after the first eluate were about 1×10^6 - 5×10^6 pfu. The second and third eluate generally yielded about 5×10^6 - 2×10^7 pfu and 5×10^7 - 1×10^{10} pfu, respectively.

The second or third eluate was plated at a suitable density for plaque identification screening and sequencing of positive clones (i.e., plated at confluency for rare clones and 200-500 plaques/plate if pure plaques were needed). Briefly, plaques grown for about 6 hours at 37°C and were overlaid with nitrocellulose filters that had been soaked in 2 mM IPTG and then briefly dried. The filters remained on the plaques overnight at room

- temperature, removed and placed in blocking solution for 1-2 hours. Following blocking, the filters were incubated in 1 μ g/ml ligand binding protein in blocking solution for 1-2 hours at room temperature. Goat
- 5 antmouse Ig-coupled alkaline phosphatase (Fisher) was added at a 1:1000 dilution and the filters were rapidly washed with 10 mls of TBS or block solution over a glass vacuum filter. Positive plaques were identified after alkaline phosphatase development for detection.
- 10 Screening of the degenerate oligonucleotide library with several different ligand binding proteins resulted in the identification of peptide sequences which bound to each of the ligands. For example, screening with an antibody to β -endorphin resulted in the detection of
- 15 about 30-40 different clones which essentially all had the core amino acid sequence known to interact with the antibody. The sequences flanking the core sequences were different showing that they were independently derived and not duplicates of the same clone. Screening with an
- 20 antibody known as 57 gave similar results (i.e., a core consensus sequence was identified but the flanking sequences among the clones were different).

EXAMPLE IV

Generation of a Left Half Random Oligonucleotide Library

- 25 This example shows the synthesis and construction of a left half random oligonucleotide library.

- A population of random oligonucleotides nine codons in length was synthesized as described in Example I except that different sequences at their 5' and 3' ends
- 30 were synthesized so that they could be easily inserted into the vector by mutagenesis. Also, the mixing and dividing steps for generating random distributions of

reaction products was performed by the alternative method of dispensing equal volumes of bead suspensions. The liquid chosen that was dense enough for the beads to remain dispersed was 100% acetonitrile.

- 5 Briefly, each column was prepared for the first coupling reaction by suspending 22 mg (1 μ mole) of 48 μ mol/g capacity beads (Genta, San Diego, CA) in 0.5 mls of 100% acetonitrile. These beads are smaller than those described in Example I and are derivatized with a guanine
 10 nucleotide. They also do not have a controlled pore size. The bead suspension was then transferred to an empty reaction column. Suspensions were kept relatively dispersed by gently pipetting the suspension during transfer. Columns were plugged and monomer coupling
 15 reactions were performed as shown in Table XII.

Table XII

	<u>Column</u>	<u>Sequence</u> <u>(5' to 3')</u>
	column 1L	AA(A/C)GGCTTTTGCCACAGG
20	column 2L	AG(A/G)GGCTTTTGCCACAGG
	column 3L	AT(A/G)GGCTTTTGCCACAGG
	column 4L	AC(A/G)GGCTTTTGCCACAGG
	column 5L	CA(G/T)GGCTTTTGCCACAGG
	column 6L	CT(G/C)GGCTTTTGCCACAGG
25	column 7L	AG(T/C)GGCTTTTGCCACAGG
	column 8L	AT(T/C)GGCTTTTGCCACAGG
	column 9L	CC(A/C)GGCTTTTGCCACAGG
	column 10L	T(A/T)TGGCTTTTGCCACAGG

- After coupling of the last monomer, the columns were
 30 unplugged as described previously and their contents were poured into a 1.5 ml microfuge tube. The columns were rinsed with 100% acetonitrile to recover any remaining beads. The volume used for rinsing was determined so

that the final volume of total bead suspension was about 100 μ l for each new reaction column that the beads would be aliquoted into. The mixture was vortexed gently to produce a uniformly dispersed suspension and then

5 divided, with constant pipetting of the mixture, into equal volumes. Each mixture of beads was then transferred to an empty reaction column. The empty tubes were washed with a small volume of 100% acetonitrile and also transferred to their respective columns. Random

10 codon positions 2 through 9 were then synthesized as described in Example I where the mixing and dividing steps were performed using a suspension in 100% acetonitrile. The coupling reactions for codon positions 2 through 9 are shown in Table XIII.

15

Table XIII

	<u>Column</u>	<u>Sequence (5' to 3')</u>
	column 1L	AA(A/C) <u>A</u>
	column 2L	AG(A/G) <u>A</u>
20	column 3L	AT(A/G) <u>A</u>
	column 4L	AC(A/G) <u>A</u>
	column 5L	CA(G/T) <u>A</u>
	column 6L	CT(G/C) <u>A</u>
	column 7L	AG(T/C) <u>A</u>
25	column 8L	AT(T/C) <u>A</u>
	column 9L	CC(A/C) <u>A</u>
	column 10L	T(A/T)T <u>A</u>

After coupling of the last monomer for the ninth codon position, the reaction products were mixed and a

30 portion was transferred to an empty reaction column. Columns were plugged and the following monomer coupling reactions were performed: 5'-CGGATGCCTCAGAAGCCCCXXA-3' (SEQ ID NO: 60). The resulting population of random oligonucleotides was purified and incorporated by

mutagenesis into the left half vector M13ED04.

M13ED04 is a modified version of the M13ED03 vector described in Example III and therefore contains all the features of that vector. The difference between M13ED03
5 and M13ED04 is that M13ED04 does not contain the five amino acid sequence (Tyr Gly Gly Phe Met) recognized by anti- β -endorphin antibody. This sequence was deleted by mutagenesis using the oligonucleotide 5'-
CGGATGCCTCAGAAGGGCTTTGCCACAGG (SEQ ID NO: 61). The
10 entire nucleotide sequence of this vector is shown in Figure 10 (SEQ ID NO: 6).

Although the invention has been described with reference to the presently preferred embodiment, it should be understood that various modifications can be
15 made without departing from the spirit of the invention. Accordingly, the invention is limited only by the claims.

SEQUENCE LISTING

(1) GENERAL INFORMATION:

- (i) APPLICANT: Huse, William D.
- (ii) TITLE OF INVENTION: SURFACE EXPRESSION LIBRARIES OF RANDOMIZED PEPTIDES
- (iii) NUMBER OF SEQUENCES: 61
- (iv) CORRESPONDENCE ADDRESS:
 - (A) ADDRESSEE: Pretty, Schroeder, Brueggemann & Clark
 - (B) STREET: 444 South Flower Street, Suite 2000
 - (C) CITY: Los Angeles
 - (D) STATE: California
 - (E) COUNTRY: United States
 - (F) ZIP: 90071
- (v) COMPUTER READABLE FORM:
 - (A) MEDIUM TYPE: Floppy disk
 - (B) COMPUTER: IBM PC compatible
 - (C) OPERATING SYSTEM: PC-DOS/MS-DOS
 - (D) SOFTWARE: PatentIn Release #1.0, Version #1.25
- (vi) CURRENT APPLICATION DATA:
 - (A) APPLICATION NUMBER:
 - (B) FILING DATE:
 - (C) CLASSIFICATION:
- (viii) ATTORNEY/AGENT INFORMATION:
 - (A) NAME: Campbell, Cathryn A
 - (B) REGISTRATION NUMBER: 31,815
 - (C) REFERENCE/DOCKET NUMBER: P31 9072
- (ix) TELECOMMUNICATION INFORMATION:
 - (A) TELEPHONE: (619) 535-9001
 - (B) TELEFAX: (619) 535-8949

(2) INFORMATION FOR SEQ ID NO:1:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 7294 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: both
 - (D) TOPOLOGY: circular

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

AATGCTACTA CTATTAGTAG AATTGATGCC ACCTTTTCAG CTCGCGCCCC AAATGAAAAT	60
ATAGCTAAAC AGGTTATTGA CCATTTCGGA AATGTATCTA ATGGTCAAAC TAAATCTACT	120
CGTTCGCAGA ATTGGGAATC AACTGTTACA TGGAATGAAA CTTCCAGACA CCGTACTTTA	180
GTTGCATATT TAAAACATGT TGAGCTACAG CACCAGATTC AGCAATTAAAG CTCTAAGCCA	240
TCTGCAAAAA TGACCTCTTA TCAAAAGGAG CAATTAAAGG TACTCTCTAA TCCTGACCTG	300
TTGGAGTTTG CTTCCGGTCT GGTTCGCTTT GAAGCTCGAA TTAAAACGCG ATATTTGAAG	360
TCTTTCGGGC TTCCTCTTAA TCTTTTGTAT GCAATCCGCT TTGCTTCTGA CTATAATAGT	420

CAGGGTAAAG ACCTGATTTT TGATTTATGG TCATTCTCGT TTTCTGAACT GTTTAAAGCA	480
TTTGAGGGGG ATTCAATGAA TATTTATGAC GATTCCGCAG TATTGGACGC TATCCAGTCT	540
AAACATTTTA CTATTACCCC CTCTGGCAA ACTTCTTTTG CAAAAGCCTC TCGCTATTTT	600
GGTTTTATC GTCGTCTGGT AAACGAGGGT TATGATAGTG TTGCTCTTAC TATGCCTCGT	660
AATTCCTTTT GCGGTATGT ATCTGCATTA GTTGAATGTG GTATTCCTAA ATCTCAACTG	720
ATGAATCTTT CTACCTGTAA TAATGTTGTT CCGTTAGTTC GTTTTATTAA CGTAGATTTT	780
TCTTCCCAAC GTCCTGACTG GTATAATGAG CCAGTTCCTA AAATCGCATA AGGTAATTCA	840
CAATGATTAA AGTTGAAATT AAACCATCTC AAGCCCAATT TACTACTCGT TCTGGTGTTC	900
CTCGTCAGGG CAAGCCTTAT TCACTGAATG AGCAGCTTTG TTACGTTGAT TTGGGTAATG	960
AATATCCGGT TCTTGTCAAG ATTACTCTTG ATGAAGGTCA GCCAGCCTAT GCGCCTGGTC	1020
TGTACACCGT TCATCTGTCC TCTTTCAAAG TTGGTCAGTT CGGTTCCCTT ATGATTGACC	1080
GTCTGCGCCT CGTTCGGCT AAGTAACATG GAGCAGGTGG CGGATTTCTGA CACAATTTAT	1140
CAGGCGATGA TACAAATCTC CGTTGTACTT TGTTCGCGC TTGGTATAAT CGCTGGGGGT	1200
CAAAGATGAG TGTTTTAGTG TATCTTTTCG CCTCTTTCTG TTTAGGTTGG TGCCTTCGTA	1260
GTGGCATTAC GTATTTTACC CGTTTAATGG AAACCTCCTC ATGAAAAAGT CTTTAGTCCT	1320
CAAAGCCTCT GTAGCCGTTG CTACCCTCGT TCCGATGCTG TCTTTCGCTG CTGAGGGTGA	1380
CGATCCCGCA AAAGCGGCCT TTAACCTCCCT GCAAGCCTCA GCGACCGAAT ATATCGGTTA	1440
TGCGTGGGCG ATGGTTGTTG TCATTGTCCG CGCAACTATC GGTATCAAGC TGTTTAAGAA	1500
ATTCACCTCG AAAGCAAGCT GATAAACCGA TACAATTAAA GGCTCCTTTT GGAGCCTTTT	1560
TTTTTGAGA TTTTCAACGT GAAAAAATTA TTATTCGCAA TTCCTTTAGT TGTTCTTTTC	1620
TATTCTCACT CCGCTGAAAC TGTGAAAGT TGTTTAGCAA AACCCCATAC AGAAAAATCA	1680
TTTACTAACG TCTGGAAAGA CGACAAAAC TTAGATCGTT ACGCTAACTA TGAGGGTTGT	1740
CTGTGGAATG CTACAGGCGT TGAGTTTGT ACTGGTGACG AAACCTCAGT TTACGGTACA	1800
TGGGTTCTTA TTGGGCTTGC TATCCCTGAA AATGAGGGTG GTGGCTCTGA GGGTGGCGGT	1860
TCTGAGGGTG GCGGTTCTGA GGGTGGCGGT ACTAAACCTC CTGAGTACGG TGATACACCT	1920
ATTCGGGGCT ATACTTATAT CAACCTCTC GACGGCACTT ATCCGCCTGG TACTGAGCAA	1980
AACCCCGCTA ATCCTAATCC TTCTCTTGAG GAGTCTCAGC CTCTTAATAC TTTCATGTTT	2040
CAGAATAATA GGTTCGAAA TAGGCAGGGG GCATTAACTG TTTATACGGG CACTGTTACT	2100
CAAGGCACTG ACCCCGTAA AACTTATTAC CAGTACACTC CTGTATCATC AAAAGCCATG	2160
TATGACGCTT ACTGGAACGG TAAATTCAGA GACTGCGCTT TCCATTCTGG CTTTAATGAA	2220
GATCCATTCT TTTGTGAATA TCAAGGCCAA TCGTCTGACC TGCCTCAACC TCCTGTCAAT	2280
GCTGGCGGCG GCTCTGGTGG TGGTTCTGGT GGCGGCTCTG AGGGTGGTGG CTCTGAGGGT	2340
GGCGGTTCTG AGGGTGGCGG CTCTGAGGGA GGCGGTTCCG GTGGTGGCTC TGGTTCCGGT	2400
GATTTTGATT ATGAAAAGAT GGCAAACGCT AATAAGGGGG CTATGACCGA AAATGCCGAT	2460

GAAAACGCGC	TACAGTCTGA	CGCTAAAGGC	AAACTTGATT	CTGTGGCTAC	TGATTACGGT	2520
GCTGCTATCG	ATGGTTTCAT	TGGTGACGTT	TCCGGGCCCTG	CTAATGGTAA	TGGTGCTACT	2580
GGTGATTTTG	CTGGCTCTAA	TTCCCAAATG	GCTCAAGTCG	GTGACGGTGA	TAATTCACCT	2640
TTAATGAATA	ATTTCCGTCA	ATATTTACCT	TCCCTCCCTC	AATCGGTTGA	ATGTCGCCCT	2700
TTTGTCTTTA	GGGCTGGTAA	ACCATATGAA	TTTTCTATTG	ATTGTGACAA	AATAAACTTA	2760
TTCCGTGGTG	TCTTTGCGTT	TCTTTTATAT	GTTGCCACCT	TTATGTATGT	ATTTTCTACG	2820
TTTGCTAACA	TACTGCGTAA	TAAGGAGTCT	TAATCATGCC	AGTTCTTTTG	GGTATTCCGT	2880
TATTATTGGC	TTTCCCTCGGT	TTCCTTCTGG	TAACTTTGTT	CGGCTATCTG	CTTACTTTTC	2940
TTAAAAAGGG	CTTCGGTAAG	ATAGCTATTG	CTATTTGATT	GTTTCTTGCT	CTTATTATTG	3000
GGCTTAACTC	AATTCTTGTT	GGTTATCTCT	CTGATATTAG	CGCTCAATTG	CCCTCTGACT	3060
TTGTTTACGG	TGTTTACGTT	ATTCTCCCGT	CTAATGCGCT	TCCCTGTTTT	TATGTTATTG	3120
TCTCTGTAAA	GGCTGCTATT	TTCATTTTTG	ACGTAAACA	AAAAATCGTT	TCTTATTG	3180
ATTGGGATAA	ATAATATGGC	TGTTTATTTT	GTAAGTGGCA	AATTAGGCTC	TGGAAAGACG	3240
CTCGTTAGCG	TTGGTAAGAT	TCAGGATAAA	ATTGTAGCTG	GGTGCAAAAT	AGCAACTAAT	3300
CTTGATTTAA	GGCTTCAAAA	CCTCCCGCAA	GTCCGGAGGT	TCGCTAAAC	GCCTCGCGTT	3360
CTTAGAATAC	CGGATAAGCC	TTCTATATCT	GATTTGCTTG	CTATTGGGCG	CGGTAATGAT	3420
TCCTACGATG	AAAATAAAAA	CGGCTTGCTT	GTTCTCGATG	AGTGCGGTAC	TTGGTTTAAT	3480
ACCCGTTCTT	GGAATGATAA	GGAAAGACAG	CCGATTATTG	ATTGGTTTCT	ACATGCTCGT	3540
AAATTAGGAT	GGGATATTAT	CTTCCTTGTT	CAGGACTTAT	CTATTGTTGA	TAAACAGGCG	3600
CGTTCTGCAT	TAGCTGAACA	TGTTGTTTAT	TGTCGTCGTC	TGGACAGAAT	TACTTTACCT	3660
TTTGTGGGTA	CTTTATATTG	TCTTATTACT	GGCTCGAAAA	TGCCTCTGCC	TAAATTACAT	3720
GTTGGCGTTG	TTAAATATGG	CGATTCTCAA	TTAAGCCCTA	CTGTTGAGCG	TTGGCTTTAT	3780
ACTGGTAAGA	ATTTGTATAA	CGCATATGAT	ACTAAACAGG	CTTTTTCTAG	TAATTATGAT	3840
TCCGGTGTTT	ATTCTTATTT	AACGCCTTAT	TTATCACACG	GTCGGTATTT	CAAACCATTG	3900
AATTTAGGTC	AGAAGATGAA	GCTTACTAAA	ATATATTTGA	AAAAGTTTTG	ACGCGTTCTT	3960
TGTCTTGCGA	TTGGATTGTC	ATCAGCATTG	ACATATAGTT	ATATAACCCA	ACCTAAGCCG	4020
GAGGTTAAAA	AGGTAGTCTC	TCAGACCTAT	GATTTTGATA	AATTCACAT	TGACTCTTCT	4080
CAGCGTCTTA	ATCTAAGCTA	TCGCTATGTT	TTCAAGGATT	CTAAGGGAAA	ATTAATTAAT	4140
AGCGACGATT	TACAGAAGCA	AGGTTATTCA	CTCACATATA	TTGATTTATG	TACTGTTTCC	4200
ATTAAAAAGG	TAATTCAAAT	GAAATTGTTA	AATGTAATTA	ATTTTGTTTT	CTTGATGTTT	4260
GTTTCATCAT	CTTCTTTTGC	TCAGGTAATT	GAAATGAATA	ATTCGCCTCT	GCGCGATTTT	4320
GTAACCTGGT	ATTCAAAGCA	ATCAGGCGAA	TCCGTTATTG	TTTCTCCCGA	TGTAAAAGGT	4380
ACTGTTACTG	TATATTCATC	TGACGTTAAA	CCTGAAAATC	TACGCAATTT	CTTTATTTCT	4440
GTTTTACGTG	CTAATAATTT	TGATATGGTT	GGTTCAATTC	CTTCCATTAT	TTAGAAGTAT	4500

AATCCAAACA ATCAGGATTA TATTGATGAA TTGCCATCAT CTGATAATCA GGAATATGAT	4560
GATAATTCCG CTCCTTCTGG TGGTTTCTTT GTTCCGCAAA ATGATAATGT TACTCAAAC	4620
TTTAAAATTA ATAACGTTCC GGCAAAGGAT TTAATACGAG TTGTCGAATT GTTTGTAAAG	4680
TCTAATACTT CTAAATCCTC AAATGTATTA TCTATTGACG GCTCTAATCT ATTAGTTGTT	4740
AGTGACCTA AAGATATTTT AGATAACCTT CCTCAATTCC TTTCTACTGT TGATTTGCCA	4800
ACTGACCAGA TATTGATTGA GGGTTTGATA TTTGAGGTTC AGCAAGGTGA TGCTTTAGAT	4860
TTTTCATTTG CTGCTGGCTC TCAGCGTGGC ACTGTTGCAG GCGGTGTAA TACTGACCGC	4920
CTCACCTCTG TTTTATCTTC TGCTGGTGGT TCGTTCCGTA TTTTAAATGG CGATGTTTAA	4980
GGGCTATCAG TTCGCGCATT AAAGACTAAT AGCCATTCAA AAATATTGTC TGTGCCACGT	5040
ATTCTTACGC TTTGAGGTCA GAAGGTTCT ATCTCTGTTG GCCAGAATGT CCCTTTTATT	5100
ACTGGTCGTG TGACTGGTGA ATCTGCCAAT GTAAATAATC CATTTTCAGAC GATTGACCGT	5160
CAAAATGTAG GTATTTCCAT GAGCGTTTTT CCTGTTGCAA TGGCTGGCGG TAATATTGTT	5220
CTGGATATTA CCAGCAAGGC CGATAGTTTG AGTTCTTCTA CTCAGGCAAG TGATGTTATT	5280
ACTAATCAAA GAAGTATTGC TACAACGGTT AATTTGCGTG ATGGACAGAC TCTTTTACTC	5340
GGTGGCCTCA CTGATTATAA AAACACTTCT CAAGATTCTG GCGTACCGTT CCTGTCTAAA	5400
ATCGCTTTAA TCGGCCTCCT GTTTAGCTCC CGCTCTGATT CCAACGAGGA AAGCAGTTA	5460
TACGTGCTCG TCAAAGCAAC CATAGTACGC GCCCTGTAGC GGCGCATTAA GCGCGCGGG	5520
TGTGGTGGTT ACGCGCAGCG TGACCGCTAC ACTTGCCAGC GCCCTAGCGC CCGCTCCTTT	5580
CGCTTTCTTC CCTTCCTTC TCGCCACGTT CGCCGGCTTT CCGGTGAAG CTCTAAATCG	5640
GGGGCTCCCT TTAGGTTCC GATTAGTGC TTTACGGCAC CTCGACCCCA AAAAAGTTGA	5700
TTTGGGTGAT GGTTACGTA GTGGGCCATC GCCCTGATAG ACGGTTTTTC GCCCTTGAC	5760
GTTGGAGTCC ACGTTCTTTA ATAGTGGACT CTTGTTCCAA ACTGGAACAA CACTCAACCC	5820
TATCTCGGGC TATTCTTTTG ATTTATAAGG GATTTTGCCG ATTTGGAAC CACCATCAAA	5880
CAGGATTTTC GCCTGCTGGG GCAAACCAGC GTGGACCGCT TGCTGCAACT CTCTAGGGC	5940
CAGGCGGTGA AGGGCAATCA GCTGTTGCCC GTCTCGCTGG TGAAAAGAAA AACCAACCTG	6000
GCGCCCAATA CGCAAACCGC CTCTCCCCGC GCGTTGGCCG ATTCATTAAT GCAGCTGGCA	6060
CGACAGGTTT CCCGACTGGA AAGCGGGCAG TGAGCGCAAC GCAATTAATG TGAGTTAGCT	6120
CACTCATTAG GCACCCAGG CTTTACACTT TATGCTTCCG GCTCGTATGT TGTGTGGAAT	6180
TGTGAGCGGA TAACAATTTT ACACAGGAAA CAGCTATGAC CAGGATGTAC GAATTCGCAG	6240
GTAGGAGAGC TCGGCGGATC CTAGGCTGAA GGCGATGACC CTGCTAAGGC TGCATTCAAT	6300
AGTTTACAGG CAAGTGCTAC TGAGTACATT GGCTACGCTT GGGCTATGGT AGTAGTTATA	6360
GTTGGTGCTA CCATAGGAT TAAATTATTC AAAAAGTTTA CGAGCAAGGC TTCTTAACCA	6420
GCTGGCGTAA TAGCGAAGAG GCGCGACCG ATCGCCCTTC CCAACAGTTG CGCAGCCTGA	6480
ATGGCGAATG GCGCTTTGCC TGGTTTCCGG CACCAGAAGC GGTGCCGGA AGCTGGCTGG	6540

AGTGGGATCT TCCTGAGGCC GATACGGTCG TCGTCCGCTC AACTGGCAG ATGCACGGTT	6600
ACGATGCGCC CATCTACACC AACGTAACCT ATCCGATTAC GGTCAATCCG CCGTTTGTTT	6660
CCACGGAGAA TCCGACGGGT TGTTACTCGC TCACATTTAA TGTGATGAA AGCTGGCTAC	6720
AGGAAGGCCA GACGCGAATT ATTTTGTATG GCGTTCCTAT TGGTTAAAAA ATGAGCTGAT	6780
TTAACAAAAA TTTAACGCGA ATTTTAACAA AATATTAACG TTTACAATTT AAATATTTGC	6840
TTATACAATC TTCCTGTTTT TGGGGCTTTT CTGATTATCA ACCGGGGTAC ATATGATTGA	6900
CATGCTAGTT TTACGATTAC CGTTCATCGA TTCTCTTGTT TGCTCCAGAC TCTCAGGCAA	6960
TGACCTGATA GCCTTTGTAG ATCTCTCAA AATAGCTACC CTCTCCGGCA TTAATTTATC	7020
AGCTAGAACG GTTGAATATC ATATTGATGG TGATTGACT GTCTCCGGCC TTTCTCACCC	7080
TTTTGAATCT TTACCTACAC ATTACTCAGG CATTGCATTT AAAATATATG AGGGTTCTAA	7140
AAATTTTTAT CCTTGCGTTG AAATAAAGGC TTCTCCCGCA AAAGTATTAC AGGGTCATAA	7200
TGTTTTTGGT ACAACCGATT TAGCTTTATG CTCTGAGGCT TTATTGCTTA ATTTTGCTAA	7260
TTCTTTGCCT TGCCTGTATG ATTTATTGGA CGTT	7294

(2) INFORMATION FOR SEQ ID NO:2:

(1) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 7320 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: both
- (D) TOPOLOGY: circular

(x1) SEQUENCE DESCRIPTION: SEQ ID NO:2:

AATGCTACTA CTATTAGTAG AATTGATGCC ACCTTTTCAG CTCGCGCCCC AAATGAAAAT	60
ATAGCTAAAC AGGTTATTGA CCATTGCGA AATGTATCTA ATGGTCAAAC TAAATCTACT	120
CGTTCGAGA ATTGGGAATC AACTGTTACA TGGAATGAAA CTTCCAGACA CCGTACTTTA	180
GTTGCATATT TAAAACATGT TGAGCTACAG CACCAGATTC AGCAATTAAG CTCTAAGCCA	240
TCTGCAAAAA TGACCTCTTA TCAAAAGGAG CAATTAAAGG TACTCTCTAA TCCTGACCTG	300
TTGGAGTTTG CTTCCGGTCT GGTTGCTTT GAAGCTCGAA TTAAAACGCG ATATTTGAAG	360
TCTTTGCGGC TTCCTCTTAA TCTTTTGTAT GCAATCCGCT TTGCTTCTGA CTATAATAGT	420
CAGGCTAAAG ACCTGATTTT TGATTTATGG TCATTCTCGT TTTCTGAACT GTTTAAAGCA	480
TTTGAGGGGG ATTCAATGAA TATTATGAC GATTCCGCAG TATTGGACGC TATCCAGTCT	540
AAACATTTTA CTATTACCCC CTCTGGCAAA ACTTCTTTTG CAAAAGCCTC TCGCTATTTT	600
GGTTTTTATC GTCGTCTGGT AAACGAGGGT TATGATAGTG TTGCTCTTAC TATGCCTCGT	660
AATTCCTTTT GGCCTTATGT ATCTGCATTA GTTGAATGTG GTATTCCTAA ATCTCAACTG	720
ATGAATCTTT CTACCTGTAA TAATGTTGTT CCGTTAGTTC GTTTTATTAA CGTAGATTTT	780
TCTTCCCAAC GTCCTGACTG GTATAATGAG CCAGTTCTTA AAATCGCATA AGGTAATTCA	840

CAATGATTAA AGTTGAAATT AAACCATCTC AAGCCCAATT TACTACTCGT TCTGGTGT	900
CTCGTCAGGG CAAGCCTTAT TCACTGAATG AGCAGCTTTG TTACGTTGAT TTGGGTAATG	960
AATATCCGGT TCTTGTCAAG ATTACTCTTG ATGAAGGTCA GCCAGCCTAT GCGCCTGGTC	1020
TGTACACCGT TCATCTGTCC TCTTTCAAAG TTGGTCAGTT CGGTTCCCTT ATGATTGACC	1080
GTCTGCGCCT CGTTCCGGCT AAGTAACATG GAGCAGGTG CCGATTTCTGA CACAATTTAT	1140
CAGGCGATGA TACAAATCTC CGTTGTACTT TGTTTCGGCGC TTGGTATAAT CGCTGGGGGT	1200
CAAAGATGAG TGTTTTAGTG TATTCTTTCC CCTCTTTCGT TTTAGGTTGG TGCCTTCGTA	1260
GTGGCATTAC GTATTTTACC CGTTTAATGG AAACCTTCCTC ATGAAAAAGT CTTTAGTCCT	1320
CAAAGCCTCT GTAGCCGTTG CTACCCTCGT TCCGATGCTG TCTTTCGCTG CTGAGGGTGA	1380
CGATCCCGCA AAAGCGGCCT TTAACCTCCCT GCAAGCCTCA GCGACCGAAT ATATCGGTTA	1440
TGCGTGGGCG ATGTTTGTG TCATTGTCGG CGCAACTATC GGTATCAAGC TGTTTAAGAA	1500
ATTCACCTCG AAAGCAAGCT GATAAACCGA TACAATTAAG GGCTCCTTTT GGAGCCTTTT	1560
TTTTTGAGA TTTTCAACGT GAAAAATTA TTATTCGCAA TTCCTTTAGT TGTTCTTTT	1620
TATTCTCACT CCGCTGAAAC TGTTGAAAGT TGTTTAGCAA AACCCCATAC AGAAAAATCA	1680
TTTACTAACG TCTGGAAAGA CGACAAAAC TTAGATCGTT ACGCTAACTA TGAGGGTTGT	1740
CTGTGGAATG CTACAGGCGT TGTAAGTTGT ACTGGTGACG AAACCTCAGT TTACGGTACA	1800
TGGGTTCTTA TTGGGCTTGC TATCCCTGAA AATGAGGGTG GTGGCTCTGA GGGTGGCGGT	1860
TCTGAGGGTG GCGGTTCTGA GGGTGGCGGT ACTAAACCTC CTGAGTACGG TGATACACCT	1920
ATTCGGGGCT ATACTTATAT CAACCCTCTC GACGGCACTT ATCCGCCTGG TACTGAGCAA	1980
AACCCCGCTA ATCCTAATCC TTCTCTTGAG GAGTCTCAGC CTCTTAATAC TTTCATGTTT	2040
CAGAATAATA GGTTCGAAA TAGGCAGGG GCATTAACTG TTTATACGGG CACTGTACT	2100
CAAGGCACTG ACCCGTTAA AACTTATTAC CAGTACACTC CTGTATCATC AAAAGCCATG	2160
TATGACGCTT ACTGGAACGG TAAATTCAGA GACTGCGCTT TCCATTCTGG CTTTAATGAA	2220
GATCCATTCTG TTTGTGAATA TCAAGGCCAA TCGTCTGACC TGCCTCAACC TCCTGTCAAT	2280
GCTGGCGGCG GCTCTGGTGG TGGTCTGGT GCGGGCTCTG AGGGTGGTGG CTCTGAGGGT	2340
GGCGGTTCTG AGGGTGGCGG CTCTGAGGGA GCGGGTTCCG GTGGTGGCTC TGTTCCGGT	2400
GATTTTGATT ATGAAAAGAT GGCAAACGCT AATAAGGGGG CTATGACCGA AAATGCCGAT	2460
GAAAACGCGC TACAGTCTGA CGCTAAAGGC AAACCTTGATT CTGTGCTAC TGATTACGGT	2520
GCTGCTATCG ATGGTTTCAT TGGTGACGTT TCCGGCCTTG CTAATGGTAA TGGTGCTACT	2580
GGTGATTTTG CTGGCTCTAA TTCCGAAATG GCTCAAGTCG GTGACGGTGA TAATTCACCT	2640
TTAATGAATA ATTTCCGTCA ATATTACCT TCCCTCCCTC AATCGGTGA ATGTCGCCCT	2700
TTTGTCTTTA GCGCTGGTAA ACCATATGAA TTTTCTATTG ATTGTGACAA AATAAACTTA	2760
TTCCGTGGTG TCTTTGCGTT TCTTTTATAT GTTGCCACCT TTATGTATGT ATTTTCTACG	2820
TTTGCTAACA TACTGCGTAA TAAGGAGTCT TAATCATGCC AGTTCTTTTG GGTATTCGGT	2880

TATTATTGCG	TTTCCTCGGT	TTCCTTCTGG	TAACCTTGTT	CGGCTATCTG	CTTACTTTTC	2940
TTAAAAAGGG	CTTCGGTAAG	ATAGCTATTG	CTATTTTATT	GTTTCTTGCT	CTTATTATTG	3000
GGCTTAACTC	AATTCTTGTG	GGTTATCTCT	CTGATATTAG	CGCTCAATTA	CCCTCTGACT	3060
TTGTTCAAGG	TGTTCAAGTA	ATTCTCCCGT	CTAATGCGCT	TCCCTGTTTT	TATGTTATTG	3120
TCTCTGTAAA	GGCTGCTATT	TTCATTTTGG	ACGTTAAACA	AAAAATCGTT	TCTTATTTGG	3180
ATTGGGATAA	ATAATATGGC	TGTTTATTTT	GTAAGTGGCA	AATTAGGCTC	TGGAAAGACG	3240
CTCGTTAGCG	TTGGTAAGAT	TTAGGATAAA	ATTGTAGCTG	GGTGCAAAAT	AGCAACTAAT	3300
CTTGATTTAA	GGCTTCAAAA	CCTCCCGCAA	GTCGGGAGGT	TCGCTAAAAC	GCCTCGCGTT	3360
CTTAGAATAC	CGGATAAGCC	TTCTATATCT	GATTTGCTTG	CTATTGGGCG	CGGTAATGAT	3420
TCCTACGATG	AAAATAAAAA	CGGCTTGCTT	GTTCTCGATG	AGTGCGGTAC	TTGGTTTAAT	3480
ACCCGTTCTT	GGAATGATAA	GGAAAGACAG	CCGATTATTG	ATTGGTTTCT	ACATGCTCGT	3540
AAATTAGGAT	GGGATATTAT	CTTCCTTGTT	CAGGACTTAT	CTATTGTTGA	TAAACAGGCG	3600
CGTTCTGCAT	TAGCTGAACA	TGTTGTTTAT	TGTCGTCGTC	TGGACAGAAT	TACTTTACCT	3660
TTTGTCGGTA	CTTTATATTC	TCTTATTACT	GGCTCGAAAA	TGCCTCTGCC	TAAATTACAT	3720
GTTGGCGTTG	TTAAATATGG	CGATTCTCAA	TTAAGCCCTA	CTGTTGAGCG	TTGGCTTTAT	3780
ACTGGTAAGA	ATTTGTATAA	CGCATATGAT	ACTAAACAGG	CTTTTTCTAG	TAATTATGAT	3840
TCCGGTGTTT	ATTCTTATTT	AACGCCTTAT	TTATCAGACG	GTCGGTATTT	CAAACCATTA	3900
AATTTAGGTC	AGAAGATGAA	ATTAACATAA	ATATATTTGA	AAAAGTTTTC	TGCGGTTCTT	3960
TGCTTTCGCA	TTGGATTGTC	ATCAGCATTT	ACATATAGTT	ATATAACCCA	ACCTAAGCCG	4020
GAGGTAAAAA	AGGTAGTCTC	TCAGACCTAT	GATTTTGATA	AATTCAGTAT	TGACTCTTCT	4080
CAGCGTCTTA	ATCTAAGCTA	TCGCTATGTT	TTCAAGGATT	CTAAGGGAAA	ATTAATTAAT	4140
AGCGACGATT	TACAGAAGCA	AGGTTATTCA	CTCACATATA	TTGATTATATG	TACTGTTTCC	4200
ATTAAAAAAG	GTAATTCAAA	TGAAATTGTT	AAATGTAATT	AATTTTGTTT	TCTTGATGTT	4260
TGTTTCATCA	TCTTCTTTTG	CTCAGGTAAT	TGAAATGAAT	AATTGCGCTC	TGCGCGATTT	4320
TGTAACCTGG	TATTCAAAGC	AATCAGGCGA	ATCCGTTATT	GTTTCTCCCG	ATGTAAAAGG	4380
TACTGTTACT	GTATATTCAT	CTGACGTTAA	ACCTGAAAAT	CTACGCAATT	TCTTTATTTT	4440
TGTTTACGTT	GCTAATAATT	TTGATATGGT	TGGTTCAATT	CCTTCCATAA	TTCAGAAGTA	4500
TAATCCAAAC	AATCAGGATT	ATATTGATGA	ATTGCCATCA	TCTGATAATC	AGGAATATGA	4560
TGATAATTCC	GCTCCTTCTG	GTGGTTTCTT	TGTTCCGCAA	AATGATAATG	TTACTCAAAC	4620
TTTTAAAAAT	AATAACGTTT	GGGCAAAGGA	TTAATACGA	GTTGTCGAAT	TGTTTGTAAG	4680
GTCTAATACT	TCTAAATCCT	CAAATGTATT	ATCTATTGAC	GGCTCTAATC	TATTAGTTGT	4740
TAGTGCACCT	AAAGATATTT	TAGATAACCT	TCCTCAATTC	CTTCTACTG	TTGATTTGCC	4800
AACTGACCAG	ATATTGATTG	AGGGTTTGAT	ATTTGAGGTT	CAGCAAGGTG	ATGCTTTAGA	4860
TTTTTCATTT	GCTGCTGGCT	CTCAGCGTGG	CACTGTTGCA	GGCGGTGTTA	ATACTGACCG	4920

CCTCACCTCT	GTTTTATCTT	CTGCTGGTGG	TTCGTTCCGT	ATTTTAAATG	GGGATGTTTT	4980
AGGGCTATCA	GTTGGCGCAT	TAAAGACTAA	TAGCCATTCA	AAAATATTGT	CTGTGCCACG	5040
TATTCTTACG	CTTTCAGGTC	AGAAGGGTTC	TATCTCTGTT	GGCCAGAATG	TCCCTTTTAT	5100
TACTGGTCGT	GTGACTGGTG	AATCTGCCAA	TGTAAATAAT	CCATTTCAGA	CGATTGAGCG	5160
TCAAAATGTA	GGTATTTCCA	TGAGCGTTTT	TCCTGTTGCA	ATGGCTGGCG	GTAATATTGT	5220
TCTGGATATT	ACCAGCAAGG	CCGATAGTTT	GAGTTCCTCT	ACTCAGGCAA	GTGATGTTAT	5280
TACTAATCAA	AGAAGTATTG	CTACAACGGT	TAATTTGCGT	GATGGACAGA	CTCTTTTACT	5340
CGGTGGCCTC	ACTGATTATA	AAAACACTTC	TCAAGATTCT	GGCGTACCGT	TCCTGTCTAA	5400
AATCCCTTTA	ATCGGCCTCC	TGTTTAGCTC	CCGCTCTGAT	TCCAACGAGG	AAAGCACGTT	5460
ATACGTGCTC	GTCAAAGCAA	CCATAGTACG	CGCCCTGTAG	CGGCGCATT	AGCGCGGCGG	5520
GTGTGGTGGT	TACGCGCAGC	GTGACCGCTA	CACTTGCCAG	CGCCCTAGCG	CCCGCTCCTT	5580
TCGCTTTCTT	CCCTTCCTTT	CTCGCCACGT	TCGCCGGGCT	TCCCCGTCAA	GCTCTAAATC	5640
GGGGGCTCCC	TTTAGGGTTC	CGATTTAGTG	CTTTACGGCA	CCTCGACCCC	AAAAAACTTG	5700
ATTTGGGTGA	TGGTTCACGT	AGTGGGCCAT	CGCCCTGATA	GACGGTTTTT	CGCCCTTTGA	5760
CGTTGGAGTC	CACGTTCTTT	AATAGTGGAC	TCTTGTTCCA	AACTGGAACA	AACTCAACC	5820
CTATCTCGGG	CTATTCTTTT	GATTTATAAG	GGATTTTGCC	GATTTGCGAA	CCACCATCAA	5880
ACAGGATTTT	CGCCTGCTGG	GGCAAACCAG	CGTGGACCGC	TTGCTGCAAC	TCTCTCAGGG	5940
CCAGGGGGTG	AAGGGCAATC	AGCTGTTGCC	CGTCTCGCTG	GTGAAAAGAA	AAACCACCCT	6000
GGCGCCCAAT	ACGCAAACCG	CCTCTCCCCG	CGCGTTGGCC	GATTCATTAA	TGCAGCTGGC	6060
ACGACAGGTT	TCCCGACTGG	AAAGCGGGCA	GTGAGCGCAA	CGCAATTAAT	GTGAGTTAGC	6120
TCACTCATTA	GGCACCCAG	GCTTTACACT	TTATGCTTCC	GGCTCGTATG	TTGTGTGGAA	6180
TTGTGAGCGG	ATAACAATTT	CACACGCCAA	GGAGACAGTC	ATAATGAAAT	ACCTATTGCC	6240
TACGGCAGCC	GCTGGATTGT	TATTACTCGC	TGCCCAACCA	GCCATGGCCG	AGCTCGTGAT	6300
GACCCAGACT	CCAGAATTCC	ATCCGGAATG	AGTGTTAATT	CTAGAACGCG	TAAGCTTGGC	6360
ACTGGCCGTC	GTTTTACAAC	GTGCTGACTG	GGAAAACCCT	GGCGTTACCC	AACTTAATCG	6420
CCTTGCAGCA	CACCCCCCTT	TCGCCAGCTG	GCGTAATAGC	GAAGAGGCCC	GCACCGATCG	6480
CGCTTCCCAA	CAGTTGCGCA	GCCTGAATGG	CGAATGGCGC	TTTGCCTGGT	TTCCGGCACC	6540
AGAAGCGGTG	CCGGAAGCT	GGCTGGAGTG	CGATCTTCCT	GAGGCCGATA	CGGTCGTCGT	6600
CCCCTCAAAC	TGGCAGATGC	ACGGTTACGA	TGCGCCCATC	TACACCAACG	TAACCTATCC	6660
CATTACGGTC	AATCCGCCGT	TTGTTCCGAC	GGAGAATCCG	ACGGGTTGTT	ACTCGCTCAC	6720
ATTTAATGTT	GATGAAAGCT	GGCTACAGGA	AGCCAGACG	CGAATTATTT	TTGATGGCGT	6780
TCCTATTGGT	TAAAAAATGA	GCTGATTTAA	CAAAAATTTA	ACGCGAATTT	TAACAAAATA	6840
TTAACGTTTA	CAATTTAAAT	ATTTGCTTAT	ACAATCTTCC	TGTTTTTGGG	GCTTTTCTGA	6900
TTATCAACCG	GGGTACATAT	GATTGACATG	CTAGTTTTAC	GATTACCGTT	CATCGATTCT	6960

CTTGTTTGCT CCAGACTCTC AGGCAATGAC CTGATAGCCT TTGTAGATCT CTCAAAAATA	7020
GCTACCTCTT CCGGCATTAA TTTATCAGCT AGAACGGTTG AATATCATAT TGATGGTGAT	7080
TTGACTGTCT CCGGCCTTTT TCACCCTTTT GAATCTTTAC CTACACATTA CTCAGGCATT	7140
GCATTTAAAA TATATGAGGG TTCTAAAAAT TTTTATCCTT GCGTTGAAAT AAAGGCTTCT	7200
CCCGCAAAAG TATTACAGGG TCATAATGTT TTTGGTACAA CCGATTTAGC TTTATGCTCT	7260
GAGGCTTTAT TGCTTAATTT TGCTAATTCT TTGCCTTGCC TGTATGATTT ATTGGACGTT	7320

(2) INFORMATION FOR SEQ ID NO:3:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 7445 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: both
- (D) TOPOLOGY: circular

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

AATGCTACTA CTATTAGTAG AATTGATGCC ACCTTTTCAG CTCGCGCCCC AAATGAAAAT	60
ATAGCTAAAC AGGTTATTGA CCATTTCGGA AATGTATCTA ATGGTCAAAC TAAATCTACT	120
CGTTCGCAGA ATTGGGAATC AACTGTTACA TGGAATGAAA CTTCCAGACA CCGTACTTTA	180
GTTGCATATT TAAAACATGT TGAGCTACAG CACCAGATTC AGCAATTAAG CTCTAAGCCA	240
TCTGCAAAAA TGACCTCTTA TCAAAAGGAG CAATTAAAGG TACTCTCTAA TCCTGACCTG	300
TTGGAGTTTG CTTCCGGTCT GGTTCGCTTT GAAGCTCGAA TTAAAACGCG ATATTTGAAG	360
TCTTTCGGGC TTCCTCTTAA TCTTTTGTAT GCAATCCGCT TTGCTTCTGA CTATAAGT	420
CAGGGTAAAG ACCTGATTTT TGATTTATGG TCATTCTCGT TTTCTGAACT GTTTAAAGCA	480
TTTGAGGGGG ATTCAATGAA TATTTATGAC GATTCCGCAG TATTGGACGC TATCCAGTCT	540
AAACATTTTA CTATTACCCC CTCTGGCAAA ACTTCTTTTG CAAAAGCCTC TCGGTATTTT	600
GGTTTTTATC GTCGTCTGGT AAACGAGGGT TATGATAGTG TTGCTCTTAC TATGCCTCGT	660
AATTCCTTTT GCGGTATGT ATCTGCATTA GTTGAATGTG GTATTCCTAA ATCTCAACTG	720
ATGAATCTTT CTACCTGTAA TAATGTTGTT CCGTTAGTTC GTTTTATTAA CGTAGATTTT	780
TCTTCCCAAC GTCCTGACTG GTATAATGAG CCAGTTCTTA AAATCGCATA AGGTAATTCA	840
CAATGATTAA AGTTGAAATT AAACCATCTC AAGCCCAATT TACTACTCGT TCTGGTGT	900
CTCGTCAGGG CAAGCCTTAT TCACTGAATG AGCAGCTTTG TTACGTTGAT TTGGGTAATG	960
AATATCCGGT TCTTGTCAG ATTACTCTTG ATGAAGTCA GCCAGCCTAT GCGCCTGGTC	1020
TGTACACCGT TCATCTGTCC TCTTTCAAAG TTGGTCAGTT CCGTTCCCTT ATGATTGACC	1080
GTCTGCGCCT CGTTCCGGCT AAGTAACATG GAGCAGGTCG CGGATTTCTGA CACAATTTAT	1140
CAGGCGATGA TACAAATCTC CGTTGTACTT TGTTTCGGCG TTGGTATAAT CGCTGGGGGT	1200
CAAAGATGAG TGTTTTAGTG TATTCTTTCT CCTCTTTCGT TTTAGGTTGG TGCCTTCGTA	1260

GTGGCATTAC	GTATTTTACC	CGTTTAATGG	AAACTTCCTC	ATGAAAAAGT	CTTTAGTCCT	1320
CAAAGCCTCT	GTAGCCGTTG	CTACCCTCGT	TCCGATGCTG	TCTTTCGCTG	CTGAGGGTGA	1380
CGATCCCGCA	AAAGCGGCCT	TTAACTCCCT	GCAAGCCTCA	GCGACCGAAT	ATATCGGTTA	1440
TGCGTGGGCG	ATGGTTGTTG	TCATTGTGGG	CGCAACTATC	GGTATCAAGC	TGTTTAAGAA	1500
ATTCACCTCG	AAAGCAAGCT	GATAAACCGA	TACAATTAAA	GGCTCCTTTT	GGAGCCTTTT	1560
TTTTTGGAGA	TTTTC AACGT	GAAAAAATTA	TTATTCGCAA	TTCCTTTAGT	TGTTCCCTTC	1620
TATTCTCACT	CCGCTGAAAC	TGTTGAAAAGT	TGTTTAGCAA	AACCCCATAC	AGAAAAATCA	1680
TTTACTAACG	TCTGGAAAGA	CGACAAAAC	TTAGATCGTT	ACGCTAACTA	TGAGGGTGT	1740
CTGTGGAATG	CTACAGGCGT	TGTAGTTTGT	ACTGGTGACG	AAACTCAGTG	TTACGGTACA	1800
TGGGTTCCCTA	TTGGGCTTGC	TATCCCTGAA	AATGAGGGTG	GTGGCTCTGA	GGGTGGCGGT	1860
TCTGAGGGTG	GCGGTTCTGA	GGGTGGCGGT	ACTAAACCTC	CTGAGTACGG	TGATACACCT	1920
ATTCGGGGCT	ATACTTATAT	CAACCCTCTC	GACGGCACTT	ATCCGCCTGG	TACTGAGCAA	1980
AACCCCGCTA	ATCCTAATCC	TTCTCTTGAG	GAGTCTCAGC	CTCTTAATAC	TTTCATGTTT	2040
CAGAATAATA	GGTTCGAAA	TAGGCAGGGG	GCATTAACTG	TTTATACGGG	CACTGTTACT	2100
CAAGGCACTG	ACCCCGTTAA	AACTTATTAC	CAGTACACTC	CTGTATCATC	AAAAGCCATG	2160
TATGAGGCTT	ACTGGAACGG	TAAATTCAGA	GACTGCGCTT	TCCATTCTGG	CTTTAATGAA	2220
GATCCATTCC	TTTGTGAATA	TCAAGGCCAA	TCGTCTGACC	TGCCTCAACC	TCCTGTCAAT	2280
GCTGGCGGGG	GCTCTGGTGG	TGGTTCTGGT	GGCGGCTCTG	AGGGTGGTGG	CTCTGAGGGT	2340
GGCGGTTCTG	AGGGTGGCGG	CTCTGAGGGA	GGCGGTTCCG	GTGGTGGCTC	TGTTCCGGT	2400
GATTTTGATT	ATGAAAAGAT	GGCAAACGCT	AATAAGGGGG	CTATGACCGA	AAATGCCGAT	2460
GAAAACGCGC	TACAGTCTGA	CGCTAAAGGC	AAACTTGATT	CTGTGCTAC	TGATTACGGT	2520
GCTGCTATCG	ATGGTTTCAT	TGGTGACGTT	TCCGGCCTTG	CTAATGGTAA	TGGTGCTACT	2580
GGTGATTTTG	CTGGCTCTAA	TTCCCAAATG	GCTCAAGTCG	GTGACGGTGA	TAATTCACCT	2640
TTAATGAATA	ATTTCCGTCA	ATATTTACCT	TCCCTCCCTC	AATCGGTTGA	ATGTCGCCCT	2700
TTTGTCTTTA	GCGCTGGTAA	ACCATATGAA	TTTTCTATTG	ATTGTGACAA	AATAAACTTA	2760
TTCCGTGGTG	TCTTTCGCTT	TCTTTTATAT	GTTGCCACCT	TTATGTATGT	ATTTTCTACG	2820
TTTGCTAACA	TACTGCGTAA	TAAGGAGTCT	TAATCATGCC	AGTTCTTTTG	GGTATTCCGT	2880
TATTATTGCC	TTTCCTCGGT	TTCTTCTGG	TAACCTTGT	CGGCTATCTG	CTTACTTTTC	2940
TTAAAAAGGG	CTTCGGTAAG	ATAGCTATTG	CTATTTTATT	GTTTCTTGCT	CTTATTATTG	3000
GGCTTAACTC	AATTCTTG	GGTTATCTCT	CTGATATTAG	CGCTCAATTA	CCCTCTGACT	3060
TTGTTTCAAGG	TGTTTCAAGT	ATTCTCCCGT	CTAATGCGCT	TCCCTGTTTT	TATGTTATTG	3120
TCTCTGTAAA	GGCTGCTATT	TTTATTTTTG	ACGTAAACA	AAAAATCGTT	TCTTATTTGG	3180
ATTGGGATAA	ATAATATGGC	TGTTTATTTT	GTAAGTGGCA	AATTAGGCTC	TGGAAAGACG	3240
CTCGTTAGCG	TTGGTAAGAT	TCAGGATAAA	ATTGTAGCTG	GGTGCAAAAT	AGCAACTAAT	3300

CTTGATTTAA GGCTTCAAAA CCTCCGCAA GTCGGGAGGT TCGCTAAAAAC GCCTCGCGTT	3360
CTTAGAATAC CGGATAAGCC TTCTATATCT GATTTGCTTG CTATTGGGCG CGGTAATGAT	3420
TCCTACGATG AAAATAAAAA CGGCTTGCTT GTTCTCGATG AGTGGCGTAC TTGGTTTAAAT	3480
ACCCGTTCTT GGAATGATAA GGAAAGACAG CCGATTATTG ATTGGTTTCT ACATGCTCGT	3540
AAATTAGGAT GGGATATTAT TTTTCTTGTT CAGGACTTAT CTATTGTTGA TAAACAGGCG	3600
CGTTCTGCAT TAGCTGAACA TGTTGTTTAT TGTGTCGTC TGGACAGAAT TACTTTACCT	3660
TTTGTGGTA CTTTATATTG TCTTATTACT GGCTCGAAAA TGCCTCTGCC TAAATTACAT	3720
GTTGGCGTTG TTAAATATGG CGATTCTCAA TTAAGCCCTA CTGTTGAGCG TTGGCTTTAT	3780
ACTGGTAAGA ATTTGTATAA CGCATATGAT ACTAAACAGG CTTTTTCTAG TAATTATGAT	3840
TCCGGTGTTT ATTCTTATTT AACGCCTTAT TTATCACACG GTCGGTATTT CAAACCATTA	3900
AATTTAGGTC AGAAGATGAA GCTTACTAAA ATATATTTGA AAAAGTTTTT ACGCGTTCTT	3960
TGTCTTGCGA TTGGATTGCG ATCAGCATTG AGATATAGTT ATATAACCCA ACCTAAGCCG	4020
GAGGTAAAA AGGTAGTCTC TCAGACCTAT GATTTTGATA AATTCACTAT TGA CTCTCT	4080
CAGCGTCTTA ATCTAAGCTA TCGCTATGTT TTCAAGGATT CTAAGGGAAA ATTAATTAAT	4140
AGCGACGATT TACAGAAGCA AGGTTATTCA CTCACATATA TTGATTTATG TACTGTTTCC	4200
ATTAATAAAG GTAATTCAAA TGAAATTGTT AAATGTAATT AATTTTGTTT TCTTGATGTT	4260
TGTTTTCATCA TCTTCTTTTG CTCAGGTAAT TGAAATGAAT AATTCGCCTC TGCGCGATT	4320
TGTAACCTGG TATTCAAAGC AATCAGGCGA ATCCGTTATT GTTCTCTCCG ATGTAAAAGG	4380
TACTGTTACT GTATATTCAT CTGACGTAA ACCTGAAAAT CTACGCAATT TCTTTATTT	4440
TGTTTTACGT GCTAATAATT TTGATATGGT TGGTTCAATT CCTTCCATAA TTCAGAAGTA	4500
TAATCCAAAC AATCAGGATT ATATTGATGA ATTGCCATCA TCTGATAATC AGGAATATGA	4560
TGATAATTCC GCTCCTTCTG GTGGTTTCTT TGTTCCGCAA AATGATAATG TTA CTCAAAC	4620
TTTTAAATT AATAACGTTT GGGCAAAGGA TTTAATACGA GTTGTGCAAT TGTTTGTA	4680
GTCTAATACT TCTAAATCCT CAAATGTATT ATCTATTGAC GGCTCTAATC TATTAGTTGT	4740
TAGTGCACCT AAAGATATTT TAGATAACCT TCCTCAATTC CTTTCTACTG TTGATTTGCC	4800
AACTGACCAG ATATTGATTG AGGGTTTGAT ATTTGAGGTT CAGCAAGGTG ATGCTTTAGA	4860
TTTTTCATTT GCTGCTGGCT CTCAGCGTGG CACTGTTGCA GGCGGTGTTA ATACTGACCG	4920
CCTCACCTCT GTTTTATCTT CTGCTGGTGG TTCGTTGCGT ATTTTAAATG GCGATGTTT	4980
AGGGCTATCA GTTCGGCGAT TAAAGACTAA TAGCCATTCA AAAATATTGT CTGTGCCACG	5040
TATTCTTACG CTTTCAGGTC AGAAGGGTTC TATCTCTGTT GGCCAGAATG TCCCTTTTAT	5100
TACTGGTCGT GTGACTGGTG AATCTGCCAA TGTAATAAAT CCATTTGAGA CGATTGAGCG	5160
TCAAAATGTA GGTATTTCGA TGAGCGTTTT TCCTGTTGCA ATGGCTGGCG GTAATATTGT	5220
TCTGGATATT ACCAGCAAGG CCGATAGTTT GAGTTCTTCT ACTCAGGCAA GTGATGTTAT	5280
TACTAATCAA AGAAGTATTG CTACAACGGT TAATTTGCGT GATGGACAGA CTCTTTTACT	5340

CGGTGGCCTC	ACTGATTATA	AAAACACTTC	TCAAGATTCT	GGCGTACCGT	TCCTGTCTAA	5400
AATCCCTTTA	ATCGGCCTCC	TGTTTAGCTC	CGGCTCTGAT	TCCAACGAGG	AAAGCACGTT	5460
ATACGTGCTC	GTCAAAGCAA	CCATAGTACG	CGCCCTGTAG	CGGCGCATT	AGCGCGGCGG	5520
GTGTGGTGGT	TACGCGCAGC	GTGACCGCTA	CACTTGCCAG	CGCCCTAGCG	CCCGCTCCTT	5580
TCGCTTTCTT	CCCTTCCTTT	CTCGCCAGGT	TCGCCGGCTT	TCCCCGTCAA	GCTCTAAATC	5640
GGGGGCTCCC	TTTAGGGTTC	CGATTTAGTG	CTTTACGGCA	CCTCGACCCC	AAAAAACTTG	5700
ATTTGGGTGA	TGGTTCACGT	AGTGGGCCAT	CGCCCTGATA	GACGGTTTTT	CGCCCTTTGA	5760
CGTTGGAGTC	CACGTTCTTT	AATAGTGGAC	TCTTGTTCCT	AACTGGAACA	ACACTCAACC	5820
CTATCTCGGG	CTATTCTTTT	GATTTATAAG	GGATTTTGCC	GATTTTCGGAA	CCACCATCAA	5880
ACAGGATTTT	CGCCTGCTGG	GGCAAACCAG	CGTGGACCGC	TTGCTGCAAC	TCTCTCAGGG	5940
CCAGGCGGTG	AAGGGCAATC	AGCTGTTGCC	CGTCTCGCTG	GTGAAAAGAA	AAACCACCCT	6000
GGCGCCCAAT	ACGCAAACCG	CCTCTCCCGG	CGCGTTGGCC	GATTCATTAA	TGCAGCTGGC	6060
ACGACAGGTT	TCCCGACTGG	AAAGCGGGCA	GTGAGCGCAA	CGCAATTAAT	GTGAGTTAGC	6120
TCACTCATTA	GGCAGCCGAG	GCTTTAGACT	TTATGCTTCC	GGCTCGTATG	TTGTGTGGAA	6180
TTGTGAGCGG	ATAACAATTT	CACACGCGTC	ACTTGGCACT	GGCCGTCGTT	TTACAACGTC	6240
GTGACTGGGA	AAACCCTGGC	GTTACCCAAG	CTTTGTACAT	GGAGAAAATA	AAGTGAACA	6300
AAGCACTATT	GCACTGGCAC	TCTTACCGTT	ACCGTTACTG	TTTACCCCTG	TGACAAAAGC	6360
CGCCCAGGTC	CAGCTGCTCG	AGTGAGGCCT	ATTGTGCCCA	GGGGATTGTA	CTAGTGGATC	6420
CTAGGCTGAA	GGCGATGACC	CTGCTAAGGC	TGCATTCAAT	AGTTTACAGG	CAAGTGCTAC	6480
TGAGTACATT	GGCTACGCTT	GGGCTATGGT	AGTAGTTATA	GTTGGTGCTA	CCATAGGGAT	6540
TAAATTATTC	AAAAAGTTTA	CGAGCAAGGC	TTCTTAAGCA	ATAGCGAAGA	GGCCCGCACC	6600
GATCGCCCTT	CCCAACAGTT	GCGCAGCCTG	AATGGCGAAT	GGCGCTTTGC	CTGGTTTCCG	6660
GCACCAGAAG	CGGTGCCGGA	AAGCTGGCTG	GAGTGCGATC	TTCTGAGGC	CGATACGGTC	6720
GTCGTCCCTT	CAAACCTGGC	GATGCACGGT	TACGATGCGC	CCATCTACAC	CAACGTAACC	6780
TATCCCATTA	CGGTCAATCC	GCCGTTTGTT	CCCACGGAGA	ATCCGACGGG	TTGTTACTCG	6840
CTCACATTTA	ATGTTGATGA	AAGCTGGCTA	CAGGAAGGCC	AGACGCGAAT	TATTTTTGAT	6900
GGCGTTCCCTA	TTGGTTAAAA	AATGAGCTGA	TTTAACAAAA	ATTTAACCGG	AATTTTAACA	6960
AAATATTAAC	GTTTACAATT	TAAATATTTG	CTTATACAAT	CTTCCTGTTT	TTGGGGCTTT	7020
TCTGATTATC	AACCGGGGTA	CATATGATTG	ACATGCTAGT	TTTACGATTA	CCGTTTATCG	7080
ATTCTCTTGT	TTGCTCCAGA	CTCTCAGGCA	ATGACCTGAT	AGCCTTTGTA	GATCTCTCAA	7140
AAATAGCTAC	CCTCTCCGGC	ATTAATTTAT	CAGCTAGAAC	GGTTGAATAT	CATATTGATG	7200
GTGATTTGAC	TGTCTCCGGC	CTTTCTCACC	CTTTTGAATC	TTTACCTACA	CATTACTCAG	7260
GCATTGCATT	TAAAAATATAT	GAGGGTTCTA	AAAATTTTTA	TCCTTGCGTT	GAAATAAAGG	7320
CTTCTCCCGC	AAAAGTATTA	CAGGGTCATA	ATGTTTTTGG	TACAACCGAT	TTAGCTTTAT	7380

GCTCTGAGGC TTTATTGCTT AATTTTGCTA ATTCTTTGCC TTGCCTGTAT GATTTATTGG 7440
ACGTT 7445

(2) INFORMATION FOR SEQ ID NO:4:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 7409 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: both
 (D) TOPOLOGY: circular

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

AATGCTACTA CTATTAGTAG AATTGATGCC ACCTTTTCAG CTCGCGCCCC AAATGAAAAT 60
 ATAGCTAAAC AGGTTATTGA CCATTTGCCA AATGTATCTA ATGGTCAAAC TAAATCTACT 120
 CGTTCCGAGA ATTGGGAATC AACTGTTACA TGGAATGAAA CTTCCAGACA CCGTACTTTA 180
 GTTGCAATATT TAAAACATGT TGAGCTACAG CACCAGATTC AGCAATTAAG CTCTAAGCCA 240
 TCTGCAAAAA TGACCTCTTA TCAAAAGGAG CAATTAAAGG TACTCTCTAA TCCTGACCTG 300
 TTGGAGTTTG CTTCGGTCT GGTTCGCTTT GAAGCTCGAA TTAAAACGCG ATATTGAAG 360
 TCTTTCGGGC TCGCTCTTAA TCTTTTTGAT GCAATCCGCT TTGCTTCTGA CTATAATAGT 420
 CAGGGTAAAG ACCTGATTTT TGATTTATGG TCATTCTCGT TTTCTGAACT GTTTAAAGCA 480
 TTTGAGGGGG ATTCAATGAA TATTTATGAC GATTCCGCAG TATTGGACGC TATCCAGTCT 540
 AAACATTTTA CTATTACCCC CTCTGGCAA ACTTCTTTTG CAAAAGCCTC TCGCTATTTT 600
 GGTTTTTATC GTCGTCTGGT AAACGAGGGT TATGATAGTG TTGCTCTTAC TATGCCTCGT 660
 AATTCCTTTT GCGGTTATGT ATCTGCATTA GTTGAATGTG GTATTCCTAA ATCTCAACTG 720
 ATGAATCTTT CTACCTGTAA TAATGTTGTT CCGTTAGTTC GTTTTATTAA CGTAGATTTT 780
 TCTTCCGAAC GTCCTGACTG GTATAATGAG CCAGTTCTTA AAATCGCATA AGGTAATTCA 840
 CAATGATTAA AGTTGAAATT AAACCATCTC AAGCCCAATT TACTACTCGT TCTGGTGTTT 900
 CTCGTGAGGG CAAGCCTTAT TCACTGAATG AGCAGCTTTG TTACGTTGAT TTGGGTAATG 960
 AATATCCGGT TCTTGTCAAG ATTACTCTTG ATGAAGGTCA GCCAGCCTAT GCGCCTGGTC 1020
 TGTACACCGT TCATCTGTCC TCTTTCAAAG TTGGTCAGTT CCGTTCCTT ATGATTGACC 1080
 GTCTGCGCCT CGTTCCGGCT AAGTAACATG GAGCAGGTCG CGGATTTCTGA CACAATTTAT 1140
 CAGGCGATGA TACAAATCTC CGTTGTACTT TGTTTCGCGC TTGGTATAAT CGCTGGGGGT 1200
 CAAAGATGAG TGTTTTAGTG TATTCCTTCG CCTCTTTCGT TTAGGTTGG TGCCTTCGTA 1260
 GTGGCATTAC GTATTTTACC CGTTTAATGG AAACCTCCTC ATGAAAAAGT CTTTAGTCCT 1320
 CAAAGCCTCT GTAGCCGTTG CTACCCTCGT TCCGATGCTG TCTTTCGCTG CTGAGGGTGA 1380
 CGATCCCGCA AAAGCGGCCT TTAACCTCCT GCAAGCCTCA GCGACCGAAT ATATCGGTTA 1440
 TGGTGCGGCG ATGGTTGTTG TCATTGTCCG CGCAACTATC GGTATCAAGC TGTTTAAGAA 1500

ATTACCTCG	AAAGCAAGCT	GATAAACCGA	TACAATTAAA	GGCTCCTTTT	GGAGCCTTTT	1560
TTTTTGGAGA	TTTTC AACGT	GA AAAAATTA	TTATT CGCAA	TTCTTTTAGT	TGTTCCCTTC	1620
TATTCTCACT	CCGCTGAAAC	TGTTGAAAGT	TGTTTAGCAA	AACCCCATAC	AGAAAATTCA	1680
TTTACTAAG	TCTGGAAAGA	CGACAAAAC	TTAGATCGTT	ACGCTAACTA	TGAGGGTTGT	1740
CTGTGGAATG	CTACAGGCGT	TGTAGTTTGT	ACTGGTGACG	AAACTCAGTG	TTACGGTACA	1800
TGGGTTCCTA	TTGGGCTTGC	TATCCCTGAA	AATGAGGGTG	GTGGCTCTGA	GGGTGGCGGT	1860
TCTGAGGGTG	GCGGTTCTGA	GGGTGGCGGT	ACTAAACCTC	CTGAGTACGG	TGATACACCT	1920
ATTCCGGGCT	ATACTTATAT	CAACCCTCTC	GACGGCACTT	ATCCGCCTGG	TACTGAGCAA	1980
AACCCCGCTA	ATCCTAATCC	TTCTCTTGAG	GAGTCTCAGC	CTCTTAATAC	TTTCATGTTT	2040
CAGAATAATA	GGTTC CGAAA	TAGGCAGGGG	GCATTAAC	TTTATACGGG	CACTGTTACT	2100
CAAGGCACTG	ACCCCGTTAA	AACTTATTAC	CAGTACACTC	CTGTATCATC	AAAAGCCATG	2160
TATGACGCTT	ACTGGAACGG	TAAATTCAGA	GACTGGGCTT	TCCATTCTGG	CTTTAATGAA	2220
GATCCATTCC	TTTGTGAATA	TCAAGGCCAA	TGCTCTGACC	TGCCTCAACC	TCCTGTCAAT	2280
GCTGGCGGGG	GCTCTGGTGG	TGGTCTGGT	GGCGGCTCTG	AGGGTGGTGG	CTCTGAGGGT	2340
GGCGGTTCTG	AGGGTGGCGG	CTCTGAGGGA	GGCGGTTCCG	GTGGTGGCTC	TGGTTCGGGT	2400
GATTTTGATT	ATGAAAAGAT	GGCAAACGCT	AATAAGGGGG	CTATGACCGA	AAATGCCGAT	2460
GAAAACGGCG	TACAGTCTGA	CGCTAAAGGC	AACTTGATT	CTGTCGCTAC	TGATTACGGT	2520
GCTGCTATCG	ATGGTTTCAT	TGGTGACGTT	TCCGGCCTTG	CTAATGGTAA	TGGTGCTACT	2580
GGTGATTTTG	CTGGCTCTAA	TTCCCAAATG	GCTCAAGTCG	GTGACGGTGA	TAATTCACCT	2640
TTAATGAATA	ATTTCCGTCA	ATATTTACCT	TCCCTCCCTC	AATCGGTTGA	ATGTCGCCCT	2700
TTTGTCTTAA	GCGCTGGTAA	ACCATATGAA	TTTTCTATTG	ATTGTGACAA	AATAAACTTA	2760
TTCCGTGGTG	TCTTTGCGTT	TCTTTTATAT	GTTGCCACCT	TTATGTATGT	ATTTTCTACG	2820
TTTGCTAACA	TACTGCGTAA	TAAGGAGTCT	TAATCATGCC	AGTTCCTTTG	GGTATTCCGT	2880
TATTATTGCG	TTTCTCGGT	TTGCTTCTGG	TAACTTTGTT	CGGCTATCTG	CTTACTTTTC	2940
TTAAAAAGGG	CTTCGGTAAG	ATAGCTATTG	CTATTTCAAT	GTTTCTTGCT	CTTATTATTG	3000
GGCTTAACTC	AATTCTTG	GGTTATCTCT	CTGATATTAG	CGCTCAATTA	CCCTCTGACT	3060
TTGTTACAGG	TGTTACAGTTA	ATTCTCCCGT	CTAATGCGCT	TCCCTGTTTT	TATGTTATTC	3120
TCTCTGTAAA	GGCTGCTATT	TTCAATTTTG	ACGTTAAACA	AAAAATCGTT	TCTTATTTGG	3180
ATTGGGATAA	ATAATATGGC	TGTTTATTTT	GTAAC TGGCA	AATTAGGCTC	TGGAAAGACG	3240
CTCGTTAGCG	TTGGTAAGAT	TCAGGATAAA	ATTGTAGCTG	GGTGCAAAAT	AGCAACTAAT	3300
CTTGATTTAA	GGCTTCAAAA	CCTCCCGCAA	GTCGGGAGGT	TCGCTAAAAAC	GCCTCGCGTT	3360
CTTAGAATAC	CGGATAAGCC	TTCTATATCT	GATTTGCTTG	CTATTGGGCG	CGGTAATGAT	3420
TCCTACGATG	AAAATAAAAA	CGGCTTGCTT	GTTCTCGATG	AGTCCGGTAC	TTGTTTAAAT	3480
ACCCGTTCTT	GGAATGATAA	GGAAAGACAG	CCGATTATTG	ATTGGTTTCT	ACATGCTCGT	3540

AAATTAGGAT	GGGATATTAT	TTTTCTTGTT	CAGGACTTAT	CTATTGTTGA	TAAACAGGCG	3600
CGTTCTGCAT	TAGCTGAACA	TGTTGTTTAT	TGTCGTCGTC	TGGACAGAAT	TACTTTACCT	3660
TTTGTCGGTA	CTTTATATTC	TCTTATTACT	GGCTCGAAAA	TGCCTCTGCC	TAAATTACAT	3720
GTTGGCGTTG	TTAAATATGG	CGATTCTCAA	TTAAGCCCTA	CTGTTGAGCG	TTGGCTTTAT	3780
ACTGGTAAGA	ATTTGTATAA	CGCATATGAT	ACTAAACAGG	CTTTTTCTAG	TAATTATGAT	3840
TCCGGTGTTT	ATTCTTATTT	AACGCCTTAT	TTATCACACG	GTCGGTATTT	CAAACCATT	3900
AATTTAGGTC	AGAAGATGAA	GCTTACTAAA	ATATATTTGA	AAAAGTTTTC	ACGCGTTCCT	3960
TGTCTTGCGA	TTGGATTTGC	ATCAGCATT	ACATATAGTT	ATATAACCCA	ACCTAAGCCG	4020
GAGGTTAAAA	AGGTAGTCTC	TCAGACCTAT	GATTTTGATA	AATTCACTAT	TGACTCTTCT	4080
CAGCGTCTTA	ATCTAAGCTA	TCGCTATGTT	TTCAAGGATT	CTAAGGAAA	ATTAATTAAT	4140
AGCGACGATT	TACAGAAGCA	AGGTTATTCA	CTCACATATA	TTGATTTATG	TACTGTTTCC	4200
ATTAAAAAAG	GTAATTCAAA	TGAAATTGTT	AAATGTAATT	AATTTTGTTT	TCTTGATGTT	4260
TGTTTCATCA	TCTTCTTTTG	CTCAGGTAAT	TGAAATGAAT	AATTCGCCTC	TGCGCGATTT	4320
TGTAACCTGG	TATTCAAAGC	AATCAGGCGA	ATCCGTTATT	GTTTCTCCCG	ATGTAAAAGG	4380
TACTGTTACT	GTATATTCA	CTGACGTTAA	ACCTGAAAAT	CTACGCAATT	TCTTTATTTT	4440
TGTTTTACGT	GCTAATAATT	TTGATATGGT	TGTTCAATT	CCTTCCATAA	TTCAGAAGTA	4500
TAATCCAAAC	AATCAGGATT	ATATTGATGA	ATTGCCATCA	TCTGATAATC	AGGAATATGA	4560
TGATAATTCC	GCTCCTTCTG	GTGGTTTCTT	TGTTCCGCAA	AATGATAATG	TTACTCAAAC	4620
TTTTAAAATT	AATAACGTTT	GGGCAAAGGA	TTTAATACGA	GTTGTCGAAT	TGTTTGTA	4680
GTCTAATACT	TCTAAATCCT	CAAATGTATT	ATCTATTGAC	GGCTCTAATC	TATTAGTTGT	4740
TAGTGACCT	AAAGATATTT	TAGATAACCT	TCCTCAATTC	CTTTCTACTG	TTGATTTGCC	4800
AACTGACCAG	ATATTGATTG	AGGGTTTGAT	ATTGAGGTT	CAGCAAGGTG	ATGCTTTAGA	4860
TTTTTCATTT	GCTGCTGGCT	CTCAGCGTGG	CACTGTTGCA	GGCGGTGTTA	ATACTGACCG	4920
CCTCACCTCT	GTTTTATCTT	CTGCTGGTGG	TTCGTTCCGGT	ATTTTAAATG	GCGATGTTTT	4980
AGGGCTATCA	GTTGCGGCAT	TAAAGACTAA	TAGCCATTCA	AAAATATTGT	CTGTGCCACG	5040
TATTCTTACG	CTTTCAGGTC	AGAAGGGTTC	TATCTCTGTT	GGCCAGAATG	TCCCTTTTAT	5100
TACTGGTCGT	GTGACTGGTG	AATCTGCCAA	TGTAAATAAT	CCATTTTACA	CGATTGAGCG	5160
TCAAAATGTA	GGTATTTCCA	TGAGCGTTTT	TCCTGTTGCA	ATGGCTGGCG	GTAATATTGT	5220
TCTGGATATT	ACCAGCAAGG	CCGATAGTTT	GAGTTCTTCT	ACTCAGGCAA	GTGATGTTAT	5280
TACTAATCAA	AGAAGTATTG	CTACAACGGT	TAATTTGCGT	GATGGACAGA	CTCTTTTACT	5340
CGGTGGCCTC	ACTGATTATA	AAAACACTTC	TCAAGATTCT	GGCGTACCGT	TCCTGTCTAA	5400
AATCCCTTTA	ATCGGCCTCC	TGTTTAGCTC	CCGCTCTGAT	TCCAACGAGG	AAAGCACGTT	5460
ATACGTGCTC	GTCAAAGCAA	CCATAGTACG	CGCCCTGTAG	CGGCGCATT	AGCGCGGCGG	5520
GTGTGGTGGT	TACGCGCAGC	GTGACCGCTA	CACTTGCCAG	CGCCCTAGCG	CCCGCTCCTT	5580

TCGCTTTCTT CCCTTCCTTT CTCGCCACGT TCGCCGGCTT TCCCGTCAA GCTCTAAATC	5640
GGGGGCTCCC TTTAGGGTTC CGATTTAGTG CTTTACGGCA CCTCGACCCC AAAAACTTG	5700
ATTTGGGTGA TGGTTCACGT AGTGGGCCAT CGCCCTGATA GACGGTTTTT CGCCCTTGA	5760
CGTIGGAGTC CACGTTCTTT AATAGTGGAC TCTGTTCGA AACTGGAACA AACTCAACC	5820
CTATCTCGGG CTATTCTTTT GATTTATAAG GGATTTTGCC GATTTGGA CCACCATCAA	5880
ACAGGATTTT CGCCTGCTGG GGCAAACCAG CGTGGACCGC TTGCTGCAAC TCTCTGAGG	5940
CCAGGCGGTG AAGGGCAATC AGCTGTTGCC CGTCTCGCTG GTGAAAAGAA AAACCACCT	6000
GGCGCCCAAT ACGCAAACCG CCTCTCCCCG CGCGTTGGCC GATTCATTAA TGCAGCTGGC	6060
ACGACAGGTT TCCCGACTGG AAAGCGGGCA GTGAGCGCAA CGCAATTAAT GTGAGTTAGC	6120
TCACTCATTG GGCACCCAG GCTTTACACT TTATGCTTCC GGCTCGTATG TTGTGTGGAA	6180
TTGTGAGCGG ATAACAATTT CACACGCGTC ACTTGGCACT GGCCGTCGTT TTACAACGTC	6240
GTGACTGGGA AAACCCTGGC GTTACCCAAG CTTTGTACAT GGAGAAAATA AAGTGAAACA	6300
AAGCACTATT GCACTGGCAC TCTTACCGTT ACTGTTTACC CCTGTGGCAA AAGCCTATGG	6360
GGGGTTTATG ACTTCTGAGG GATCCGGAGC TGAAGGCGAT GACCCTGCTA AGGCTGCATT	6420
CAATAGTTTA CAGGCAAGTG CTAAGTGA TAATTGGCTAC GCTTGGGCTA TGGTAGTAGT	6480
TATAGTTGGT GCTACCATAG GGATTAATTT ATTCAAAAAG TTTACGAGCA AGGCTTCTTA	6540
AGCAATAGCG AAGAGGCCCG CACCGATCGC CCTTCCCAAC AGTTGCGCAG CCTGAATGGC	6600
GAATGGCGCT TTGCCTGGTT TCCGGCACCA GAAGCGGTGC CGGAAAGCTG GCTGAGTGC	6660
GATCTTCCTG AGGCCGATAC GGTGTCGTC CCCTCAAAC GGCAGATGCA CGGTTACGAT	6720
GCGCCCATCT ACACCAACGT AACCTATCCC ATTACGGTCA ATCCGCCGTT TGTTCCACG	6780
GAGAATCCGA CGGGTTGTTA CTCGCTCACA TTAAATGTTG ATGAAAGCTG GCTACAGGAA	6840
GGCCAGACGC GAATTATTTT TGATGGCGTT CCTATTGGTT AAAAAATGAG CTGATTTAAC	6900
AAAAATTTAA CGCGAATTTT AACAAAATAT TAACGTTTAC AATTTAAATA TTTGCTTATA	6960
CAATCTTCCT GTTTTGGGG CTTTCTGAT TATCAACCGG GGTACATATG ATTGACATGC	7020
TAGTTTACG ATTACCGTTC ATCGATTCTC TTGTTTGCTC CAGACTCTCA GGCAATGACC	7080
TGATAGCCTT TGTAGATCTC TCAAAAATAG CTACCCTCTC CGGCATTAAT TTATCAGCTA	7140
GAACGGTTGA ATATCATATT GATGGTGATT TGAAGTCTC GGGCCTTTCT CACCTTTTG	7200
AATCTTTACC TACACATTAC TCAGGCATTG CATTTAAAAT ATATGAGGGT TCTAAAAATT	7260
TTTATCCTTG CGTTGAAATA AAGGCTTCTC CCGCAAAAGT ATTACAGGGT CATAATGTTT	7320
TTGGTACAAC CGATTTAGCT TTATGCTCTG AGGCTTTATT GCTTAATTTT GCTAATTCTT	7380
TGCCTTGCCT GTATGATTTA TTGGACGTT	7409

(2) INFORMATION FOR SEQ ID NO:5:

- (1) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 7294 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: both
 (D) TOPOLOGY: circular

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

AATGCTACTA CTATTAGTAG AATTGATGCC ACCTTTTCAG CTCGCGCCCC AAATGAAAAT	60
ATAGCTAAAC AGGTTATTGA CCATTGCGGA AATGTATCTA ATGGTCAAAC TAAATCTACT	120
CGTTCCGAGA ATTGGGAATC AACTGTTACA TGAATGAAA CTTCCAGACA CCGTACTTTA	180
GTTGCATATT TAAAACATGT TGAGCTACAG CACCAGATTC AGCAATTAAG CTCTAAGCCA	240
TCTGCAAAAA TGACCTCTTA TCAAAGGAG CAATTAAAGG TACTCTCTAA TCCTGACCTG	300
TTGGAGTTTG CTTCCGGTCT GGTTCGCTTT GAAGCTCGAA TTAAAACGCG ATATTTGAAG	360
TCTTTCGGGC TTCCTCTTAA TCTTTTGAT GCAATCCGCT TTGCTTCTGA CTATAATAGT	420
CAGGGTAAAG ACCTGATTTT TGATTTATGG TCATTCTCGT TTTCTGAACT GTTTAAAGCA	480
TTTGAGGGGG ATTCAATGAA TATTTATGAC GATTCCGCAG TATTGGACGC TATCCAGTCT	540
AAACATTTTA CTATTACCCC CTCTGGCAA ACTTCTTTTG CAAAAGCCTC TCGCTATTTT	600
GGTTTTTATC GTCGTCTGGT AAACGAGGGT TATGATAGTG TTGCTCTTAC TATGCCTCGT	660
AATTCCTTTT GCGGTTATGT ATCTGCATTA GTTGAATGTG GTATTCCTAA ATCTCAACTG	720
ATGAATCTTT CTACCTGTAA TAATGTTGTT CCGTTAGTTC GTTTTATTAA CGTAGATTTT	780
TCTTCCGAAC GTCCTGACTG GTATAATGAG CCAGTTCTTA AAATCGCATA AGGTAATTCA	840
CAATGATTAA AGTTGAAATT AAACCATCTC AAGCCCAATT TACTACTCGT TCTGGTGTTC	900
CTCGTCAGGG CAAGCCTTAT TCACTGAATG AGCAGCTTTG TTACGTTGAT TTGGGTAATG	960
AATATCCGGT TCTTGTCAG ATTACTCTTG ATGAAGGTCA GCCAGCCTAT GCGCCTGGTC	1020
TGTACACCGT TCATCTGTCC TCTTTCAAAG TTGGTCAGTT CGGTTCCCTT ATGATTGACC	1080
GTCTGGGCCT CGTTCCGGCT AAGTAACATG GAGCAGGTGG CGGATTTCCA CACAATTTAT	1140
CAGGCGATGA TACAAATCTC CGTTGTACTT TGTTTCGCGC TTGGTATAAT CGCTGGGGGT	1200
CAAAGATGAG TGTTTTAGTG TATTCTTTTG CCTCTTTCTG TTTAGGTTGG TGCCTTCGTA	1260
GTGGCATTAC GTATTTTACC CGTTTAATGG AAACCTCCTC ATGAAAAAGT CTTTAGTCCT	1320
CAAAGCCTCT GTAGCCGTTG CTACCCTCGT TCCGATGCTG TCTTTCGCTG CTGAGGGTGA	1380
CGATCCGCA AAAGCGGCCT TTAACCTCCT GCAAGCCTCA GCGACCGAAT ATATCGGTTA	1440
TGCGTGGGCG ATGGTTGTTG TCATTGTCGG CGCAACTATC GGTATCAAGC TGTTTAAGAA	1500
ATTACCTCG AAAGCAAGCT GATAAACCGA TACAATTAAA GGCTCCTTTT GGAGCCTTTT	1560
TTTTTGAGA TTTTCAACGT GAAAAAATTA TTATTCGCAA TTCCTTTAGT TGTTCCTTTC	1620
TATTCTCACT CCGCTGAAAC TGTTGAAAGT TGTTTAGCAA AACCCCATAC AGAAAAATTCA	1680

TTTACTAACG	TCTGGAAAGA	CGACAAAAC	TTAGATCGTT	ACGCTAACTA	TGAGGGTTGT	1740
CTGTGGAATG	CTACAGGCGT	TGTAGTTTGT	ACTGGTGACG	AAACTCAGTG	TTACGGTACA	1800
TGGGTTCCCTA	TTGGGCTTGC	TATCCCTGAA	AATGAGGGTG	GTGGCTCTGA	GGGTGGCGGT	1860
TCTGAGGGTG	GCGGTTCTGA	GGGTGGCGGT	ACTAAACCTC	CTGAGTACGG	TGATACACCT	1920
ATTCCGGGCT	ATACTTATAT	CAACCCTCTC	GACGGCACTT	ATCCGCCTGG	TACTGAGCAA	1980
AACCCCGCTA	ATCCTAATCC	TTCTCTTGAG	GAGTCTCAGC	CTCTTAATAC	TTTCATGTTT	2040
CAGAATAATA	GGTTCGAAA	TAGGCAGGGG	GCATTAAC	TTTATACGGG	CACTGTTACT	2100
CAAGGCACTG	ACCCCGTTAA	AACTTATTAC	CAGTACACTC	CTGTATCATC	AAAAGCCATG	2160
TATGACGCTT	ACTGGAACGG	TAAATTGAGA	GACTGCGCTT	TCCATTCTGG	CTTTAATGAA	2220
GATCCATTCTG	TTTGTGAATA	TCAAGGCCAA	TCGTCTGACC	TGCCTCAACC	TCCTGTCAAT	2280
GCTGGCGGCG	GCTCTGGTGG	TGGTCTGGT	GGCGGCTCTG	AGGGTGGTGG	CTCTGAGGGT	2340
GGCGGTTCTG	AGGGTGGCGG	CTCTGAGGGA	GGCGGTTCCG	GTGGTGGCTC	TGGTTCGGGT	2400
GATTTTGATT	ATGAAAAGAT	GGCAAACGCT	AATAAGGGGG	CTATGACCGA	AAATGCCGAT	2460
GAAGACGCGC	TACAGTCTGA	CGCTAAAGGC	AAACTTGATT	CTGTGCTAC	TGATTACGGT	2520
GCTGCTATCG	ATGGTTTCAT	TGGTGACGTT	TCCGGCCTTG	CTAATGGTAA	TGGTGCTACT	2580
GGTGATTTTG	CTGGCTCTAA	TTCCCAAATG	GCTCAAGTCG	GTGACGGTGA	TAATTCACCT	2640
TTAATGAATA	ATTTCCGTCA	ATATTTACCT	TCCCTCCCTC	AATCGGTTGA	ATGTCGCCCT	2700
TTTGTCTTTA	GCGCTGGTAA	ACCATATGAA	TTTCTATTG	ATTGTGACAA	AATAAACTTA	2760
TTCCGTGGTG	TCTTTGCGTT	TCTTTTATAT	GTTGCCACCT	TTATGTATGT	ATTTTCTACG	2820
TTTGCTAACA	TACTGCGTAA	TAAGGAGTCT	TAATCATGCC	AGTTCTTTTG	GGTATTCCGT	2880
TATTATTGCG	TTTCCTCGGT	TTCTTCTGG	TAACTTTGTT	CGGCTATCTG	CTTACTTTTC	2940
TTAAAAAGGG	CTTCGGTAAG	ATAGCTATTG	CTATTTTCATT	GTTTCTTGCT	CTTATTATTG	3000
GGCTTAACTC	AATTCTTG	GGTTATCTCT	CTGATATTAG	CGCTCAATTA	CCCTCTGACT	3060
TTGTTGAGGG	TGTTGAGTTA	ATTCTCCCGT	CTAATGCGCT	TCCCTGTTTT	TATGTTATTC	3120
TCTCTGTAAA	GGCTGCTATT	TTTATTTTGG	ACGTAAACA	AAAAATCGTT	TCTTATTTGG	3180
ATTGGGATAA	ATAATATGGC	TGTTTATTTT	GTAAGTGGCA	AATTAGGCTC	TGGAAAGACG	3240
CTCGTTAGCG	TTGGTAAGAT	TCAGGATAAA	ATTGTAGCTG	GGTGCAAAAT	AGCAACTAAT	3300
CTTGATTTAA	GGCTTCAAAA	CCTCCCGCAA	GTGCGGAGGT	TGCTAAAAC	GCCTCGCGTT	3360
CTTAGAATAC	CGGATAAGCC	TTCTATATCT	GATTTGCTTG	CTATTGGGCG	CGGTAATGAT	3420
TCCTACGATG	AAAATAAAAA	CGGCTTGCTT	GTTCTCGATG	AGTGCGGTAC	TTGGTTTAAT	3480
ACCCGTTCTT	GGAATGATAA	GGAAAGACAG	CCGATTATTG	ATTGGTTTCT	ACATGCTCGT	3540
AAATTAGGAT	GGGATATTAT	CTTCCTTGTT	CAGGACTTAT	CTATTGTTGA	TAAACAGGCG	3600
CGTTCTGCAT	TAGCTGAACA	TGTTGTTTAT	TGTCGTCGTC	TGGACAGAAT	TACTTTACCT	3660
TTTGTGCGTA	CTTTATATTG	TCTTATTACT	GGCTCGAAAA	TGCCTCTGCC	TAAATTACAT	3720

GTGGCGTTG	TTAAATATGG	CGATTCTCAA	TTAAGCCCTA	CTGTTGAGCG	TTGGCTTTAT	3780
ACTGGTAAGA	ATTTGTATAA	CGCATATGAT	ACTAAACAGG	CTTTTTCTAG	TAATTATGAT	3840
TCCGGTGTTT	ATTCTTATTT	AACGCCTTAT	TTATCACACG	GTCGGTATTT	CAAACCATTA	3900
AATTTAGGTC	AGAAGATGAA	GCTTACTAAA	ATATATTTGA	AAAAGTTTTC	ACGCGTTCTT	3960
TGTCTTGCGA	TTGGATTTGC	ATCAGCATTT	ACATATAGTT	ATATAACCGA	ACCTAAGCCG	4020
GAGGTTAAAA	AGGTAGTCTC	TCAGACCTAT	GATTTTGATA	AATTCACAT	TGACTCTTCT	4080
CAGCGTCTTA	ATCTAAGCTA	TGGCTATGTT	TTCAAGGATT	CTAAGGAAAA	ATTAATTAAT	4140
AGCGACGATT	TACAGAAGCA	AGGTTATTCA	CTCACATATA	TTGATTTATG	TACTGTTTCC	4200
ATTAAAAAGG	TAATTCAAAT	GAAATTGTTA	AATGTAATTA	ATTTTGTTTT	CTTGATGTTT	4260
GTTTCATCAT	CTTCTTTTGC	TCAGGTAATT	GAAATGAATA	ATTGGCCTCT	GCGCGATTTT	4320
GTAAC TTGGT	ATTCAAAGCA	ATCAGGCGAA	TCCGTTATTG	TTTCTCCCGA	TGTA AAAGGT	4380
ACTGTTACTG	TATATTCATC	TGACGT TAAA	CCTGAAAATC	TACGCAATTT	CTTTATTTCT	4440
GTTTTACGTG	CTAATAATTT	TGATATGGTT	GGTTC AATTC	CTTCCATTAT	TTAGAAGTAT	4500
AATCCAAACA	ATCAGGATTA	TATTGATGAA	TTGCCATCAT	CTGATAATCA	GGAATATGAT	4560
GATAATTCCG	CTCCTTCTGG	TGGTTTCTTT	GTTCCGCAAA	ATGATAATGT	TACTCAA ACT	4620
TTTAA AATTA	ATAACGTT CG	GGCAAAGGAT	TTAATACGAG	TTGT CGAATT	GTTTGTAAG	4680
TCTAATACTT	CTAAATCCTC	AAATGTATTA	TCTATTGACG	GCTCTAATCT	ATTAGTTGTT	4740
AGTGCACCTA	AAGATATTTT	AGATAACCTT	CCTCAATTCC	TTTCTACTGT	TGATTTGCCA	4800
ACTGACCAGA	TATTGATTGA	GGGTTTGATA	TTTGAGGTTT	AGCAAGGTGA	TGCTTTAGAT	4860
TTTTCATTTG	CTGCTGGCTC	TCAGCGTGGC	ACTGTTGCAG	GCGGTGTTAA	TACTGACCGC	4920
CTCACCTCTG	TTTTATCTTC	TGCTGGTGGT	TGTTTCGGTA	TTTTTAATGG	CGATGTTTTA	4980
GGGCTATCAG	TTCGCGCATT	AAAGACTAAT	AGCCATTCAA	AAATATTGTC	TGTGCCACGT	5040
ATTCTTACGC	TTTCAGGTCA	GAAGGGTTCT	ATCTCTGTTG	GCCAGAATGT	CCCTTTTATT	5100
ACTGGTCGTG	TGACTGGTGA	ATCTGCCAAT	GTAATAATC	CATTT CAGAC	GATTGAGCGT	5160
CAAAATGTAG	GTATTTCCAT	GAGCGTTTTT	CCTGTTGCAA	TGGCTGGCGG	TAATATTGTT	5220
CTGGATATTA	CCAGCAAGGC	CGATAGTTTG	AGTTCTTCTA	CTCAGGCAAG	TGATGTTATT	5280
ACTAATCAAA	GAAGTATTGC	TACAACGGTT	AATTTGCGTG	ATGGACAGAC	TCTTTTACTC	5340
GGTGGCCTCA	CTGATTATAA	AAACACTTCT	CAAGATTCTG	GCGTACCGTT	CCTGTCTAAA	5400
ATCCCTTTAA	TCGGCCTCCT	GTTTAGCTCC	CGCTCTGATT	CCAACGAGGA	AAGCACGTTA	5460
TACGTGCTCG	TCAAAGCAAC	CATAGTACGC	GCCCTGTAGC	GGCGCATTAA	GCGCGGCGGG	5520
TGTGGTGGTT	ACGCGCAGCG	TGACCGCTAC	ACTTGCCAGC	GCCCTAGCGC	CCGCTCCTTT	5580
CGCTTTCTTC	CCTTCCTTTC	TCGCCACGTT	CGCCGGCTTT	CCCGTCAAG	CTCTAAATCG	5640
GGGGCTCCCT	TTAGGGTTCC	GATTTAGTGC	TTTACGGCAC	CTCGACCCCA	AAAACTTGA	5700
TTTGGGTGAT	GGTTCACGTA	GTGGGCCATC	GCCCTGATAG	ACGGTTTTTC	GCCCTTTGAC	5760

GTTGGAGTCC	ACGTTCTTTA	ATAGTGGACT	CTTGTTCCAA	ACTGGAACAA	CACTCAACCC	5820
TATCTCGGGC	TATTCCTTTG	ATTTATAAGG	GATTTTGCCG	ATTCGGAAC	CACCATCAAA	5880
CAGGATTTTC	GCCTGCTGGG	GCAAACCAGC	GTGGACCGCT	TGCTGCAACT	CTCTCAGGGC	5940
CAGGCGGTGA	AGGGCAATCA	GCTGTTGCCC	GTCTCGCTGG	TGAAAAGAAA	AACCACCCTG	6000
GCGGCCAATA	CGCAAACCGC	CTCTCCCCGC	GCGTTGGCCG	ATTCATTAAT	GCAGCTGGCA	6060
CGACAGGTTT	CCCGACTGGA	AAGCGGGCAG	TGAGCGCAAC	GCAATTAATG	TGAGTTAGCT	6120
CACTCATTAG	GCACCCAGG	CTTTACACTT	TATGCTTCCG	GCTCGTATGT	TGTGTGGAAT	6180
TGTGAGCGGA	TAACAATTTT	ACACAGGAAA	CAGCTATGAC	CAGGATGTAC	GAATTCGCAG	6240
GTAGGAGAGC	TGGGCGGATC	CGAGGCTGAA	GGCGATGACC	CTGCTAAGGC	TGCATTCAAT	6300
AGTTTACAGG	CAAGTGCTAC	TGAGTACATT	GGCTACGCTT	GGGCTATGGT	AGTAGTTATA	6360
GTTGGTGCTA	CCATAGGGAT	TAAATTATTC	AAAAAGTTTA	CGAGCAAGGC	TTCTTAACCA	6420
GCTGGCGTAA	TAGCGAAGAG	GCCCGCACCG	ATCGCCCTTC	CCAACAGTTG	CGCAGCCTGA	6480
ATGGCGAATG	GCGCTTTGCC	TGGTTTCCGG	CACCAGAAGC	GGTGCCGGAA	AGCTGGCTGG	6540
AGTGCGATCT	TCCTGAGGCC	GATACGGTCG	TCGTCCCTTC	AAACTGGCAG	ATGCACGTTT	6600
ACGATGCGCC	CATCTACACC	AACGTAACCT	ATCCCATTAC	GGTCAATCCG	CCGTTTGTTT	6660
CCACGGAGAA	TCCGACGGGT	TGTTACTCGC	TCACATTTAA	TGTTGATGAA	AGCTGGCTAC	6720
AGGAAGGCCA	GACGCCAATT	ATTTTTGATG	GCGTTCCTAT	TGGTTAAAAA	ATGAGCTGAT	6780
TTAACAAAAA	TTTAACGCGA	ATTTTAACAA	AATATTAACG	TTTACAATTT	AAATATTTGC	6840
TTATACAATC	TTCTGTTTTT	TGGGGCTTTT	CTGATTATCA	ACCGGGGTAC	ATATGATTGA	6900
CATGCTAGTT	TTACGATTAC	CGTTCATCGA	TTCTCTTGTT	TGCTCCAGAC	TCTCAGGCAA	6960
TGACCTGATA	GCCTTTGTAG	ATCTCTCAAA	AATAGCTACC	CTCTCGGGCA	TTAATTTATC	7020
AGCTAGAACG	GTTGAATATC	ATATTGATGG	TGATTTGACT	GTCTCGGGCC	TTTCTCACCC	7080
TTTTGAATCT	TTACCTACAC	ATTACTCAGG	CATTGCATT	AAAATATATG	AGGGTTCTAA	7140
AAATTTTAT	CCTTGCGTTG	AAATAAAGGC	TTCTCCCGCA	AAAGTATTAC	AGGGTCATAA	7200
TGTTTTTGGT	ACAACCGATT	TAGCTTTATG	CTCTGAGGCT	TTATTGCTTA	ATTTTGCTAA	7260
TTCTTTGCCT	TGCCTGTATG	ATTTATTGGA	CGTT			7294

(2) INFORMATION FOR SEQ ID NO:6:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 7394 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: both
- (D) TOPOLOGY: circular

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

AATGCTACTA CTATTAGTAG AATTGATGCC ACCTTTTCAG CTCGCGCCCC AAATGAAAAT 60

ATAGCTAAAC AGGTTATTGA CCATTTGCCA AATGTATCTA ATGGTCAAAC TAAATCTACT	120
CGTTCGCAGA ATTGGGAATC AACTGTTACA TGGAATGAAA CTTCCAGACA CCGTACTTTA	180
GTTGCATATT TAAAACATGT TGAGCTACAG CACCAGATTC AGCAATTAAG CTCTAAGCCA	240
TCTGCAAAAA TGACCTCTTA TCAAAAGGAG CAATTAAAGG TACTCTCTAA TCCTGACCTG	300
TTGGAGTTTG CTTCCGGTCT GGTTCGCTTT GAAGCTCGAA TTAAAACGCG ATATTTGAAG	360
TCTTTCCGGC TTCCTCTTAA TCTTTTTGAT GCAATCCGCT TTGCTTCTGA CTATAATAGT	420
CAGGGTAAAG ACCTGATTTT TGATTTATGG TCATTCTCGT TTTCTGAACT GTTTAAAGCA	480
TTTGAGGGGG ATTCAATGAA TATTTATGAC GATTCCGCAG TATTGGACGC TATCCAGTCT	540
AAACATTTTA CTATTACCCC CTCTGGCAA ACTTCTTTTG CAAAAGCCTC TCGCTATTTT	600
GGTTTTTATC GTCGTCTGGT AAACGAGGGT TATGATAGTG TTGCTCTTAC TATGCCTCGT	660
AATTCCTTTT GCGGTTATGT ATCTGCATTA GTTGAATGTG GTATTCCTAA ATCTCAACTG	720
ATGAATCTTT CTACCTGTAA TAATGTTGTT CCGTTAGTTC GTTTTATTAA CGTAGATTTT	780
TCTTCCCAAC GTCCTGACTG GTATAATGAG CCAGTTCTTA AAATCGCATA AGGTAATTCA	840
CAATGATTAA AGTTGAAATT AAACCATCTC AAGCCCAATT TACTACTCGT TCTGGTGTTC	900
CTCGTCAGGG CAAGCCTTAT TCACTGAATG AGCAGCTTTG TTACGTTGAT TTGGGTAATG	960
AATATCCGGT TCTTGTCAAG ATTACTCTTG ATGAAGGTCA GCCAGCCTAT GCGCCTGGTC	1020
TGTACACCGT TCATCTGTCC TCTTTCAAAG TTGGTCAGTT CGGTTCCCTT ATGATTGACC	1080
GTCTGCGCCT CGTTCGGGCT AAGTAACATG GAGCAGGTCG CGGATTTCGA CACAATTTAT	1140
CAGGCGATGA TACAAATCTC CGTTGTACTT TGTTTTCGCGC TTGGTATAAT CGCTGGGGGT	1200
CAAAGATGAG TGTTTTAGTG TATTCCTTCG CCTCTTTCGT TTTAGGTTGG TGCCTTCGTA	1260
GTGGCATTAC GTATTTTACC CGTTTAATGG AAACCTCCTC ATGAAAAAGT CTTTAGTCCT	1320
CAAAGCCTCT GTAGCCGTTG CTACCCTCGT TCCGATGCTG TCTTTCGCTG CTGAGGGTGA	1380
CGATCCCGCA AAAGCGGCCT TTAACCTCCT GCAAGCCTCA GCGACCGAAT ATATCGGTTA	1440
TGCGTGGGCG ATGGTTGTTG TCATTGTCCG CGCAACTATC GGTATCAAGC TGTTTAAGAA	1500
ATTCACCTCG AAAGCAAGCT GATAAACCGA TACAATTAAA GGCTCCTTTT GGAGCCTTTT	1560
TTTTTGAGAG TTTTGAACGT GAAAAAATTA TTATTGCAA TTCCTTTAGT TGTTCTTTTC	1620
TATTCTCACT CCGCTGAAAC TGTTGAAAGT TGTTTAGCAA AACCCCATAC AGAAAAATCA	1680
TTTACTAACG TCTGGAAAGA CGACAAAAC TTAGATCGTT ACGCTAACTA TGAGGGTTGT	1740
CTGTGGAATG CTACAGGCCG TGTAAGTTGT ACTGGTGACG AAACCTCAGT TTACGGTACA	1800
TGGGTTCCCTA TTGGGCTTGC TATCCCTGAA AATGAGGGTG GTGGCTCTGA GGGTGGCCGT	1860
TCTGAGGGTG GCGGTTCTGA GGGTGGCGGT ACTAAACCTC CTGAGTACGG TGATACACCT	1920
ATCCGGGGCT ATACTTATAT CAACCTCTC GACGGCACTT ATCCGCCTGG TACTGAGCAA	1980
AACCCCGCTA ATCCTAATCC TTCTCTTGAG GAGTCTCAGC CTCTTAATAC TTTCATGTTT	2040
CAGAATAATA GGTTCGAAA TAGGCAGGGG GCATTAACTG TTTATACGGG CACTGTTACT	2100

CAAGGCACTG	ACCCCGTTAA	AACTTATTAC	CAGTACACTC	CTGTATCATC	AAAAGCCATG	2160
TATGAGCCTT	ACTGGAACGG	TAAATTCAGA	GA CTGCGCTT	TCCATTCTGG	CTTTAATGAA	2220
GATCCATTCC	TTTGTGAATA	TCAAGGCCAA	TCGTCTGACC	TGCCTCAACC	TCCTGTCAAT	2280
GCTGGCGGCG	GCTCTGGTGG	TGGTTCTGGT	GGCGGCTCTG	AGGGTGGTGG	CTCTGAGGGT	2340
GGCGGTTCTG	AGGGTGGCGG	CTCTGAGGGA	GGCGGTTCCG	GTGGTGGCTC	TGGTTCCGGT	2400
GATTTTGATT	ATGAAAAGAT	GGCAAACGCT	AATAAGGGGG	CTATGACCGA	AAATGCCGAT	2460
GAAAACGCGC	TACAGTCTGA	CGCTAAAGGC	AAACTTGATT	CTGTGCGTAC	TGATTACGGT	2520
GCTGCTATCG	ATGGTTTCAT	TGGTGACGTT	TCCGGCCTTG	CTAATGGTAA	TGGTGCTACT	2580
GGTGATTTTG	CTGGCTCTAA	TTCCCAAATG	GCTCAAGTCG	GTGACGGTGA	TAATTCACCT	2640
TTAATGAATA	ATTTCCGTGA	ATATTTACCT	TCCCTCCCTC	AATCGGTTGA	ATGTCGCCCT	2700
TTTGTCTTTA	GCGCTGGTAA	ACCATATGAA	TTTTCTATTG	ATTGTGACAA	AATAAACTTA	2760
TTCCGTGGTG	TCTTTGCGTT	TCTTTTATAT	GTTGCCACCT	TTATGTATGT	ATTTTCTACG	2820
TTTGCTAACA	TACTGCGTAA	TAAGGAGTCT	TAATCATGCC	AGTTCTTTTG	GGTATTCCGT	2880
TATTATTGCG	TTTCCTCGGT	TTGCTTCTGG	TAAC TTTGTT	CGGCTATCTG	CTTACTTTTC	2940
TTAAAAAGGG	CTTCGGTAAG	ATAGCTATTG	CTATTTTCATT	GTTTCTTGCT	CTTATTATTG	3000
GGCTTAACTC	AATTCTTG TG	GGTTATCTCT	CTGATATTAG	CGCTCAATTA	CCCTCTGACT	3060
TTGTTCAAGG	TGTTCAAGTTA	ATTCTCCCGT	CTAATGCGCT	TCCCTGTTTT	TATGTTATTG	3120
TCTCTGTAAA	GGCTGCTATT	TTCA TTTTGG	ACGTTAAACA	AAAAATCGTT	TCTTATTTGG	3180
ATTGGGATAA	ATAATATGGC	TGTTTATTTT	GTAAC TGGCA	AATTAGGCTC	TGGAAAGACG	3240
CTCGTTAGCG	TTGGTAAGAT	TTAGGATAAA	ATTGTAGCTG	GCTGCAAAAT	AGCAACTAAT	3300
CTTGATT TAA	GGCTTCAAAA	CCTCCCGCAA	GTCCGGAGGT	TCGCTAAAAC	GCCTCGCGTT	3360
CTTAGAATAC	CGGATAAGCC	TTCTATATCT	GATTTGCTTG	CTATTGGGCG	CGGTAATGAT	3420
TCCTACGATG	AAAATAAAAA	CGGCTTGCTT	GTTCTCGATG	AGTGC GG TAC	TTGGTTTAA T	3480
ACCCGTTCTT	GGAATGATAA	GGAAAGACAG	CCGATTATTG	ATTGGTTTCT	ACATGCTCGT	3540
AAATTAGGAT	GGGATATTAT	TTTTCTTGTT	CAGGACTTAT	CTATTGTTGA	TAAACAGGCG	3600
CGTTCTGCAT	TAGCTGAACA	TGTTGTTTAT	TGTCGTCGTC	TGGACAGAAT	TACTTTACCT	3660
TTTGTGGGTA	CTTTATATTG	TCTTATTACT	GGCTCGAAAA	TGCCTCTGCC	TAAATTACAT	3720
GTGCGGTTG	TTAAATATGG	CGATTCTCAA	TAAAGCCCTA	CTGTTGAGCG	TTGGCTTTAT	3780
ACTGGTAAGA	ATTTGTATAA	CGCATATGAT	ACTAAACAGG	CTTTTTCTAG	TAATTATGAT	3840
TCCGGTGTTT	ATTCTTATTT	AACGCCTTAT	TTATCACACG	GTCCGTATTT	CAAACCATTA	3900
AATTTAGGTC	AGAAGATGAA	GCTTACTAAA	ATATATTTGA	AAAAGTTTTT	ACGCGTTCTT	3960
TGTCTTGCGA	TTGGATTTGC	ATCAGCATTT	ACATATAGTT	ATATAACCCA	ACCTAAGCCG	4020
GAGGT TAAAA	AGGTAGTCTC	TCAGACCTAT	GATTTTGATA	AATTC ACTAT	TGACTCTTCT	4080
CAGCGTCTTA	ATCTAAGCTA	TCGCTATGTT	TTCAAGGATT	CTAAGGGAAA	ATTAATTAAT	4140

AGCGACGATT	TACAGAAGCA	AGGTTATTCA	CTCACATATA	TTGATTTATG	TACTGTTTCC	4200
ATTAAAAAAG	GTAATTCAAA	TGAAATTGTT	AAATGTAATT	AATTTTGTTT	TCTTGATGTT	4260
TGTTTCATCA	TCTTCTTTTG	CTCAGGTAAT	TGAAATGAAT	AATTCGCCTC	TGCGCGATTT	4320
TGTAACCTGG	TATTCAAAGC	AATCAGGCCA	ATCCGTTATT	GTTTCTCCCG	ATGTAAAAGG	4380
TACTGTTACT	GTATATTCAT	CTGACGTAA	ACCTGAAAAT	CTACGCAATT	TCTTTATTTT	4440
TGTTTTACGT	GCTAATAATT	TTGATATGGT	TGGTTCAATT	CCTTCCATAA	TTCAGAAGTA	4500
TAATCCAAAC	AATCAGGATT	ATATTGATGA	ATTGCCATCA	TCTGATAATC	AGGAATATGA	4560
TGATAATTCC	GCTCCTTCTG	GTGGTTTCTT	TGTTCCGCAA	AATGATAATG	TACTCAAAC	4620
TTTTAAAATT	AATAACGTTT	GGGCAAAGGA	TTTAATACGA	GTTGTGCAAT	TGTTTGTAAG	4680
GTCTAATACT	TCTAAATCCT	CAAATGTATT	ATCTATTGAC	GGCTCTAATC	TATTAGTTGT	4740
TAGTGCACCT	AAAGATATTT	TAGATAACCT	TCCTCAATTC	CTTTCTACTG	TTGATTTGCC	4800
AACTGACCAG	ATATTGATTG	AGGGTTTGAT	ATTGAGGTT	CAGCAAGGTG	ATGCTTTAGA	4860
TTTTTCATTT	GCTGCTGGCT	CTCAGCGTGG	CACTGTTGCA	GGCGGTGTTA	ATACTGACCG	4920
CCTCACCTCT	GTTTTATCTT	CTGCTGGTGG	TTCGTTCCGT	ATTTTAAATG	GCGATGTTTT	4980
AGGGCTATCA	GTTCCGCGCAT	TAAAGACTAA	TAGCCATTCA	AAAATATTGT	CTGTGCCACG	5040
TATTCTTACG	CTTTCAGGTC	AGAAGGGTTC	TATCTCTGTT	GGCCAGAATG	TCCCTTTTAT	5100
TACTGGTCGT	GTGACTGGTG	AATCTGCCAA	TGTAAATAAT	CCATTTTCTA	CGATTGAGCG	5160
TCAAAATGTA	GGTATTTCCA	TGAGCGTTTT	TCCTGTTGCA	ATGGCTGGCG	GTAATATTGT	5220
TCTGGATATT	ACCAGCAAGG	CCGATAGTTT	GAGTTCTTCT	ACTCAGGCAA	GTGATGTTAT	5280
TACTAATCAA	AGAAGTATTG	CTACAACGGT	TAATTTGCGT	GATGGACAGA	CTCTTTTACT	5340
CGGTGGCCTC	ACTGATTATA	AAAACACTTC	TCAAGATTCT	GGCGTACCGT	TCCTGTCTAA	5400
AATCCCTTTA	ATCGGCCTCC	TGTTTAGCTC	CCGCTCTGAT	TCCAACGAGG	AAAGCACGTT	5460
ATACGTGCTC	GTCAAAGCAA	CCATAGTACG	CGCCCTGTAG	CGGCGCATT	AGCGCGGCGG	5520
GTGTGGTGGT	TACGCGCAGC	GTGACCGCTA	CACTTGCCAG	CGCCCTAGCG	CCCGCTCCTT	5580
TCGCTTTCTT	CCCTTCCTTT	CTCGCCACGT	TCGCGGCTT	TCCCCGTCAA	GCTCTAAATC	5640
GGGGGCTCCC	TTTAGGGTTC	CGATTTAGTG	CTTTACGGCA	CCTCGACCCC	AAAAAAGTTG	5700
ATTTGGGTGA	TGGTTCACGT	AGTGGGCCAT	CGCCCTGATA	GACGGTTTTT	CGCCCTTTGA	5760
CGTTGGAGTC	CACGTTCTTT	AATAGTGGAC	TCTTGTTCCA	AACTGGAACA	ACACTCAACC	5820
CTATCTCGGG	CTATTCTTTT	GATTTATAAG	GGATTTTGCC	GATTTTCGGAA	CCACCATCAA	5880
ACAGGATTTT	CGCCTGCTGG	GGCAAACCAG	CGTGGACCGC	TTGCTGCAAC	TCTCTCAGGG	5940
CCAGGCGGTG	AAGGGCAATC	AGCTGTTGCC	CGTCTCGCTG	GTGAAAAGAA	AAACCACCCT	6000
GGCGCCCAAT	ACGCAAACCG	CCTCTCCCGG	CGCGTTGGCC	GATTCATTAA	TGCAGCTGGC	6060
ACGACAGGTT	TCCCGACTGG	AAAGCGGGCA	GTGAGCGCAA	CGCAATTAAT	GTGAGTTAGC	6120
TCACTCATT	GGCACCCAG	GCTTTACACT	TTATGCTTCC	GGCTCGTATG	TTGTGTGGAA	6180

92

TTGTGAGCGG	ATAACAATTT	CACACGCGTC	ACTTGGCACT	GGCCGTCGTT	TTACAACGTC	6240
GTGACTGGGA	AAACCCTGGC	GTTACCCAAG	CTTTGTACAT	GGAGAAAATA	AAGTGAAACA	6300
AAGCACTATT	GCACTGGCAC	TCTTACCGTT	ACTGTTTACC	CCTGTGGCAA	AAGCCCTTCT	6360
GAGGCATCCG	GGAGCTGAAG	GCGATGACCC	TGCTAAGGCT	GCATTCAATA	GTTTACAGGC	6420
AAGTGCTACT	GAGTACATTG	GCTACGCTTG	GGCTATGGTA	GTAGTTATAG	TTGGTGCTAC	6480
CATAGGGATT	AAATTATTCA	AAAAGTTTAC	GAGCAAGGCT	TCTTAAGCAA	TAGCGAAGAG	6540
GCCCGCACCG	ATCGCCCTTC	CCAACAGTTG	CGCAGCCTGA	ATGGCGAATG	GCGCTTTGCC	6600
TGGTTTCCGG	CACCAGAAGC	GGTGCCGGAA	AGCTGGCTGG	AGTGCGATCT	TCCTGAGGCC	6660
GATACGGTGG	TCGTCCCCTC	AAACTGGCAG	ATGCACGGTT	ACGATGCGCC	CATCTACACC	6720
AACGTAACCT	ATCCGATTAC	GGTCAATCCG	CCGTTTGTTT	CCACGGAGAA	TCCGACGGGT	6780
TGTTACTCGC	TCACATTTAA	TGTTGATGAA	AGCTGGCTAC	AGGAAGGCCA	GACGCGAATT	6840
ATTTTGTATG	GCGTTCCTAT	TGGTTAAAAA	ATGAGCTGAT	TTAACAAAAA	TTTAACGCGA	6900
ATTTTAACAA	AATATTAACG	TTTACAATTT	AAATATTTCG	TTATACAATC	TTCCTGTTTT	6960
TGGGGCTTTT	CTGATTATCA	ACCGGGGTAC	ATATGATTGA	CATGCTAGTT	TTACGATTAC	7020
CGTTCATCGA	TTCTCTTGTT	TGCTCCAGAC	TCTCAGGCAA	TGACCTGATA	GCCTTTGTAG	7080
ATCTCTCAAA	AATAGCTACC	CTCTCCGGCA	TTAATTTATC	AGCTAGAACG	GTTGAATATC	7140
ATATTGATGG	TGATTGACT	GTCTCCGGCC	TTTCTCACCC	TTTTGAATCT	TTACCTACAC	7200
ATTACTCAGG	CATTGCATTT	AAAATATATG	AGGGTCTTAA	AAATTTTTAT	CCTTGCGTTG	7260
AAATAAAGGC	TTCTCCCGCA	AAAGTATTAC	AGGGTGATAA	TGTTTTTGGT	ACAACCGATT	7320
TAGCTTTATG	CTCTGAGGCT	TTATTGCTTA	ATTTTGCTAA	TTCTTTGCCT	TGCCTGTATG	7380
ATTTATTGGA	CGTT					7394

(2) INFORMATION FOR SEQ ID NO:7:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 37 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

GATCCTAGGC TGAAGGCGAT GACCCTGCTA AGGCTGC

37

(2) INFORMATION FOR SEQ ID NO:8:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 35 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

ATTCAATAGT TTACAGGCAA GTGCTACTGA GTACA

35

(2) INFORMATION FOR SEQ ID NO:9:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 35 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

TTGGCTACGC TTGGGCTATG GTAGTAGTTA TAGTT

35

(2) INFORMATION FOR SEQ ID NO:10:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 35 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

GGTGCTACCA TAGGGATTAA ATTATTCAAA AAGTT

35

(2) INFORMATION FOR SEQ ID NO:11:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 18 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

TAGGAGCAAG GCTTCTTA

18

(2) INFORMATION FOR SEQ ID NO:12:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 39 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

AGCTTAAGAA GCCTTGCTCG TAAACTTTTT GAATAATTT

39

(2) INFORMATION FOR SEQ ID NO:13:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 36 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

AATCCCTATG GTAGCACCAA CTATAACTAC TACCAT

36

(2) INFORMATION FOR SEQ ID NO:14:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 35 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

AGCCCAAGCG TAGCCAATGT ACTCAGTAGC ACTTG

35

(2) INFORMATION FOR SEQ ID NO:15:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 34 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:

CCTGTAAACT ATTGAATGCA GCCTTAGCAG GGTC

34

(2) INFORMATION FOR SEQ ID NO:16:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 16 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:16:

ATCGCCTTCA GCCTAG

16

(2) INFORMATION FOR SEQ ID NO:17:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 27 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:17:
CTCGAATTCTG TACATCCTGG TCATAGC 27
- (2) INFORMATION FOR SEQ ID NO:18:
- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 21 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:18:
CATTTTGTCA GATGGCTTAG A 21
- (2) INFORMATION FOR SEQ ID NO:19:
- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 18 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:19:
TAGCATTAAC GTCCAATA 18
- (2) INFORMATION FOR SEQ ID NO:20:
- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 26 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:20:
ATATATTTTA GTAAGCTTCA TCTTCT 26
- (2) INFORMATION FOR SEQ ID NO:21:
- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 23 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:21:
GACAAAGAAC GCGTGAAAAC TTT 23

(2) INFORMATION FOR SEQ ID NO:22:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 35 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:22:

GCGGGCCTCT TCGCTATTGC TTAAGAAGCC TTGCT

35

(2) INFORMATION FOR SEQ ID NO:23:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 48 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:23:

TTCAGCCTAG GATCCGCCGA GCTCTCCTAC CTGCGAATTC GTACATCC

48

(2) INFORMATION FOR SEQ ID NO:24:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 24 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:24:

TGGATTATAC TTCTAAATAA TGGA

24

(2) INFORMATION FOR SEQ ID NO:25:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 36 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:25:

TAACACTCAT TCCGGATGGA ATTCTGGAGT CTGGGT

36

(2) INFORMATION FOR SEQ ID NO:26:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 22 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:26:

AATTCGCCAA GGAGACAGTC AT

22

(2) INFORMATION FOR SEQ ID NO:27:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 39 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:27:

AATGAAATAC CTATTGCCTA CGGCAGCCGC TGGATTGTT

39

(2) INFORMATION FOR SEQ ID NO:28:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 39 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:28:

ATTACTCGCT GCCCAACCAG CCATGGCCGA GCTCGTGAT

39

(2) INFORMATION FOR SEQ ID NO:29:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 39 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:29:

GACCCAGACT CCAGATATCC AACAGGAATG AGTGTTAAT

39

(2) INFORMATION FOR SEQ ID NO:30:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 13 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:30:

TCTAGAACGC GTC

13

(2) INFORMATION FOR SEQ ID NO:31:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 35 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:31:

ACGTGACGCG TTCTAGAATT AACACTCATT CCTGT

35

(2) INFORMATION FOR SEQ ID NO:32:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 39 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:32:

TGGATATCTG GAGTCTGGGT CATCAGGAGC TCGGCCATG

39

(2) INFORMATION FOR SEQ ID NO:33:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 39 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:33:

GCTGGTTGGG CAGCGAGTAA TAACAATCCA GCGGCTGCC

39

(2) INFORMATION FOR SEQ ID NO:34:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 37 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:34:

GTAGGCAATA GGTATTTTCAT TATGACTGTC CTTGGCG

37

(2) INFORMATION FOR SEQ ID NO:35:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 30 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:35:
TGACTGTCTC CTTGGCGTGT GAAATTGTIA 30
- (2) INFORMATION FOR SEQ ID NO:36:
- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 36 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:36:
TAACACTCAT TCCGGATGGA ATTCTGGAGT CTGGGT 36
- (2) INFORMATION FOR SEQ ID NO:37:
- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 25 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:37:
CAATTTTATC CTAAATCTTA CCAAC 25
- (2) INFORMATION FOR SEQ ID NO:38:
- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 21 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:38:
CATTTTTGCA GATGGCTTAG A 21
- (2) INFORMATION FOR SEQ ID NO:39:
- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 21 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:39:
CGAAAGGGGG GTGTGCTGCA A 21

100

(2) INFORMATION FOR SEQ ID NO:40:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 18 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:40:

TAGCATTAAC GTCCAATA

18

(2) INFORMATION FOR SEQ ID NO:41:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 43 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:41:

AAACGACGGC CAGTGCCAAG TGACGCGTGT GAAATTGTTA TCC

43

(2) INFORMATION FOR SEQ ID NO:42:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 43 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:42:

GGCGAAAGGG AATTCTGCAA GGCGATTAAG CTTGGGTAAC GCC

43

(2) INFORMATION FOR SEQ ID NO:43:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 36 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:43:

GGCGTTACCC AAGCTTTGTA CATGGAGAAA ATAAAG

36

101

(2) INFORMATION FOR SEQ ID NO:44:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 42 bas pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:44:

TGAAACAAAG CACTATTGCA CTGGCACTCT TACCGTTACC GT

42

(2) INFORMATION FOR SEQ ID NO:45:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 42 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:45:

TACTGTTTAC CCCTGTGACA AAAGCCGCCC AGGTCCAGCT GC

42

(2) INFORMATION FOR SEQ ID NO:46:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 44 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:46:

TCGAGTCAGG CCTATTGTGC CCAGGGATTG TACTAGTGGA TCCG

44

(2) INFORMATION FOR SEQ ID NO:47:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 38 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:47:

TGGCGAAAGG GAATTCGGAT CCACTAGTAC AATCCCTG

38

(2) INFORMATION FOR SEQ ID NO:48:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 42 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:48:

GGCACAATAG GCCTGACTCG AGCAGCTGGA CCAGGGCGGC TT

42

(2) INFORMATION FOR SEQ ID NO:49:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 42 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:49:

TTGTCACAGG GGTAACAGT AACGGTAACG GTAAGTGTGC CA

42

(2) INFORMATION FOR SEQ ID NO:50:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 42 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:50:

GTGCAATAGT GCTTTGTTTC ACTTTATTTT CTCCATGTAC AA

42

(2) INFORMATION FOR SEQ ID NO:51:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 21 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:51:

TAACGGTAAG AGTGCCAGTG C

21

(52) INFORMATION FOR SEQ ID NO:52:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 68 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ix) FEATURE:

- (A) NAME/KEY: misc difference
- (B) LOCATION: r place(25, "")
- (D) OTHER INFORMATION: /note= "M REPRESENTS AN EQUAL MIXTURE OF A AND C AT THIS LOCATION AND AT LOCATIONS 28, 31, 34, 37, 40, 43, 46 & 49"

103

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:52:

AGCTCCCGGA TGCCTCAGAA GATGMNNMNN MNNMNNMNNM NNMNNMNNMN NGGCTTTTGC 60
CACAGGGG 68

(2) INFORMATION FOR SEQ ID NO:53:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 54 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ix) FEATURE:

- (A) NAME/KEY: misc difference
- (B) LOCATION: replace(17, "")
- (D) OTHER INFORMATION: /note= "M REPRESENTS AN EQUAL MIXTURE OF A AND C AT THIS LOCATION AND AT LOCATIONS 20, 23, 26, 29, 32, 35, 38, 41, 44 & 50"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:53:

CAGCCTCGGA TCCGCCMNNM NNMNNMNNMN NMNNMNNMNN MNNMNNATGM GAAT 54

(2) INFORMATION FOR SEQ ID NO:54:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 27 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:54:

GGTAAACACT AACGGTAAGA GTGCCAG 27

(2) INFORMATION FOR SEQ ID NO:55:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 19 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:55:

GGGCTTTTGC CACAGGGGT 19

(2) INFORMATION FOR SEQ ID NO:56:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 63 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

104

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:56:

AGGGTCATCG CCTTCAGCTC CGGATCCCTC AGAAGTCATA AACCCCCCAT AGGCTTTTGC 60
CAC 63

(2) INFORMATION FOR SEQ ID NO:57:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 47 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:57:

TGGCCTTCAG CTCCCGGATG CCTCAGAAGC ATGAACCCCC CATAGGC 47

(2) INFORMATION FOR SEQ ID NO:58:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 25 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:58:

CAATTTTATC CTAAATCTTA CCAAC 25

(2) INFORMATION FOR SEQ ID NO:59:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 21 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:59:

GCCTTCAGCC TCGGATCCGC C 21

(2) INFORMATION FOR SEQ ID NO:60:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 21 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:60:

CGGATGCCTC AGAAGCCCCN N 21

105

(2) INFORMATION FOR SEQ ID NO:61:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 30 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:61:

CGGATGCCTC AGAAGGGCTT TTGCCACAGG

30

I CLAIM:

1. A composition of matter comprising a plurality of cells containing a diverse population of expressible oligonucleotides operationally linked to expression elements, said expressible oligonucleotides
5 having a desirable bias of random codon sequences produced from random combinations of first and second oligonucleotide precursor populations having a desirable bias of random codon sequences.
2. The composition of claim 1, wherein the desirable bias of random codon sequences of said first and second oligonucleotides is unbiased.
3. The composition of claim 1, wherein the desirable bias of random codon sequences of said first and second oligonucleotides is biased toward a predetermined sequence.
4. The composition of claim 1, wherein said first and second oligonucleotides having random codon sequences have at least one specified codon at a predetermined position.
5. The composition of claim 1, wherein said cells are procaryotes.
6. The composition of claim 1, wherein said cells are E. coli.

7. A kit for the preparation of vectors useful for the expression of a diverse population of random peptides from combined first and second oligonucleotides having a desirable bias of random codon sequences,
5 comprising: two vectors: a first vector having a cloning site for said first oligonucleotides and a pair of restriction sites for operationally combining first oligonucleotides with second oligonucleotides; and a
10 second vector having a cloning site for said second oligonucleotides and a pair of restriction sites complementary to those on said first vector, one or both vectors containing expression elements capable of being operationally linked to said combined first and second oligonucleotides.

8. The kit of claim 7, wherein said vectors are in a filamentous bacteriophage.

9. The kit of claim 8, wherein said filamentous bacteriophage are M13.

10. The kit of claim 7, wherein said vectors are plasmids.

11. The kit of claim 7, wherein said vectors are phagemids.

12. The kit of claim 7, wherein the desirable bias of random codon sequences of said first and second oligonucleotides is unbiased.

13. The kit of claim 7, wherein the desirable bias of random codon sequences of said first and second oligonucleotides is diverse but biased toward a predetermined sequenc .

14. The kit of claim 7, wherein said first and second oligonucleotides having a desirable bias of random codon sequences have at least one specified codon at a predetermined position.

15. The kit of claim 7, wherein said pair of restriction sites are Fok I.

16. A cloning system for expressing random peptides from diverse populations of combined first and second oligonucleotides having a desirable bias of random codon sequences, comprising: a set of first vectors
5 having a diverse population of first oligonucleotides having a desirable bias of random codon sequences and a set of second vectors having a diverse population of second oligonucleotides having a desirable bias of random codon sequences, said first and second vectors each
10 having a pair of restriction sites so as to allow the operational combination of first and second oligonucleotides into a contiguous oligonucleotide having a desirable bias of random codon sequences.

17. The cloning system of claim 16, wherein the desirable bias of random codon sequences of said first and second oligonucleotides is unbiased.

18. The cloning system of claim 16, wherein the desirable bias of random codon sequences of said first and second oligonucleotides is diverse but biased toward a predetermined sequence.

19. The cloning system of claim 16, wherein said first and second oligonucleotides having a desirable bias of random codon sequences have at least one specified codon at a pr determined position.

20. The cloning system of claim 16, wherein said combined first and second vectors is through a pair of restriction sites.

21. The cloning system of claim 16, wherein said pair of restriction sites are Fok I.

22. A composition of matter comprising a plurality of cells containing a diverse population of expressible oligonucleotides operationally linked to expression elements, said expressible oligonucleotides
5 having a desirable bias of random codon sequences.

23. The composition of claim 22, wherein said cells are procaryotes.

24. The composition of claim 22, wherein said expressible oligonucleotides are expressed as peptide fusion proteins on the surface of a filamentous bacteriophage.

25. The composition of claim 22, wherein said filamentous bacteriophage is M13.

26. The composition of claim 22, wherein said fusion protein contains the product of gene VIII.

27. The composition of claim 22, wherein said diverse population of oligonucleotides having a desirable bias of random codon sequences are produced from the combination of diverse populations of first and second
5 oligonucleotides having a desirable bias of random codon sequences.

28. The composition of claim 22, wherein the desirable bias of random codon sequences of said oligonucleotides is unbiased.

29. The composition of claim 22, wherein the desirable bias of random codon sequences of said oligonucleotides is diverse but biased toward a predetermined sequence.

30. The composition of claim 22, wherein said oligonucleotides having a desirable bias of random codon sequences have at least one specified codon at a predetermined position.

31. A plurality of vectors containing a diverse population of expressible oligonucleotides having a desirable bias of random codon sequences.

32. The vectors of claim 31, wherein said oligonucleotides are expressible as fusion proteins on the surface of filamentous bacteriophage.

33. The vectors of claim 31, wherein said filamentous bacteriophage is M13.

34. The vectors of claim 31, wherein said fusion protein contains the product of gene VIII.

35. The vectors of claim 31, wherein the desirable bias of random codon sequences of said oligonucleotides is unbiased.

36. The vectors of claim 31, wherein the desirable bias of random codon sequences of said oligonucleotides is diverse but biased toward a predetermined sequence.

37. The vectors of claim 31, wherein said oligonucleotides having a desirable bias of random codon sequences have at least one specified codon at a predetermined position.

38. A composition of matter, comprising a diverse population of oligonucleotides having a desirable bias of random codon sequences produced from random combinations of two or more oligonucleotide precursor
5 populations having a desirable bias of random codon sequences.

39. A method of constructing a diverse population of vectors having combined first and second oligonucleotides having a desirable bias of random codon sequences capable of expressing said combined
5 oligonucleotides as random peptides, comprising the steps of:

- 10 (a) operationally linking sequences from a diverse population of first oligonucleotides having a desirable bias of random codon sequences to a first vector;
- 15 (b) operationally linking sequences from a diverse population of second oligonucleotides having a desirable bias of random codon sequences to a second vector; and
- 20 (c) combining the vector products of steps (a) and (b) under conditions where said populations of first and second oligonucleotides are joined together into a population of combined vectors capable of being expressed.

40. The method of claim 39, wherein the desirable bias of random codon sequences of said first and second oligonucleotides is unbiased.

41. The method of claim 39, wherein the desirable bias of random codon sequences of said first and second oligonucleotides is diverse but biased toward a predetermined sequence.

42. The method of claim 39, wherein said first and second oligonucleotides having a desirable bias of random codon sequences have at least one specified codon at a predetermined position.

43. The method of claim 38, wherein steps (a) through (c) are repeated two or more times.

44. A method of selecting a peptide capable of being bound by a ligand binding protein from a population of random peptides, comprising:

- 5 (a) operationally linking a diverse population of first oligonucleotides having a desirable bias of random codon sequences to a first vector;
- 10 (b) operationally linking a diverse population of second oligonucleotides having a desirable bias of random codon sequences to a second vector;
- 15 (c) combining the vector products of steps (a) and (b) under conditions where said populations of first and second oligonucleotides are joined together into a population of combined vectors;
- 20 (d) introducing said population of combined vectors into a compatible host under conditions sufficient for expressing said population of random peptides; and
- (e) determining the peptide which binds to said ligand binding protein.

45. The method of claim 44, wherein the desirable bias of random codon sequences of said first and second oligonucleotides is unbiased.

46. The method of claim 44, wherein the desirable bias of random codon sequences of said first and second oligonucleotides is diverse but biased toward a predetermined sequence.

47. The method of claim 44, wherein said first and second oligonucleotides having a desirable bias of random codon sequences have at least one specified codon at a predetermined position.

48. The method of claim 44, wherein steps (a) through (c) are repeated two or more times.

49. A method for determining the nucleic acid sequence encoding a peptide capable of being bound by a ligand binding protein which is selected from a population of random peptides, comprising:

- 5 (a) operationally linking a diverse population
of first oligonucleotides having a
desirable bias of random codon sequences
to a first vector;
- 10 (b) operationally linking a diverse population
of second oligonucleotides having a
desirable bias of random codon sequences
to a second vector;
- 15 (c) combining the vector products of steps (a)
and (b) under conditions where said
populations of first and second
oligonucleotides are joined together into
a population of combined vectors;
- 20 (d) introducing said population of combined
vectors into a compatible host under
conditions sufficient for expressing said
population of random peptides;
- 25 (e) determining the peptide which binds to
said ligand binding protein;
- (f) isolating the nucleic acid encoding said
peptide; and
- (g) sequencing said nucleic acid.

50. The method of claim 49, wherein the desirable bias of random codon sequences of said first and second oligonucleotides is unbiased.

51. The method of claim 49, wherein the desirable bias of random codon sequences of said first and second oligonucleotides is diverse but biased toward a predetermined sequence.

52. The method of claim 49, wherein said first and second oligonucleotides having a desirable bias of random codon sequences have at least one specified codon at a predetermined position.

53. The method of claim 49, wherein steps (a) through (c) are repeated two or more times.

54. A method of constructing a diverse population of vectors containing expressible oligonucleotides having a desirable bias of random codon sequences, comprising operationally linking a diverse
5 population of oligonucleotides having a desirable bias of random codon sequences to expression elements.

55. The method of claim 54, wherein said oligonucleotides are expressible as fusion proteins on the surface of filamentous bacteriophage.

56. The method of claim 54, wherein said filamentous bacteriophage are M13.

57. The method of claim 54, wherein said fusion protein contains the product of gene VIII.

58. The method of claim 54, wherein the desirable bias of random codon sequences of said oligonucleotides is unbiased.

59. The method of claim 54, wherein the desirable bias of random codon sequences of said oligonucleotides is diverse but biased toward a predetermined sequence.

60. The method of claim 54, wherein said oligonucleotides having a desirable bias of random codon sequences have at least one specified codon at a predetermined position.

61. The method of claim 54, wherein said operationally linking further comprising the steps of:

5 (a) operationally linking a diverse population of first oligonucleotides having a desirable bias of random codon sequences to a first vector;

10 (b) operationally linking a diverse population of second oligonucleotides having a desirable bias of random codon sequences to a second vector; and

15 (c) combining the vector products of steps (a) and (b) under conditions where said populations of first and second oligonucleotides are joined together into a population of combined vectors.

62. The method of claim 61, wherein steps (a) through (c) are repeated two or more times.

63. A method of selecting a peptid capable of being bound by a binding protein from a population of random peptides, comprising:

- 5 (a) operationally linking a diverse population of oligonucleotides having a desirable bias of random codon sequences to expression elements;
- 10 (b) introducing said population of vectors into a compatible host under conditions sufficient for expressing said population of random peptides; and
- (c) determining the peptide which binds to said ligand binding protein.

64. The method of claim 63, wherein said population of random peptides are expressed as fusion proteins on the surface of filamentous bacteriophage.

65. The method of claim 63, wherein said filamentous bacteriophage are M13.

66. The method of claim 63, wherein said fusion protein contains the product of gene VIII.

67. The method of claim 63, wherein the desirable bias of random codon sequences of said oligonucleotides is unbiased.

68. The method of claim 63, wherein the desirable bias of random codon sequences of said oligonucleotides is diverse but biased toward a predetermined sequence.

69. The method of claim 63, wherein said oligonucleotides having a desirable bias of random codon sequences have at least one specified codon at a predetermined position.

70. The method of claim 63, wherein step (a) further comprises:

- 5 (a1) operationally linking a diverse population of first oligonucleotides having a desirable bias of random codon sequences to a first vector;
- 10 (a2) operationally linking a diverse population of second oligonucleotides having a desirable bias of random codon sequences to a second vector; and
- 15 (a3) combining the vector products of steps (a) and (b) under conditions where said populations of first and second oligonucleotides are joined together into a population of combined vectors.

71. The method of claim 70, wherein steps (a1) through (a3) are repeated two or more times.

72. A method of determining the nucleic acid sequence encoding a peptide capable of being bound by a ligand binding protein which is selected from a population of random peptides, comprising:

- 5 (a) operationally linking a diverse population of oligonucleotides having a desirable bias of random codon sequences to expression elements.
- 10 (b) introducing said population of vectors into a compatible host under conditions sufficient for expressing said population of random peptides;
- (c) determining the peptide which binds to said ligand binding protein;
- 15 (d) isolating the nucleic acid encoding said peptide; and
- (e) sequencing said nucleic acid.

73. The method of claim 72, wherein said population of random peptides are expressed as fusion proteins on the surface of filamentous bacteriophage.

74. The method of claim 72, wherein said filamentous bacteriophage are M13.

75. The method of claim 72, wherein said fusion protein contains the product of gene VIII.

76. The method of claim 72, wherein the desirable bias of random codon sequences of said oligonucleotides is unbiased.

77. The method of claim 72, wherein the desirable bias of random codon sequences of said oligonucleotides is diverse but biased toward a predetermined sequence.

78. The method of claim 72, wherein said oligonucleotides having a desirable bias of random codon sequences have at least one specified codon at a predetermined position.

79. The method of claim 72, wherein step (a) further comprises:

5 (a1) operationally linking a diverse population of first oligonucleotides having a desirable bias of random codon sequences to a first vector;

10 (a2) operationally linking a diverse population of second oligonucleotides having a desirable bias of random codon sequences to a second vector; and

15 (a3) combining the vector products of steps (a) and (b) under conditions where said populations of first and second oligonucleotides are joined together into a population of combined vectors.

80. The method of claim 78, wherein steps (a1) through (a3) are repeated two or more times.

81. A vector comprising two copies of a gene encoding a filamentous bacteriophage coat protein, both copies encoding substantially the same amino acid sequence but having different nucleotide sequences.

82. The vector of claim 81, wherein said filamentous bacteriophage is M13.

83. The vector of claim 81, wherein said gene is gene VIII.

84. The vector of claim 81, wherein said vector has substantially the sequence shown in Figure 5 (SEQ ID NO: 1).

85. A vector comprising two copies of a gene encoding a filamentous bacteriophage coat protein, one copy of said gene capable of being operationally linked to an oligonucleotide wherein said oligonucleotide can be
5 expressed as a fusion protein on the surface of said filamentous bacteriophage or as a soluble peptide.

86. The vector of claim 84, wherein said one copy of said gene is expressed on the surface of said filamentous bacteriophage.

87. The vector of claim 84, wherein said bacteriophage coat protein is M13 gene VIII.

1 / 16




FIG. 1

SUBSTITUTE SHEET

2 / 16






FIG. 2

3 / 16



**FIG. 4**

5 / 16

	10	20	30	40	50	60
1	AATGCTACTA	CTATTAGTAG	AATTGATGCC	ACCTTTTCAG	CTCGCGCCCC	AAATGAAAAAT
61	ATAGCTAAAC	AGGTTATTGA	CCATTTGCGA	AATGTATCTA	ATGGTCAAAC	TAAATCTACT
121	CGTTCGCAGA	ATTGGGAATC	AACTGTTACA	TGGAATGAAA	CTTCCAGACA	CCGTACTTTA
181	GTTGCATATT	TAAAAACATG	TGAGCTACAG	CACCAGATTG	AGCAATTAAG	CTCTAAGCCA
241	TCTGCAAAAA	TGACCTCTTA	TCAAAAAGGAG	CAATTAAGG	IACCTCTTAA	TCCTGACCTG
301	TTGGAGTTTG	CTTCCGGTCT	GGTTCGCTTT	GAAGCTCGAA	TTAAAACGCG	ATATTTGAAG
361	TCTTTCGGGC	TTCTCTTAA	TCTTTTGTAT	GCAATCCGCT	TTGCTTCTGA	CTATAATAGT
421	CAGGGTAAAG	ACCTGATTTT	TGATTTATGG	TCATTCCTGT	TTTCTGAAGT	GTTTAAAGCA
481	TTTGAGGGGG	ATTCAATGAA	TATTTATGAC	GATTCGCGAG	TATTGGACGC	TATCCAGTCT
541	AAACATTTTA	CTATTACCCC	CTCTGGCAAA	ACTTCTTTTG	CAAAAGCCTC	TCGCTATTTT
601	GGTTTTTATC	GTCGTCTGGT	AAACGAGGGT	TATGATAGTG	TTGCTCTTAC	TATGCCTCGT
661	AATTCCTTTT	GGCGTTATGT	ATCTGCATTA	GTGAAATGTG	GTATTCCTAA	ATCTCAACTG
721	ATGAATCTTT	CTACCTGTAA	TAATGTTGTT	CCGTTAGTTC	GTTTTATTAA	CGTAGATTTT
781	TCTTCCCAAC	GTCCTGACTG	GTATAATGAG	CCAGTTCCTA	AAATCGCATC	AGGTAATTCA
841	CAATGATTAA	AGTTGAAATT	AAACCATCTC	AAGCCCAATT	TACTACTCGT	TCTGGTGTTC
901	CTCGTCAGGG	CAAGCCTTAT	TCACTGAATG	AGCAGCTTGT	TTACGTTGAT	TTGGGTAATG
961	AATATCCGGT	TCTTGTCAAG	ATTACTCTTG	ATGAAGGTCA	GCCAGCCTAT	GCGCCTGGTC
1021	TGTACACCGT	TCATCTGTCC	TCTTTCAAAG	TTGGTCAGTT	CGGTTCCCTT	ATGATTGACC
1081	GTCTGCGCCT	CGTTCGGGCT	AAGTAACATG	GAGCAGGTCG	CGGATTTCTA	CACAATTTAT
1141	CAGGCGATGA	TACAAATCTC	CGTTGTACTT	TGTTTTGCGC	TTGGTATAAT	CGCTGGGGGT
1201	CAAAGATGAG	TGTTTTAGTG	TATTTCTTCG	CCTCTTTTCG	TTTAGGTTGG	TGCTTTCGTA
1261	GTGGCATTAC	GTATTTTACC	CGTTTAATGG	AAACTTCCTC	ATGAAAAAGT	CTTTAGTCTT
1321	CAAAGCCTCT	GTAGCCGTTG	CTACCCTCGT	TCCGATGCTG	TCTTTCGCTG	CTGAGGGTGA
1381	CGATCCCGCA	AAAGCGGCGT	TTAACTCCCT	GCAAGCCTCA	GCGACCGAAT	ATATCGGTTA
1441	TGCGTGGGCG	ATGGTTGTTG	TCATTGTGCG	CGCAACTATC	GGTATCAAGC	TGTTTAAGAA
1501	ATTCACCTCG	AAAGCAAGCT	GATAAAACCG	TACAATTAAT	GGCTCCTTTT	GGAGCCTTTT
1561	TTTTTGAGAG	TTTTCAACGT	GAAAAAATTA	TTATTCGCAA	TTCTTTTAGT	TGTTCTTTTC
1621	TATTTCTACT	CCGCTGAAAC	TGTTGAAAGT	TGTTTAGCAA	AACCCCATAC	AGAAAAATCA
1681	TTTACTAACG	TCTGGAAAGA	CGACAAAAC	TTAGATCGTT	ACGCTAACTA	TGAGGGTTGT
1741	CTGTGGAATG	CTACAGGCGT	TGTAGTTTGT	ACTGGTGACG	AAACTCAGTG	TTACGGTACA
1801	TGGGTTCCTA	TTGGGCTTGC	TATCCCTGAA	AATGAGGGTG	GTGGCTCTGA	GGGTGGCGGT
1861	TCTGAGGGTG	GCGGTTCTGA	GGGTGGCGGT	ACTAAACCTC	CTGAGTACGG	TGATACACCT
1921	ATTCCGGGCT	ATACTTATAT	CAACCCTCTC	GACGGCACTT	ATCCGCCTGG	TACTGAGCAA
1981	AACCCCGCTA	ATCCTAATCC	TTCTCTTGAG	GAGTCTCAGC	CTCTTAATAC	TTTATGTTT
2041	CAGAATAATA	GGTTCCGAAA	TAGGCAGGGG	GCATTAACCTG	TTTATACGGG	CACTGTTACT
2101	CAAGGCACTG	ACCCCGTTAA	AACCTTATTAC	CAGTACACTC	CTGTATCATC	AAAAGCCATG
2161	TATGACGCTT	ACTGGAACGG	TAAATTCAGA	GACTGCGCTT	TCCATTCTGG	CTTTAATGAA
2221	GATCCACTCG	TTTGTGAATA	TCAAGGCCAA	TCGTCTGACC	TCGGTCAACC	TCCTGTCAAT
2281	GCTGGCGGCG	GCTCTGGTGG	TGGTTCTGGT	GCGCGGCTCTG	AGGGTGGTGG	CTCTGAGGGT
2341	GGCGGTTCTG	AGGGTGGCGG	CTCTGAGGGA	GGCGGTTCCG	GTGGTGGCTC	TGGTTCGGGT
2401	GATTTTGATT	ATGAAAAGAT	GGCAAACGCT	AATAAGGGGG	CTATGACCGA	AAATGCCGAT
2461	GAAAACGCGC	TACAGTCTGA	CGCTAAAGGC	AAACTTGATT	CTGTCGCTAC	TGATTACGGT
2521	GCTGCTATCG	ATGGTTTCAT	TGGTGACGTT	TCCGGCCTTG	CTAATGGTAA	TGGTGCTACT
2581	GGTGATTTTG	CTGGCTCTAA	TTCCCAATAG	GCTCAAGTCG	GTGACGGTGA	TAATTCACCT
2641	TTAATGAATA	ATTTCCGTCA	ATATTTACCT	TCCCTCCCTC	AATCGGTTGA	ATGTCGCCCT
2701	TTTGTCTTTA	GCGCTGGTAA	ACCATATGAA	TTTTCTATTG	ATTGTGACAA	AATAAACCTT
2761	TTCCGTGGTG	TCTTTGCGTT	TCTTTTATAT	GTTGCCACCT	TTATGTATGT	ATTTTCTACG
2821	TTTGCTAACA	TACTGCGTAA	TAAGGAGTCT	TAATCATGCC	AGTTCTTTTG	GGTATTCCGT
2881	TATTATTGCG	TTTCTCTGGT	TTCTTCTGG	TAATTTTGT	GCCGTATCTG	CTTACTTTTC
2941	TTAAAAAGGG	CTTCGGTAAG	ATAGCTATTG	CTATTTTCAT	GTTTCTTGCT	CTTATTATTG
3001	GGCTTAACTC	AATTCTTGTG	GGTTATCTCT	CTGATATTAG	CGCTCAATTA	CCCTCTGACT
3061	TTGTTTCAGGG	TGTTTCAGTTA	ATTCTCCCGT	CTAATGCGCT	TCCCTGTTTT	TATGTTATTCT
3121	TCTCTGTAAA	GGCTGCTATT	TTTATTTTGT	ACGTAAACAA	AAAAATCGTT	TCTTATTTGG
3181	ATTGGGATAA	ATAATATGGC	TGTTTATTTT	GTAAGTGGCA	AATTAGGCTC	TGGAAAGACG
3241	CTCGTTAGCG	TTGGTAAGAT	TCAGGATAAA	ATTGTAGCTG	GGTGCAAAAT	AGCAACTAAT
3301	CCTGATTTAA	GGCTTCAAAA	CCTCCCGCAA	GTCGGGAGGT	TCGCTAAAC	GCCTCGCGTT
3361	CTTAGAATAC	CGGATAAGCC	TTCTATATCT	GATTTGCTTG	CTATTGGGCG	CGGTAATGAT
3421	TCCTACGATG	AAAATAAAAA	CGGCTTGCTT	GTTCTCGATG	AGTGGGTTAT	TTGGTTTAAAT
3481	ACCCGTTCTT	GGAATGATAA	GGAAGACAG	CCGATTATTG	ATTGGTTTCT	ACATGCTCGT
3541	AAATTAGGAT	GGGATATTAT	CTTCTTGTG	CAGGACTTAT	CTATTGTTGA	TAAACAGGCG
3601	CGTTCTGCAT	TAGCTGAACA	TCTTGTGTTT	TGTCGTGCTC	TGGACAGAAT	TACTTTACCT
3661	TTTGTGCGTA	CTTTATATTC	TCTTATTACT	GGCTCGAAAA	TGCCTCTGCC	TAAATTACAT
3721	GTTGGCGTTG	TTAAATATGG	CGATTCTCAA	TTAAGCCCTA	CTGTTGAGCG	TTGGCTTTAT
3781	ACTGGTAAGA	ATTTGTATAA	CGCATATGAT	ACTAAACAGG	CTTTTTCTAG	TAATTATGAT

FIG. 5-1

SUBSTITUTE SHEET

6 / 16

3841	TCCGGTGTTT	ATTCTTATTT	AACGCCTTAT	TTATCACACG	GTCGGTATTT	CAAACCATTA	3900
3901	AATTTAGGTC	AGAAGATGAA	GCTTACTAAA	ATATATTTGA	AAAAGTTTTT	ACGCGTTCCT	3960
3961	TGTCTTGCGA	TTGGATTTGC	ATCAGCATTT	ACATATAGTT	ATATAACCCA	ACCTAAGCCG	4020
4021	GAGGTTAAAA	AGGTAGTCTC	TCAGACCTAT	GATTTTGATA	AATTCATAT	TGACTCTTCT	4080
4081	CAGCGTCTTA	ATCTAAGCTA	TCGCTATGTT	TTCAAGGATT	CTAAGGGAAA	ATTAATTAAT	4140
4141	AGCGACGATT	TACAGAAGCA	AGGTTATTCA	CTCACATATA	TTGATTTATG	TACTGTTTCC	4200
4201	ATTAAAAAGG	TAATTCAAAT	GAATTTGTTA	AATGTAATTA	ATTTTGTTTT	CTTGATGTTT	4260
4261	GTTTCATCAT	CTTCTTTTGC	TCAGGTAATT	GAATGAATA	ATTCGCCTCT	GCGCGATTTT	4320
4321	GTAACCTGGT	ATTCAAAAGCA	ATCAGGCGAA	TCCGTTATTG	TTTCTCCCGA	TGAAAAAGGT	4380
4381	ACTGTTACTG	TATATTCATC	TGACGTTAAA	CCTGAAAATC	TACGCAATTT	CTTTATTTCT	4440
4441	GTTTTACGTG	CTAATAATTT	TGATATGGTT	GGTTCAATTC	CTTCCATTAT	TTAGAAGTAT	4500
4501	AATCCAAACA	ATCAGGATTA	TATTGATGAA	TTGCCATCAT	CTGATAATCA	GGAATATGAT	4560
4561	GATAATTCCG	CTCCTTCTGG	TGGTTTCTTT	GTTCCGCAAA	ATGATAATGT	TACTCAAACCT	4620
4621	TTTTAAAAAT	ATAACGTTTC	GGCAAAGGAT	TTAATACGAG	TTGTGCAATT	GTTTGTAAGT	4680
4681	TCTAATACTT	CTAAATCCTC	AAATGTATTA	TCTATTGACG	GCTCTAATCT	ATTAGTTGTT	4740
4741	AGTGCACCTA	AAGATATTTT	AGATAACCTT	CCTCAATTCC	TTTCTACTGT	TGATTTGCCA	4800
4801	ACTGACCAGA	TATTGATTGA	GGGTTTGATA	TTTGAGGTTT	AGCAAGGTGA	TGCTTTAGAT	4860
4861	TTTTCATTTG	CTGCTGGCTC	TCAGCGTGGC	ACTGTTGCAG	GCGGTGTTAA	TACTGACCGC	4920
4921	CTCACCTCTG	TTTTATCTTC	TGCTGGTGGT	TCGTTCCGTA	TTTTTAATGG	CGATGTTTTA	4980
4981	GGGCTATCAG	TTCCGCGCAT	AAAGACTAAT	AGCCATTCAA	AAATATTGTC	TGTGCCACGT	5040
5041	ATTCTTACGC	TTTCAGGTCA	GAAGGTTTCT	ATCTCTGTTG	GCCAGAATGT	CCCTTTTATT	5100
5101	ACTGGTCGTG	TGACTGGTGA	ATCTGCCAAT	GTAATAATC	CATTTACAGC	GCTTGAGCGT	5160
5161	CAAAATGTAG	GTATTTCCAT	GAGCGTTTTT	CCTGTTGCAA	TGGCTGGCGG	TAATATTGTT	5220
5221	CTGGATATTA	CCAGCAAGGC	CGATAGTTTG	AGTTCTTCTA	CTCAGGCAAG	TGATGTTATT	5280
5281	ACTAATCAAA	GAAGTATTGC	TACAACGGTT	AATTTGCGTG	ATGGACAGAC	TCTTTTACTC	5340
5341	GGTGGCCTCA	CTGATTATAA	AAACACTTCT	CAAGATTCTG	GCGTACCGTT	CCTGTCTAAA	5400
5401	ATCCCTTTAA	TCGGCCTCCT	GTTTAGCTCC	CGCTCTGATT	CCAACGAGGA	AAGCAGGTTA	5460
5461	TACGTGCTCG	TCAAAGCAAC	CATAGTACGC	GCCCTGTAGC	GCGCGATTAA	GCGCGGCGGG	5520
5521	TGTGGTGGTT	ACGCGCAGCG	TGACCGCTAC	ACTTGCCAGC	GCCCTAGCGC	CCGCTCCTTT	5580
5581	CGCTTTCTTC	CCTTCCTTTC	TCGCCACGTT	CGCCGGCTTT	CCCCGTCAAG	CTCTAAATCG	5640
5641	GGGGCTCCCT	TTAGGGTTCC	GATTTAGTGC	TTTACGGCAC	CTCGACCCCA	AAAAACTTGA	5700
5701	TTTGGGTGAT	GGTTCACGTA	GTGGGCCATC	GCCCTGATAG	ACGGTTTTTC	GCCCTTTGAC	5760
5761	GTTGGAGTCC	ACGTTCTTTA	ATAGTGGACT	CTTGTTCCAA	ACTGGAACAA	CACTCAACCC	5820
5821	TATCTCGGGC	TATTCTTTTG	ATTTATAAGG	GATTTTGCCG	ATTTTCGGAAC	CACCATCAAA	5880
5881	CAGGATTTTC	GCCTGCTGGG	GCAAACCAGC	GTGGACCGCT	TGCTGCAACT	CTCTCAGGCG	5940
5941	CAGGCGGTGA	AGGGCAATCA	GCTGTTGCCG	GTCTCGCTGG	TGAAAAGAAA	AACCACCCTG	6000
6001	GCGCCCAATA	CGCAAACCGC	CTCTCCCCGC	GCGTTGGCCG	ATTCATTAAT	GCAGCTGGCA	6060
6061	CGACAGGTTT	CCCGACTGGA	AAGCGGGCAG	TGAGCGCAAC	GCAATTAATG	TGAGTTAGCT	6120
6121	CACCTATTAG	GCACCCACAG	CTTTACACTT	TATGCTTCCG	GCTCGTATGT	TGTGTGGAAT	6180
6181	TGTGAGCGGA	TAACAATTTC	ACACAGGAAA	CAGCTATGAC	CAGGATGTAC	GAATTCGCAG	6240
6241	GTAGGAGAGC	TCGGCGGATC	CTAGGCTGAA	GGCGATGACC	CTGCTAAGGC	TGCATTCAAT	6300
6301	AGTTTACAGG	CAAGTGCTAC	TGAGTACATT	GGCTACGCTT	GGGCTATGGT	AGTAGTTATA	6360
6361	GTTGGTGCTA	CCATAGGGAT	TAAATTATTC	AAAAAGTTTA	CGAGCAAGGC	TTCTTAACCA	6420
6421	GCTGGCGTAA	TAGCGAAGAG	GCCCGCACCG	ATCGCCCTTC	CCAACGATTG	CGCAGCCTGA	6480
6481	ATGGCGAATG	GCGCTTTGCC	TGGTTTCCGG	CACCAGAAGC	GGTGCCGGAA	AGCTGGCTGG	6540
6541	AGTGCGATCT	TCCTGAGGCC	GATACGGTCG	TCCGTCCTTC	AAACTGGCAG	ATGCACGGTT	6600
6601	ACGATGCGCC	CATCTACACC	AACGTAACCT	ATCCCATTAC	GGTCAATCCG	CCGTTTGTTT	6660
6661	CCACGGAGAA	TCCGACGGGT	TGTTACTCGC	TCACATTTAA	TGTTGATGAA	AGCTGGCTAC	6720
6721	AGGAAGGCCA	GACGCGAATT	ATTTTGTATG	GCGTTCCTAT	TGGTTAAAAA	ATGAGCTGAT	6780
6781	TTAACAATAA	TTTAACGCGA	ATTTTAACAA	AATATTAACG	TTTACAATTT	AAATATTTGC	6840
6841	TTATACAATC	TTCTGTGTTT	TGGGGCTTTT	CTGATTATCA	ACCGGGGTAC	ATATGATTGA	6900
6901	CATGCTAGTT	TTACGATTAC	CGTTTCATCGA	TTCTTTGTTT	TGCTCCAGAC	TCTCAGGCAA	6960
6961	TGACCTGATA	GCCTTTGTAG	ATCTCTCAAA	AATAGCTACC	CTCTCCGGCA	TTAATTTATC	7020
7021	AGCTAGAACG	GTTGAATATC	ATATTGATGG	TGATTTGACT	GTCTCCGGCC	TTTCTCACCC	7080
7081	TTTTGAATCT	TTACCTACAC	ATTACTCAGG	CATTGCATTT	AAAATATATG	AGGGTTCTAA	7140
7141	AAATTTTTAT	CCTTGCCTTG	AAATAAAGGC	TTCTCCCGCA	AAAGTATTAC	AGGGTCATAA	7200
7201	TGTTTTGGT	ACAACCGATT	TAGCTTTATG	CTCTGAGGCT	TTATTGCTTA	ATTTTGCTAA	7260
7261	TTCTTTGCCT	TGCCTGTATG	ATTTATTGGA	CGTT			7294

FIG. 5-2

SUBSTITUTE SHEET

7 / 16

	10	20	30	40	50	60
1	AATGCTACTA	CTATTAGTAG	AATTGATGCC	ACCTTTTTCAG	CTCGCGCCCC	AAATGAAAAT
61	ATAGCTAAAC	AGGTTATTGA	CCATTTGCGA	AATGTATCTA	ATGGTCAAAC	TAAATCTACT
121	CGTTCGCAGA	ATTGGGAATC	AACTGTTACA	TGGAATGAAA	CTTCCAGACA	CCGTACTTTA
181	GTTGCATATT	TAAAACATGT	TGAGCTACAG	CACCAGATTC	AGCAATTAAG	CTCTAAGCCA
241	TCTGCAAAAA	TGACCTCTTA	TCAAAAGGAG	CAATTAAGG	TACTCTCTAA	TCCTGACCTG
301	TTGGAGTTTG	CTTCCGGTCT	GGTTGCGTTT	GAAGCTCGAA	TTAAACCGCG	ATATTTGAAG
361	TCTTTCGGGC	TTCTCTTTAA	TCTTTTTGAT	GCAATCCGCT	TTGCTTCTGA	CTATAATAGT
421	CAGGGTAAAG	ACCTGATTTT	TGATTTATGG	TCATTCTCGT	TTTCTGAACT	GTTTAAAGCA
481	TTTGAGGGGG	ATTCAATGAA	TATTTATGAC	GATTCCGCAG	TATTGGACGC	TATCCAGTCT
541	AAACATTTTA	CTATTACCCC	CTCTGGCAAA	ACTTCTTTTG	CAAAAGCCTC	TCGCTATTTT
601	GGTTTTTATC	GTCTGTCTGG	AAACGAGGGT	TATGATAGTG	TTGCTCTTAC	TATGCCTCGT
661	AATTCCTTTT	GGCGTTATGT	ATCTGCTATTA	GTTGAATGTG	GTATTCCTAA	ATCTCAACTG
721	ATGAATCTTT	CTACCTGTAA	TAATGTTGTT	CCGTTAGTTC	GTTTTATTAA	CGTAGATTTT
781	TCTTCCCAAC	GTCTGACTG	GTATAATGAG	CCAGTTCTTA	AAATCGCATA	AGGTAATTCA
841	CAATGATTAA	AGTTGAAATT	AAACCATCTC	AAGCCCAATT	TACTACTCGT	TCTGGTGTTC
901	CTCGTCAGGG	CAAGCCTTAT	TCACTGAATG	AGCAGCTTTG	TTACGTTGAT	TTGGGTAATG
961	AATATCCGGT	TCTTGTCAAG	ATTACTCTTG	ATGAAGGTCA	GCCAGCCTAT	GCGCCTGGTC
1021	TGTACACCGT	TCATCTGTCC	TCTTTCAAAG	TTGGTCAGTT	CGGTTCCCTT	ATGATTTGAC
1081	GTCTGCGCCT	CGTTCCGGCT	AAGTAACATG	GAGCAGGTCG	CGGATTTTCA	CACAATTTAT
1141	CAGGCGATGA	TACAAATCTC	CGTTGTACTT	TGTTTCGCGC	TTGGTATAAT	CGCTGGGGGT
1201	CAAAGATGAG	TGTTTTAGTG	TATTCTTTTCG	CCTCTTTCGT	TTTAGGTTGG	TGCCTTCGTA
1261	GTGGCATTAC	GTATTTTACC	CGTTTAATGG	AAACTTCCTC	ATGAAAAAGT	CTTTAGTCTC
1321	CAAAGCTCTC	GTAGCCGTTG	CTACCTCTCG	TCCGATGCTG	TCTTTCGCTG	CTGAGGGTGA
1381	CGATCCCGCA	AAAGCGGCCT	TAACTCCCT	GCGACGCTCA	GCGACCGAAT	ATATCGGTTA
1441	TGCGTGGGCG	ATGGTTGTTG	TCATTGTCGG	CGCAACTATC	GGTATCAAGC	TGTTTAAAGAA
1501	ATTCACCTCG	AAAGCAAGCT	GATAAACCGA	TACAATTAAT	GGCTCCTTTT	GGAGCCTTTT
1561	TTTTTGGAGA	TTTTCAACGT	GAAAAAATTA	TTATTTCGCA	TTCTTTTAGT	TGTTCTTTTC
1621	TATTTCTACT	CCGCTGAAAC	TGTTGAAAGT	TGTTTAGCAA	AAACCCATAC	AGAAAAATTCA
1681	TTTACTAACG	CTGGAAGAA	CGACAAAAT	TTAGATCGTT	ACGCTAACTA	TGAGGGTTGT
1741	CTGTGGAATG	CTACAGGCGT	TGTAGTTTGT	ACTGGTGACG	AAACTCAGTG	TTACGGTACA
1801	TGGGTTTCTA	TTGGGCTTGC	TATCCCTGAA	AATGAGGGTG	GTGGCTCTGA	GGGTGGCGGT
1861	TCTGAGGGTG	GCGGTTCTGA	GGGTGGCGGT	ACTAAACCTC	CTGAGTACGG	TGATACACCT
1921	ATTCGGGGCT	ATACTTATAT	CAACCTCTCT	GACGGCACTT	ATCCGCTTGG	TACTGAGCAA
1981	AACCCGCTA	ATCCTAATCC	TTCTCTTGAG	GAGTCTCAGC	CTCTTAATAC	TTTCATGTTT
2041	CAGAATAATA	GGTTCCGAAA	TAGGCAAGGG	GCATTAACCTG	TTTATACGGG	CACTGTTACT
2101	CAAGGCACTG	ACCCCGTTAA	AACTTATTAC	CAGTACACTC	CTGTATCATC	AAAAGCCATG
2161	TATGACGCTT	ACTGGAACGG	TAAATTCAGA	GACTGCGCTT	TCCATTCTGG	CTTTAATGAA
2221	GATCCATTCTG	TTTGTGAATA	TCAAGGCCAA	TCGTCTGACC	TGCCTCAACC	TCCTGTCAAT
2281	GCTGGCGGCG	GCTCTGGTGG	TGTTTCTGGT	GGCGGCTCTG	AGGGTGGTGG	CTCTGAGGGT
2341	GGCGGTTCTG	AGGGTGGCGG	CTCTGAGGGA	GGCGGTTCCG	GTGGTGGCTC	TGGTTCCGGT
2401	GATTTTGATT	ATGAAAAGAT	GGCAAAACGT	AATAAGGGGG	CTATGACCGA	AAATGCCGAT
2461	GAAAACGCGC	TACAGTCTGA	CGCTAAAGGC	AAACTTGATT	CTGTGCTAC	TGATACCGGT
2521	GCTGCTATCG	ATGGTTTCAT	TGGTGACGTT	TCCGGCCTTG	CTAATGGTAA	TGGTGCTACT
2581	GGTGATTTTG	CTGGCTCTAA	TTCCCAAATG	GCTCAAGTCG	GTGACGGTGA	TAATTCACCT
2641	TTAATGAATA	ATTTCCGTCA	ATATTTACCT	TCCCTCCCTC	AATCGGTTGA	ATGTCGCCCT
2701	TTTGTCTTTA	GCGCTGGTAA	ACCATATGAA	TTTTCTATTG	ATTGTGACAA	AATAAACTTA
2761	TTCCGTGGTG	TCTTTGCGTT	TCTTTTATAT	GTTGCCACCT	TTATGTATGT	ATTTTCTACG
2821	TTTGCTAAGA	TACTGCGTAA	TAAGGAGTCT	TAATCATGCC	AGTTCTTTTG	GGTATTCCGT
2881	TATTATTGCG	TTTCCTCGGT	TTCTTCTGG	TAACCTTGTG	CGGCTATCTG	CTTACTTTTC
2941	TTAAAAAGGG	CTTCGGTAAG	ATAGCTATTG	CTATTTTCAAT	GTTTCTTGCT	CTTATTATTG
3001	GGCTTAACTC	AATTCTTGTTG	GGTTATCTCT	CTGATATTAG	CGCTCAATTA	CCCTCTGACT
3061	TTGTTTCAGG	TGTTTCAGTTA	ATTCTCCCGT	GTAATGCGCT	TCCCTGTTTT	TATGTTATTC
3121	TCTCTGTAAA	GGCTGCTATT	TTTCTTTTGG	ACGTTAAACA	AAAAATCGTT	TCTTATTTGG
3181	ATTGGGATAA	ATAATATGGC	TGTTTATTTT	TGTTAGTGCA	AATTAGGCTC	TGGAAAGACG
3241	CTCGTTAGCG	TTGGTAAGAT	TAGGATAAAA	ATTGTAGCTG	GGTGCAAAAT	AGCAACTAAT
3301	CTTGATTTAA	GGCTTCAAAA	CCTCCCGCAA	GTCGGGAGGT	TCGCTAAAAC	GCCTCGCGTT
3361	CTTAGAATAC	CGGATAAGCC	TTCTATATCT	GATTTGCTTG	CTATTGGGCG	CGGTAATGAT
3421	TCCTACGATG	AAAATAAAAA	CGGCTTGCTT	GTTCTCGATG	AGTGCGGTAC	TTGGTTTAAAT
3481	ACCCGTTCTT	GGAAATGATA	GGAAAGACAG	CCGATTATTG	ATTGGTTTCT	ACATGCTCGT
3541	AAATTAGGAT	GGGATATTAT	CTTCTTTGTT	CAGGACTTAT	CTATTGTTGA	TAAACAGGCG
3601	CGTTCTGCAT	TAGCTGAACA	TGTTGTTTAT	TGTCGTCGTC	TGGACAGAAAT	TACTTTACCT
3661	TTTGTGCGTA	CTTTATATTC	TCTTATTACT	GGCTCGAAAA	TGCCTCTGCC	TAAATTACAT
3721	GTTGGCGTTG	TTAAATATGG	CGATTCTCAA	TTAAGCCCTA	CTGTTGAGCG	TTGGCTTTAT
3781	ACTGGTAAGA	ATTTGTATAA	CGCATATGAT	ACTAAACAGG	CTTTTCTAG	TAATTATGAT

FIG. 6-1

SUBSTITUTE SHEET

8 / 16

3841	TCCGGTGTTT	ATTCTTATTT	AACGCCTTAT	TTATCACACG	GTCGGTATTT	CAAACCATTA	3900
3901	AATTTAGGTC	AGAAGATGAA	ATTAACATAA	ATATATTTGA	AAAAGTTTTT	TCGCGTTCTT	3960
3961	TGTCCTTGC	TTGGATTTGC	ATCAGCATTT	ACATATAGTT	ATATAACCCA	ACCTAAGCCG	4020
4021	GAGGTTAAAA	AGGTAGTCTC	TCAGACCTAT	GATTTTGATA	AATTCATAT	TGACTCTTCT	4080
4081	CAGCGTCTTA	ATCTAAGCTA	TCGCTATGTT	TTCAAGGATT	CTAAGGGAAA	ATTAATTAAT	4140
4141	AGCGACGATT	TACAGAAGCA	AGGTTATTCA	CTCACATATA	TTGATTTATG	TACTGTTTCC	4200
4201	ATTAATAAAG	GTAATTCAAA	TGAAATTGTT	AAATGTAATT	AATTTTGTTT	TCTTGATGTT	4260
4261	TGTTTACGTC	TCTTCTTTTG	CTCAGGTAAT	TGAAATGAAT	AATTCGCCTC	TGCGCGATT	4320
4321	TGTAACCTGG	TATTCAAAAGC	AATCAGGCGA	ATCCGTTATT	GTTTCTCCCG	ATGTAAAAGG	4380
4381	TACTGTTACT	GTATATTCA	CTGACGTTAA	ACCTGAAAAT	CTACGCAATT	TCTTTATTTT	4440
4441	TGTTTTACGT	GCTAATAATT	TTGATATGGT	TGGTTCAATT	CCTTCCATAA	TTCAGAAGTA	4500
4501	TAATCCAAAC	AATCAGGATT	ATATTGATGA	ATTGCCATCA	TCTGATAATC	AGGAATATGA	4560
4561	TGATAAATCC	GCTCCTTCTG	GTGGTTTCTT	TGTTCCGCAA	AATGATAATG	TTACTCAAAC	4620
4621	TTTTAAAAAT	AATAACGTTT	GGGCAAAGGA	TTTAATACGA	GTTGTCGAAT	TGTTTGTAAT	4680
4681	TGCTAATACT	TCTAAATCCT	CAAAATGTAT	ATCTATTGAC	GGCTCTAATC	TATTAGTTGT	4740
4741	TAGTGACCTT	AAAGATATTT	TAGATAACCT	TCTCAATTC	CCTTCTACTG	TTGATTTGCC	4800
4801	AACTGACCAG	ATATTGATTG	AGGGTTTGAT	ATTTGAGGTT	CAGCAAGGTG	ATGCTTTAGA	4860
4861	TTTTTCATTT	GCTGCTGGCT	CTCAGCGTGG	CACGTGTTGA	GGCGGTGTTA	ATACTGACCG	4920
4921	CCTCACCTCT	GTTTTATCTT	CTGCTGGTGG	TTCGTTCCGG	ATTTTTAATG	GCGATGTTTT	4980
4981	AGGGCTATCA	GTTTCGCGAT	TAAAGACTAA	TAGCCATTCA	AAAATATTGT	CTGTGCCACG	5040
5041	TATTTCTTACG	CTTTCAGGTC	AGAAGGGTTT	TATCTCTGTT	GGCCAGAATG	TCCCTTTTAT	5100
5101	TACTGGTCGT	GTGACTGGTG	AATCTGCCAA	TGTAATAAAT	CCATTTTACA	CGATTGAGCG	5160
5161	TCAAAATGTA	GGTATTTCCA	TGAGCGTTTT	TCCTGTTGCA	ATGGCTGGCG	GTAATATTGT	5220
5221	TCTGGATATT	ACCAGCAAGG	CCGATAGTTT	GAGTCTTCT	ACTCAGGCAA	GTGATGTTAT	5280
5281	TACTAATCAA	AGAAGTATTG	CTACAACGGT	TAATTTGCGT	GATGGACAGA	CTCTTTTACT	5340
5341	CGGTGGCCTC	ACTGATTATA	AAAACACTTC	TCAAGATTCT	GGCGTACCGT	TCCTGTCTAA	5400
5401	AATCCCTTTA	ATCGGCCTCC	TGTTAGCTC	CGCGCTGTAT	TCCAACGAGG	AAAGCACGTT	5460
5461	ATACGTGCTC	GTCAAAGCAA	CCATAGTACG	CGCCCTGTAG	CGGCGCATTA	AGCGCGGCGG	5520
5521	GTGTGGTGGT	TACGCGCAGC	GTGACCGCTA	CACCTGCCAG	CGCCCTAGCG	CCCGCTCCTT	5580
5581	TCGCTTTCTT	CCCTTCCCTT	CTCGCCACGT	TCGCCGGCTT	TCCCGTCAA	GCTCTAAATC	5640
5641	GGGGGCTCCC	TTTAGGGTTC	CGATTTAGTG	CTTTACGGCA	CCTCGACCCC	AAAAAACTTG	5700
5701	ATTTGGGTGA	TGGTTCACGT	AGTGGGCCAT	CGCCCTGATA	GACGGTTTTT	CGCCCTTTGA	5760
5761	CTTTGGAGTC	CACGTTCTTT	AATAGTGGAC	TCTTGTTCCT	AACTGGAACA	ACACTCAACT	5820
5821	CTATCTCGGG	CTATTCTTTT	GATTTATAAG	GGATTTTGCC	GATTTGCGAA	CCACCATCAA	5880
5881	ACAGGATTTT	CGCCTGCTGG	GGCAAACAG	CGTGGACCGC	TTGCTGCAAC	TCTCTCAGGG	5940
5941	CCAGGCGGTG	AAGGGCAATC	AGCTGTTGCC	CGTCTCGCTG	GTGAAAAGAA	AAACCACCCT	6000
6001	GGCGCCCAAT	ACGCAAAACG	CCTCTCCCGG	CGCGTTGGCC	GATTCATTAA	TGCAGCTGGC	6060
6061	ACGACAGGTT	TCCCGACTGG	AAAGCGGGCA	GTGAGCGCAA	CGCAATTAAT	GTGAGTTAGC	6120
6121	TCACTCATTA	GGCACCCAG	GCTTTACACT	TTATGCTTCC	GGCTCGTATG	TTGTGTGGAA	6180
6181	TTGTGAGCGG	ATAACAATTT	CACAGGCCAA	GGAGACAGTC	ATAATGAAAT	ACCTATTGCC	6240
6241	TACGGCAGCC	GCTGGATTGT	TATTACTCGC	TGCCCAACCA	GCCATGGCCG	AGCTCGTGAT	6300
6301	GACCCAGACT	CCAGAATTCC	ATCCGGAATG	AGTGTTAATT	CTAGAACGCG	TAAGCTTGGC	6360
6361	ACTGGCCGTC	GTTTTACAAC	GTCGTGACTG	GGAAAACCCCT	GGCGTTACCC	AACTTAATCG	6420
6421	CCTTGACAGCA	CACCCCTT	TCGCCAGCTG	GCGTAATAGC	GAAGAGGCCC	GCACCGATCG	6480
6481	CCCTTCCCAA	CAGTTGCGCA	GCCTGAATGG	CGAATGGCGC	TTTGCTGGT	TTCCGGCACC	6540
6541	AGAAGCGGTG	CCGAAAAGCT	GGCTGGAGTG	CGATCTTCCT	GAGGCCGATA	CGGTGCTCGT	6600
6601	CCCCTCAAAC	TGGCAGATGC	ACGGTTACGA	TGCGCCCATC	TACACCAACG	TAACCTATCC	6660
6661	CATTACGGTC	AATCCGCCGT	TTGTTCCAC	GGAGAATCCG	TCGGGTGTTT	ACTCGCTCAC	6720
6721	ATTTAATGTT	GATGAAAGCT	GGCTACAGGA	AGGCCAGACG	CGAATTATTT	TTGATGGCGT	6780
6781	TCCTATTGGT	TAAAAATGA	GCTGATTTAA	CAAAAATTTA	ACGCGAATTT	TAACAAAATA	6840
6841	TTAACGTTTA	CAATTTAAAT	ATTTGCTTAT	ACAATCTTCC	TGTTTTTGGG	GCTTTTCTGA	6900
6901	TTATCAACCG	GGGTACATAT	GATTGACATG	CTAGTTTTAC	GATTACCGTT	CATCGATTCT	6960
6961	CTTGTTTGCT	CCAGACTCTC	AGGCAATGAC	CTGATAGCCT	TTGTAGATCT	CTCAAAAATA	7020
7021	GCTACCCCTCT	CCGGCATTAA	TTTATCAGCT	AGAACGGTTG	AATATCATAT	TGATGGTGAT	7080
7081	TTGACTGTCT	CCGGCCTTTC	TCACCCTTTT	GAATCTTTAC	CTACACATTA	CTCAGGCATT	7140
7141	GCATTTAAAA	TATATGAGGG	TTCTAAAAAT	TTTTATCCTT	GCGTTGAAAT	AAAGGCTTCT	7200
7201	CCGCAAAAAG	TATTACAGGG	TCATAATGTT	TTTGGTACAA	CCGATTTAGC	TTTATGCTCT	7260
7261	GAGGCTTTAT	TGCTTAATTT	TGCTAATTTT	TTGCTTGCC	TGTATGATTT	ATTGGACGTT	7320

FIG. 6-2

9 / 16

	10	20	30	40	50	60	
1	AATGCTACTA	CTATTAGTAG	AATTGATGCC	ACCTTTTCAG	CTCGCGCCCC	AAATGAAAAAT	60
61	ATAGCTAAAC	AGGTTATTGA	CCATTTGCGA	AATGTATCTA	ATGGTCAAAC	TAAATCTACT	120
121	CGTTCGCAGA	ATTGGGAATC	AACTGTTACA	TGGAATGAAA	CTTCCAGACA	CCGTACTTTA	180
181	GTTGCATATT	TAAAACATGT	TGAGCTACAG	CACCAGATTG	AGCAATTAAG	CTCTAAGCCA	240
241	TCTGCAAAAA	TGACCTCTTA	TCAAAAGGAG	CAATTAAGG	TACTCTCTAA	TCCTGACCTG	300
301	TTGGAGTTTG	CTTCCGGTCT	GGTTCGCTTT	GAAGCTCGAA	TTAAACGCGC	ATATTTGAAG	360
361	TCTTTCGGGC	TTCCTCTTAA	TCTTTTTGAT	GCAATCCGCT	TIGCTTCTGA	CTATAATAGT	420
421	CAGGGTAAAG	ACCTGATTTT	TGATTTATGG	TCATTCTCGT	TTTCTGAACT	GTTTAAAGCA	480
481	TTTGAGGGGG	ATTCAATGAA	TATTTATGAC	GATTCCGCAG	TATTGGACGC	TATCCAGTCT	540
541	AAACATTTTA	CTATTACCCC	CTCTGGCAAA	ACTTCTTTTG	CAAAAGCCTC	TCGCTATTTT	600
601	GGTTTTTATC	GTCGTCTGGT	AAACGAGGGT	TATGATAGTG	TTGCTCTTAC	TATGCCTCGT	660
661	AATTCCTTTT	GGCGTTATGT	ATCTGCATTA	GTTGAATGTG	GTATTCCTAA	ATCTCAACTG	720
721	ATGAATCTTT	CTACCTGTAA	TAATGTTGTT	CCGTTAGTTC	GTTTTATTAA	CGTAGATTTT	780
781	TCTTCCCAAC	GTCCTGACTG	GTATAATGAG	CCAGTTCTTA	AAATCGCATA	AGGTAATTCA	840
841	CAATGATTAA	AGTTGAAATT	AAACCATCTC	AAGCCCAATT	TACTACTCGT	TCTGGTGTTT	900
901	CTCGTCAGGG	CAAGCCTTAT	TCACTGAATG	AGCAGCTTTG	TTACGTTGAT	TTGGGTAATG	960
961	AATATCCGGT	TCTTGTCAAG	ATTACTCTTG	ATGAAGGTCA	GCCAGCCTAT	GCGCCTGGTC	1020
1021	TGTACACCGT	TCATCTGTCC	TCTTTCAAAG	TTGGTCAGTT	CGGTTCCCTT	ATGATTGACC	1080
1081	GTCTGCGCCT	CGTTCGGGCT	AAGTAACATG	GAGCAGGTCG	CGGATTTCTG	CACAATTTAT	1140
1141	CAGGCGATGA	TACAAATCTC	CGTTGTACTT	TGTTTCGCGC	TTGGTATAAT	CGCTGGGGGT	1200
1201	CAAAGATGAG	TGTTTTAGTG	TATTTCTTTC	CCTCTTTCTG	TTTAGGTTGG	TGCTTTCGTA	1260
1261	GTGGCATTAC	GTATTTTACC	CGTTTAATGG	AAACTTCCTC	ATGAAAAAGT	CTTTAGTCCT	1320
1321	CAAAGCCTCT	GTAGCCGTTG	CTACCCTCGT	TCCGATGCTG	TCTTTCGCTG	CTGAGGGTGA	1380
1381	CGATCCCGCA	AAAGCGGCCCT	TTAACTCCCT	GCAAGCCTCA	GCGACCGAAT	ATATCGGTTA	1440
1441	TGCGTGGGGC	ATGGTTGTTG	TCATTGTGCG	CGCAACTATC	GGTATCAAGC	TGTTTAAGAA	1500
1501	ATTCACTCTG	AAAGCAAGCT	GATAAAACCG	TACAATTAAT	GGCTCCTTTT	GGAGCCTTTT	1560
1561	TTTTTGGAGA	TTTTCAACGT	GAAAAAATTA	TTATTCGCAA	TTCTTTAGT	TTCTCTTTC	1620
1621	TATTCTCACT	CCGCTGAAAC	TGTTGAAAGT	TGTTTAGCAA	AACCCCATAC	AGAAAATTCA	1680
1681	TTTACTAACG	TCTGGAAAGA	CGACAAAAC	TTAGATCGTT	ACGCTAACTA	TGAGGGTTGT	1740
1741	CTGTGGAATG	CTACAGGCCT	TGTAGTTTGT	ACTGGTGACG	AAACTCAGTG	TTACGGTACA	1800
1801	TGGGTTCCCT	TTGGGCTTGC	TATCCCTGAA	AATGAGGGTG	GTGGCTCTGA	GGGTGGCGGT	1860
1861	TCTGAGGGTG	GCGGTTCTGA	GGGTGGCGGT	ACTAAACCTC	CTGAGTACGG	TGATACACCT	1920
1921	ATTCGGGGCT	ATACTTATAT	CAACCCTCTC	CAGCGCACTT	ATCCGCCTGG	TACTAGCAA	1980
1981	AACCCCGCTA	ATCCTAATCC	TTCTCTTGAG	GAGTCTCAGC	CTCTTAATAC	TTTCATGTTT	2040
2041	CAGAATAATA	GGTTCCGAAA	TAGGCAGGGG	GCATTAAC	TTTATACGGG	CACTGTTACT	2100
2101	CAAGGCACTG	ACCCCGTTAA	AACCTTATTAC	CAGTACACTC	CTGTATCATC	AAAAGCCATG	2160
2161	TATGACGCTT	ACTGGAACGG	TAAATTGAGA	GACTGCGCTT	TCCATTCTGG	CTTTAATGAA	2220
2221	GATCCATTCT	TTTGTGAATA	TCAAGGCCAA	TCGTCTGACC	TGCCTCAACT	TCTGTCAAT	2280
2281	GCTGGCGGGC	GCTCTGGTGG	TGTTTCTGGT	GGCGGCTCTG	AGGGTGGTGG	CTCTGAGGGT	2340
2341	GGCGGTTCTG	AGGGTGGCGG	CTCTGAGGGA	GGCGGTTCCG	GTGGTGGCTC	TGGTTCGGGT	2400
2401	GATTTTGATT	ATGAAAAGAT	GGCAAACGCT	AATAAGGGGG	CTATGACCGA	AAATGCCGAT	2460
2461	GAAAACGCGC	TACAGTCTGA	CGCTAAAGGC	AAACTTGATT	CTGTCGCTAC	TGATTACGGT	2520
2521	GCTGCTATCG	ATGGTTTCAT	TGGTGACGTT	TCCGGCCTTG	CTAATGGTAA	TGGTGCTACT	2580
2581	GGTGATTTTG	CTGGCTCTAA	TTCCCAATG	TCCCAAGTCG	GTGACGGTGA	TAATTACACT	2640
2641	TTAATGAATA	ATTTCCGTCA	ATATTTACCT	TCCCTCCCTC	AATCGGTTGA	ATGTCGCCCT	2700
2701	TTTGTCTTTA	GCGCTGGTAA	ACCATATGAA	TTTTCTATTG	ATTGTGACAA	AATAAACTTA	2760
2761	TTCCGTGGTG	TCTTTGCGTT	TCTTTTATAT	GTTGCCACCT	TTATGTATGT	ATTTTCTACG	2820
2821	TTTGCTAACA	TACTGCGTAA	TAAGGAGTCT	TAATCATGCC	AGTTCTTTTG	GGTATTCCGT	2880
2881	TATTATTGCG	TTTCTCTGGT	TTCTTTCTGG	TAACTTTGTG	CGGCTATCTG	CTTACTTTTC	2940
2941	TTAAAAAGGG	CTTCGGTAAG	ATAGCTATTG	CTATTTCTAT	GTTTCTTGCT	CTTATTATTG	3000
3001	GGCTTAACCT	AATTCTTGTT	GGTTATCTCT	CTGATATTAG	CGCTCAATTA	CCCTCTGACT	3060
3061	TTGTTCAAGG	TGTTCAAGTA	ATTCTCCCGT	CTAATGCGCT	TCCCTGTTTT	TATGTTATTC	3120
3121	TCTCTGTAAA	GGCTGCTATT	TTCATTTTTG	ACGTTAAACA	AAAAATCGTT	TCCTATTTGG	3180
3181	ATTGGGATAA	ATAATATGGC	TGTTTATTTT	GTAAGTGGCA	AATTAGGCTC	TGGAAAGACG	3240
3241	CTCGTTAGCG	TTGGTAAGAT	TCAGGATAAA	ATTGTAGCTG	GGTGCAAAAT	AGCAACTAAT	3300
3301	CTTGATTTAA	GGCTTCAAAA	CCTCCCGCAA	GTCGGGAGGT	TCGCTAAAAA	GCCTCGCGTT	3360
3361	CTTAGAATAC	CGGATAAGCC	TTCTATATCT	GATTTGCTTG	CTATTGGGCG	CGGTAATGAT	3420
3421	TCCTACGATG	AAAATAAAAA	CGGCTTGCTT	GTTCTCGATG	AGTGCGGTAC	TTGGTTTAAAT	3480
3481	ACCCGTTCTT	GGAATGATAA	GGAAAGACAG	CCGATTATTG	ATTGGTTTCT	ACATGCTCGT	3540
3541	AAATTAGGAT	GGGATATTAT	TTTTCTTGTT	CAGGACTTAT	CTATTGTTGA	TAAACAGGCG	3600
3601	CGTTCTGCAT	TAGCTGAACA	TGTTGTTTAT	TGTCGTCGTC	TGGACAGAAT	TACTTTACCT	3660
3661	TTTGTGCGGT	CTTTATATTC	TCTTATTACT	GGCTCGAAAA	TGCCTCTGCC	TAAATTACAT	3720
3721	GTTGGCGTTG	TTAAATATGG	CGATTCTCAA	TTAAGCCCTA	CTGTTGAGCG	TTGGCTTTAT	3780

FIG. 7-1

SUBSTITUTE SHEET

10 / 16

3781	ACTGGTAAGA	ATTTGTATAA	CGCATATGAT	ACTAAACAGG	CTTTTTCTAG	TAATTATGAT	3840
3841	TCCGGTGTTT	ATTCTTATTT	AACGCCTTAT	TTATCACACG	GTCGGTATTT	CAAACCATTA	3900
3901	AATTTAGGTC	AGAAGATGAA	GCTTACTAAA	ATATATTTGA	AAAAGTTTTT	ACGCGTTCTT	3960
3961	TGTCTTGCGA	TTGGATTTGC	ATCAGCATTT	ACATATAGTT	ATATAACCCA	ACCTAAGCCG	4020
4021	GAGGTTAAAA	AGGTAGTCTC	TCAGACCTAT	GATTTTGATA	AATTCACAT	TGACTCTTCT	4080
4081	CAGCGTCTTA	ATCTAAGCTA	TCGCTATGTT	TTCAAGGATT	CTAAGGGAAA	ATTAATTAAT	4140
4141	AGCGACGATT	TACAGAAGCA	AGGTTATTCA	CTCACATATA	TTGATTTATG	TACTGTTTCC	4200
4201	ATTAATAAAG	GTAATTCAAA	TGAAATTGTT	AAATGTAATT	AATTTTGTTT	TCTTGATGTT	4260
4261	TGTTTCATCA	TCTTCTTTTG	CTCAGGTAAT	TGAAATGAAT	AATTCGCCTC	TGCGCGATT	4320
4321	TGTAACCTGG	TATTCAAAGC	AATCAGGCGA	ATCCGTTATT	GTTTCTCCCG	ATGTAATAAGG	4380
4381	TACTGTTACT	GTATATTCAT	CTGACGTTAA	ACCTGAAAT	CTACGCAATT	TCTTTATTTT	4440
4441	TGTTTTACGT	GCTAATAAAT	TTGATATGGT	TGGTTCAATT	CCTTCCATAA	TTCAGAAGTA	4500
4501	TAATCCAAAC	AATCAGGATT	ATATTGATGA	ATTGCCATCA	TCTGATAATC	AGGAATATGA	4560
4561	TGATAATTCC	GCTCCTTCTG	GTGGTTTCTT	TGTTCCGCAA	AATGATAATG	TTACTCAAAC	4620
4621	TTTTAAATTT	AATAACGTTT	GGGCAAAGGA	TTTAATACGA	GTTGTCGAAT	TGTTTGTAAT	4680
4681	GTCTAATACT	TCTAAATCCT	CAAATGTATT	ATCTATTGAC	GGCTCTAATC	TATTAGTTGT	4740
4741	TAGTGCACCT	AAAGATATTT	TAGATAACCT	TCCTCAATTC	CTTCTACTG	TTGATTTGCC	4800
4801	AACTGACCA	ATATTGATTG	AGGGTTTGAT	ATTTGAGGTT	CAGCAAGGTT	ATGCTTTAGA	4860
4861	TTTTTCATTT	GCTGCTGGCT	CTCAGCGTGG	CACCTGTTGA	GGCGGTGTTA	ATACTGACCG	4920
4921	CCTCACCTCT	GTTTTATCTT	CTGCTGGTGG	TTGTTCCGGT	ATTTTTAATG	GCGATGTTTT	4980
4981	AGGGCTATCA	GTTTCGCGAT	TAAAGACTAA	TAGCCATTCA	AAAATATTGT	CTGTGCCACG	5040
5041	TATTTCTACG	CTTTCAGGTC	AGAAGGGTTC	TATCTCTGTT	GGCCAGAATG	TCCCTTTTAT	5100
5101	TACTGGTCGT	GTGACTGGTG	AATCTGCCAA	TGTAATAAAT	CCATTTTCTA	CGATTGAGCG	5160
5161	TCAAAATGTA	GGTATTTCCA	TGAGCGTTTT	TCCTGTTGCA	ATGGCTGGCG	GTAAATTTGT	5220
5221	TCTGGATATT	ACCAGCAAGG	CCGATAGTTT	GAGTTCTTCT	ACTCAGGCAA	GTGATGTTAT	5280
5281	TACTAATCAA	AGAAGTATTG	CTACAACGGT	TAATTTGCGT	GATGGACAGA	CTCTTTTACT	5340
5341	CGGTGGCCTC	ACTGATTATA	AAAACACTTC	TCAAGATTCT	GGCGTACCGT	TCCTGTCTAA	5400
5401	AATCCCTTTA	ATCGGCCTCC	TGTTTAGCTC	CCGCTCTGAT	TCCAACGAGG	AAAGCACGTT	5460
5461	ATACGTGCTC	GTCAAAGCAA	CCATAGTACG	CGCCCTGTAG	CGGCGCATTA	AGCGCGGCGG	5520
5521	TGTTGGTGCT	TACGCGCAGC	GTGACGCTGA	CACCTGCCAG	CGCCCTACCG	CCCGCTCCTT	5580
5581	TCGCTTTCTT	CCCTTCCTTT	CTCGCCACGT	TCGCGGCTT	TCCCGTCAA	GCTCTAAATC	5640
5641	GGGGGCTCCC	TTTAGGGTTC	CGATTTAGTG	CTTTACGGCA	CCTCGACCCC	AAAAAACTTG	5700
5701	ATTTGGGTGA	TGGTTCACGT	AGTGGGCCAT	CGCCCTGATA	GACGGTTTTT	CGCCCTTTGA	5760
5761	CGTTGGAGTC	CACGTTCTTT	AATAGTGGAC	TCTTGTTCCA	AACTGGAACA	ACACTCAACC	5820
5821	CTATCTCGGG	CTATTCTTTT	GATTTATAAG	GGATTTTGCC	GATTTTCGGA	CCACCATCAA	5880
5881	ACAGGATTTT	CGCCTGCTGG	GGCAAACCG	CGTGGACCGC	TTGCTGCAAC	TCTCTCAGGG	5940
5941	CCAGGCGTTG	AAGGGCAATC	AGCTGTTGCC	CGTCTCGCTG	GTGAAAAGAA	AAACCAACCT	6000
6001	GGCGCCCAAT	ACGCAAACCG	CCTCTCCCCG	CGCGTTGGCC	GATTCATTAA	TGCAGCTGGC	6060
6061	ACGACAGGTT	TCCCGACTGG	AAAGCGGGCA	GTGAGCGCAA	CGCAATTAAT	GTGAGTTAGC	6120
6121	TCACTCATT	GGCACCCAG	GCTTTACACT	TTATGCTTCC	GGCTCGTATG	TTGTGTGGAA	6180
6181	TTGTGAGCGG	ATAACAATTT	CACACGCGTC	ACTTGGCACT	GGCCGTCGTT	TTACAACGTC	6240
6241	GTGACTGGGA	AAACCCTGGC	GTTACCCAAG	CTTTGTACAT	GGAGAAAATA	AAGTGAACA	6300
6301	ATGACACTATT	GCACTGGCAC	TCTTACCGTT	ACCGTTACTG	TTTACCCCTG	TGACAAAAGC	6360
6361	CGCCAGGTC	CAGCTGCTCG	AGTCAGGCCT	ATTGTGCCCA	GGGGATTGTA	CTAGTGGATC	6420
6421	CTAGGCTGAA	GGCGATGACC	CTGCTAAGGC	TGCATTCAAT	AGTTTACAGG	CAAGTGCTAC	6480
6481	TGAGTACATT	GGCTACGCTT	GGGCTATGGT	AGTAGTTATA	GTTGGTGCTA	CCATAGGGAT	6540
6541	TAAATTATTC	AAAAAGTTTA	CGAGCAAGGC	TTCTTAAGCA	ATAGCGAAGA	GGCCCGCACC	6600
6601	GATCGCCCTT	CCCAACAGTT	GGCGAGCCTG	AATGGCGAAT	GGCGCTTTGC	CTGGTTTCCG	6660
6661	GCACCAAGAG	CGGTGCCGGA	AAGCTGGCTG	GAGTGCAGTC	TTCTGAGGCG	CGATACGGTC	6720
6721	GTCGTCCCTT	CAAACCTGGC	GATGCACGGT	TACGATGCGC	CCATCTACAC	CAACGTAACC	6780
6781	TATCCCATTA	CGGTCAATCC	GCCGTTTGTT	CCCACGGAGA	ATCCGACGGG	TTGTTACTCG	6840
6841	CTCACATTTA	ATGTTGATGA	AAGCTGGCTA	CAGGAAGGCC	AGACGCGAAT	TATTTTTGAT	6900
6901	GGCGTTTCTA	TTGGTTAAAA	AATGAGCTGA	TTTAACAAAA	ATTTAACGCG	AATTTTAACA	6960
6961	AAATATTAAC	GTTTACAATT	TAAATATTTG	CTTATACAAT	CTTCTGTTT	TTGGGGCTTT	7020
7021	TCTGATTATC	AACCGGGGTA	CATATGATTG	ACATGCTAGT	TTTACGATTA	CCGTTTCTCG	7080
7081	ATTCTCTTGT	TTGCTCCAGA	CTCTCAGGCA	ATGACCTGAT	AGCCTTTGTA	GATCTCTCAA	7140
7141	AAATAGCTAC	CCTCTCCGGC	ATTAATTTAT	CAGCTAGAAC	GGTTGAATAT	CATATTGATG	7200
7201	GTGATTTGAC	TGTCTCCGGC	CTTTCTCACC	CTTTTGAATC	TTTACCTACA	CATTACTCAG	7260
7261	GCATTGCATT	TAAAATATAT	GAGGGTCTTA	AAAATTTTTA	TCCTTGCGTT	GAAATAAAGG	7320
7321	CTTCTCCCGC	AAAAGTATTA	CAGGGTCATA	ATGTTTTTGG	TACAACCGAT	TTAGCTTTAT	7380
7381	GCTCTGAGGC	TTTATTGCTT	AATTTTGCTA	ATTCTTTGCC	TTGCCTGTAT	GATTTATTGG	7440
7441	ACGTT						7445

FIG. 7-2

SUBSTITUTE SHEET

11 / 16

	10	20	30	40	50	60	
1	AATGCTACTA	CTATTAGTAG	AATTGATGCC	ACCTTTTCAG	CTCGCGCCCC	AAATGAAAAT	60
61	ATAGCTAAAC	AGGTTATTGA	CCATTTGCGA	AATGTATCTA	ATGGTCAAAC	TAAATCTACT	120
121	CGTTTCGAGA	ATTGGGAATC	AAGTGTACAC	TGGAATGAAA	CTTCCAGACA	CCGTACTTTA	180
181	GTTGCATATT	TAAAACATGT	TGAGCTACAG	CACCAGATTG	AGCAATTAAG	CTCTAAGCCA	240
241	TCTGCAAAAA	TGACCTCTTA	TCAAAAGGAG	CAATTAAGG	TACTCTCTAA	TCCTGACCTG	300
301	TTGGAGTTTG	CTTCCGGTCT	GGTTTCGCTT	GAAGCTCGAA	TTAAAACGCG	ATATTTGAAG	360
361	TCTTTTCGGGC	TTCTCTTTAA	TCTTTTTGAT	GCAATCCGCT	TTGCTTCTGA	CTATAATAGT	420
421	CAGGGTAAAG	ACCTGATTTT	TGATTTATGG	TCATTCTCGT	TTTCTGAACT	GTTTAAAGCA	480
481	TTTGAGGGGG	ATTCAATGAA	TATTTATGAC	GATTCGCGAG	TATTGGACGC	TATCCAGTCT	540
541	AAACATTTTA	CTATTACCCC	CTCTGGCAAA	ACTTCTTTTG	CAAAAGCCTC	TCGCTATTTT	600
601	GGTTTTTATC	GTCGTCTGGT	AAACGAGGGT	TATGATAGTG	TTGCTCTTAC	TATGCCTCGT	660
661	AATTCCTTTT	GGCGTTATGT	ATCTGCATTA	GTTGAATGTG	GTATTCCTAA	ATCTCAACTG	720
721	ATGAATCTTT	CTACCTGTAA	TAATGTTGTT	CCGTTAGTTC	GTTTTATTAA	CGTAGATTTT	780
781	TCTTCCCAAC	GTCTGACTG	GTATAATGAG	CCAGTCTTAA	AAATCGCATA	AGGTAATTCA	840
841	CAATGATTAA	AGTTGAAATT	AAACCATCTC	AAGCCCAATT	TACTACTCGT	TCTGGTGTTC	900
901	CTCGTCAGGG	CAAGCCTTAT	TCACTGAATG	AGCAGCTTTG	TTACGTTGAT	TTGGGTAATG	960
961	AATATCCGGT	TCTTGTCAAG	ATTACTCTTG	ATGAAGGTCA	GCCAGCCTAT	GCGCCTGGTC	1020
1021	TGTACACCGT	TCATCTGTCC	TCTTTCAAAG	TTGGTCAGTT	CGGTTCCCTT	ATGATTGACC	1080
1081	GTCTGCGCCT	CGTTCCGGCT	AAGTAACATG	GAGCAGGTCTG	CGGATTTCTGA	CACAATTTAT	1140
1141	CAGGCGATGA	TACAAATCTC	CGTTGTACTT	TGTTTCGCGC	TTGGTATAAT	CGCTGGGGGT	1200
1201	CAAGATGAG	TGTTTTAGTG	TATTTCTTTCG	CCTCTTTCTG	TTTAGGTTGG	TGCCTTCGTA	1260
1261	GTGGCATTAC	GTATTTTACC	CGTTTAAATG	AAACTTCCTC	ATGAAAAAGT	CTTTAGTCTT	1320
1321	CAAGCCTCT	GTAGCCGTG	CTACCCTCGT	TCCGATGCTG	TCTTTCGCTG	CTGAGGTTGA	1380
1381	CGATCCCGCA	AAAGCGGCCT	TTAACTCCCT	GCAAGCCTCA	GCGACCGAAT	ATATCGGTTA	1440
1441	TGCGTGGGCG	ATGGTTGTTG	TCATTGTCGG	CGCAACTATC	GGTATCAAGC	TGTTTAAGAA	1500
1501	ATTCACCTCG	AAAGCAAGCT	GATAAACCGA	TACAATTAAT	GGCTCCTTTT	GGAGCCTTTT	1560
1561	TTTTTGGAAG	TTTTCAACGT	GAAAAAATTA	TTATTCGCAA	TTCTTTTAGT	TGTTCTTTTC	1620
1621	TATTTCTACT	CCGCTGAAAC	TGTTGAAAGT	TGTTTAGCAA	AACCCCATAC	AGAAAAATTA	1680
1681	TTTTACTAACG	TCTGAAAGAG	CGACAAAGCT	TTAGATCGTT	ACGCTAACTA	TGAGGTTGTG	1740
1741	CTGTGGAATG	CTACAGGCGT	TGTAGTTTGT	ACTGGTGACG	AAACTCAGTG	TTACGGTACA	1800
1801	TGGGTTCTTA	TTGGGCTTGC	TATCCCTGAA	AATGAGGGTG	GTGGCTCTGA	GGGTGGCGGT	1860
1861	TCTGAGGGTG	GCGGTTCTGA	GGGTGGCGGT	ACTAAACCTC	CTGAGTACGG	TGATACACCT	1920
1921	ATTCGCGGCT	ATACTTATAT	CAACCCTCTC	GACGGCACTT	ATCCGCTGCG	TACTGAGCAA	1980
1981	AACCCCGCTA	ATCCTAATCC	TTCTCTTGAG	GAGTCTCAGC	CTCTTAATAC	TTTCTATGTT	2040
2041	CAGAATAATA	GTTTCCGAAA	TAGGCAGGGG	GCATTAACGT	TTTATACGGG	CACCTGTTACT	2100
2101	CAAGGCACTG	ACCCCGTTAA	AACCTATTAC	CAGTACACTC	CTGTATCATT	AAAAGCCATG	2160
2161	TATGACGCTT	ACTGGAACGG	TAAATTCAGA	GACTGCGCTT	TCCATTCTGG	CTTTAATGAA	2220
2221	GATCCATTCTG	TTTGTGAATA	TCAAGGCCAA	TCGTCTGACC	TGCCTCAACC	TCCTGTCAAT	2280
2281	GCTGGCGGCG	CGTCTGGTGG	TGGTCTGGT	GGCGGCTCTG	AGGGTGGTGG	CTCTGAGGGT	2340
2341	GGCGGTTCTG	AGGGTGGCGG	CTCTGAGGGA	GGCGGTTCCG	GTGGTGGCTC	TGGTTCCGGT	2400
2401	GATTTTGATT	ATGAAAAGAT	GGCAAAACGT	AAATAGGGGG	CTATGACCGA	AAATGCCGAT	2460
2461	GAAAACGCGC	TACAGTCTGA	CGCTAAAGGC	AAACTTGATT	CTGTGCTAC	TGATTACGGT	2520
2521	GCTGCTATCG	ATGGTTTCAT	TGGTGACGTT	TCCGGCCTTG	CTAATGGTAA	TGGTGCTACT	2580
2581	GGTGATTTTG	CTGGCTCTAA	TTCCCAAATG	GCTCAAGTCG	GTGACGGTGA	TAATTCACCT	2640
2641	TTAATGAATA	ATTTCCGTCA	ATATTTACCT	TCCCTCCCTC	AATCGGTTGA	ATGTCGCCCT	2700
2701	TTTGCTTTTA	GCGCTGGTAA	ACCATATGAA	TTTTCTATTG	ATTGTGACAA	AATAAACTTA	2760
2761	TTCCGTGGTG	TCTTTGCGTT	TCTTTTATAT	GTTGCCACCT	TTATGTATGT	ATTTTCTACG	2820
2821	TTTGCTAACA	TACTGCGTAA	TAAGGAGTCT	TAATCATGCC	AGTTCTTTTG	GGTATTCGGT	2880
2881	TATTATTGCG	TTTCCTCGGT	TTCTTTCTGG	TAACCTTTGT	CGGCTATCTG	CTTACTTTTC	2940
2941	TTAAAAAGGG	CTTCGGTAAG	ATAGCTATTG	CTATTTTCATT	GTTTCTTGCT	CTTATTATTG	3000
3001	GGCTTAACCT	AATTCCTGTG	GGTTATCTCT	CTGATATTAG	CGCTCAATTA	CCCTCTGACT	3060
3061	TTGTTTCAGG	TGTTTCAGTTA	ATTCTCCCGT	CTAATCCGCT	TCCCTGTTTT	TATGTTATTTC	3120
3121	TCTCTGTAAA	GGCTGCTATT	TTTCAATTTT	ACGTTAAACA	AAAAATCGTT	TCTTATTTGG	3180
3181	ATTGGGATAA	ATAATATGGC	TGTTTATTTT	GTAACCTGGC	AATTAGGCTC	TGGAAAGACG	3240
3241	CTCGTTAGCG	TTGGTAAGAT	TTAGGATAAA	ATTGTAGCTG	GGTGCAAAAT	AGCAACTAAT	3300
3301	CTTGATTTAA	GGCTTCAAAA	CCTCCGCAAA	GTCGGGAGGT	TCGCTAAAAC	GCCTCGCGTT	3360
3361	CTTAGAATAC	CGGATAAGCC	TTCTATATCT	GATTTGCTTG	CTATTGGGCG	CGGTAATGAT	3420
3421	TCCTACGATG	AAAATAAAAA	CGGCTTGCTT	GTTCTCGATG	AGTGCGGTAC	TTGGTTTAAAT	3480
3481	ACCCGTTCTT	GGAATGATAA	GGAAGACAGG	CCGATTATTG	ATTGGTTTCT	ACATGCTCGT	3540
3541	AAATTAGGAT	GGGATATTAT	TTTTCTTGTT	CAGGACTTAT	CTATTGTTGA	TAAACAGGCG	3600
3601	CGTTCTGCAT	TAGCTGAACA	TGTTGTTTAT	TGTCGTCGTC	TGGACAGAAT	TACTTTACCT	3660
3661	TTTGTGCGTA	CTTTATATTG	TCTTATTACT	GGCTCGAAAA	TGCCTCTGCC	TAAATTACAT	3720
3721	GTTGGCGTTG	TTAAATATGG	CGATTCTCAA	TTAAGCCCTA	CTGTTGAGCG	TTGGCTTTAT	3780
3781	ACTGGTAAGA	ATTTGTATAA	CGCATATGAT	ACTAAACAGG	CTTTTTCTAG	TAATTATGAT	3840

FIG. 8-1

SUBSTITUTE SHEET

12 / 16

3841	TCCGGTGT	TTT	ATTCTTAT	TTT	AACGCCTT	AT	TTATCACACG	GTCGGTAT	TTT	CAAACCAT	TA	3900	
3901	AATTTAGG	TC	AGAAGATG	AA	GCTTACTA	AAA	ATATATTT	GA	AAAAGTTT	TC	ACGCGTTC	TT	3960
3961	TGTCTTGC	GA	TTGGATTT	GC	ATCAGCAT	TT	ACATATAG	TT	ATATAACCC	CA	ACCTAAGCC	G	4020
4021	GAGGTTAA	AA	AGGTAGTC	TC	TCAGACCT	AT	GATTTTGAT	A	AATTCAC	TAT	TGACTCTTC	T	4080
4081	CAGCGTCT	TA	ATCTAAGC	TA	TCGCTATG	TT	TTCAAGGAT	T	CTAAGGGAA	AA	ATTAATTA	AT	4140
4141	AGCGACGA	TT	TACAGAAG	CA	AGGTTATT	CA	CTCACATAT	A	TTGATTTAT	G	TACTGTTTC	C	4200
4201	ATTA AAAA	AG	GTAATTCAA	AA	TGAAATTG	TT	AAATGTAAT	T	AATTTTGTT	T	TCTTGATG	TT	4260
4261	TGTTTCAT	CA	TCTTCTTT	TG	CTCAGGTA	AT	TGAAATGA	AT	AATTCGCCT	C	TGCGCGATT	T	4320
4321	TGTAAC	TTGG	TATTCAAAG	C	AATCAGGCG	AA	ATCCGTTAT	T	GTTTCTCCG	G	ATGTAAAAG	G	4380
4381	TACTGTTA	CT	GTATATTCA	T	CTGACGTT	AA	ACCTGAAAT	T	CTACGCAAT	T	TCTTTATTC	C	4440
4441	TGTTTTAC	GT	GCTAATAAT	T	TTGATATG	GT	TGGTTCAT	T	CCTTCCATA	A	TTCAGAAG	TA	4500
4501	TAATCCAA	AC	AATCAGGAT	T	ATATTGAT	GA	ATTGCCAT	CA	TCTGATAAT	C	AGGAATAT	GA	4560
4561	TGATAATT	CC	GCTCCTTC	TG	GTGGTTTC	TT	TGTTCCGCA	AA	AATGATAAT	G	TTACTCAA	AC	4620
4621	TTTTAAAT	T	AATAACGTT	C	GGGCAAAG	GA	TTTAATAC	GA	GTTGTCGA	AT	TGTTTGTA	AA	4680
4681	GTCTAATA	CT	TCTAAATCT	T	CAAATGTAT	T	ATCTATTGA	C	GGCTCTAAT	C	TATTAGTT	GT	4740
4741	TAGTGCAC	CT	AAAGATATT	T	TAGATAACT	T	TCTCAATTC	T	CTTTCTACT	G	TTGATTG	CC	4800
4801	AACTGACC	AG	ATATTGATT	G	AGGGTTTG	AT	ATTTGAGGT	T	CAGCAAGGT	G	ATGCTTTAG	A	4860
4861	TTTTTCAT	TT	GCTGCTGG	CT	CTCAGCGT	GG	CACGTGTG	CA	GGCGGTGT	TA	ATACTGACC	G	4920
4921	CCTCACCT	CT	GTTTTATCT	T	CTGCTGGT	GG	TTCGTTCCG	GT	ATTTTTAAT	G	GCGATGTTT	T	4980
4981	AGGGCTAT	CA	GTTTCGCG	CAT	TAAAGACT	AA	TAGCCATT	CA	AAAATATT	GT	CTGTGCCAC	G	5040
5041	TATTTCTA	CG	CITTCAGGT	C	AGAAGGGT	TT	TATCTCTGT	T	GGCCAGAAT	G	TCCCTTTAT	T	5100
5101	TACTGGTC	GT	GTGACTGG	TG	AATCTGCC	AA	TGTAAATA	AT	CCATTTTCA	G	ATCTGAGCG	G	5160
5161	TCAAAATG	TA	GGTATTTCC	A	TGAGCGTT	TT	TCCTGTTG	CA	ATGGCTGG	CG	GTAAATAT	GT	5220
5221	TCTGGATAT	T	ACCAGCAAG	G	CCGATAGTT	T	GAGTTCTTC	T	ACTCAGGCA	AA	GTGATGTT	AT	5280
5281	TACTAATCA	A	AGAAGTATT	G	CTACAACG	GT	TAATTTGCG	T	GATGGACAG	A	CTCTTTTAC	T	5340
5341	CGGTGGCC	TC	ACTGATTAT	A	AAAACACT	TC	TCAAGATT	CT	GGCGTACCG	T	TCCTGTCT	AA	5400
5401	AATCCCTTT	A	ATCGGCCT	CC	TGTTTAGCT	C	CCGCTCTGA	T	TCCAACGAG	G	AAAGCACGT	T	5460
5461	ATACGTTG	CT	GTCAAAGCA	A	CCATAGTAC	G	CGCCCTGT	AG	CGGCGCATT	A	AGCGCGCG	G	5520
5521	GTGTGGTG	GT	TACGCGCAG	C	GTGACCGC	TA	CACCTGCC	AG	CGCCCTAG	CG	CCCGCTCCT	T	5580
5581	TCGCTTTC	TT	CCCTTCCT	TT	CTCGCCAC	GT	TCGCCGGCT	T	TCCCCGTCA	AA	GCTCTAAAT	C	5640
5641	GGGGGCTCC	C	TTTAGGGTT	C	CGATTTAG	TG	CTTTACGG	CA	CCTCGACCC	C	AAAAAATCT	TG	5700
5701	ATTTGGGTG	A	TGGTTCAC	GT	AGTGGGCC	AT	CGCCCTGAT	A	GACGGTTTT	T	CGCCCTTTG	A	5760
5761	CGTTGGAG	TC	CACGTTCT	TT	AATAGTGG	AC	TCTTGTTCC	A	AACTGGAAC	A	ACACTCAAC	C	5820
5821	CTATCTCGG	G	CTATTCCT	TT	GATTTATA	AG	GGATTTTGC	C	GATTTTCGG	AA	CCACCATCA	A	5880
5881	ACAGGATTT	T	CGCCTGCT	GG	GGCAAACC	AG	CGTGGACC	GC	TTGCTGCA	AC	TCTCTCAGG	G	5940
5941	CCAGGCGGT	G	AAGGGCAAT	C	AGCTGTTG	CC	CGTCTCGCT	G	GTGAAAAG	AA	AAACCACC	CT	6000
6001	GGCGCCCA	AT	ACGCAAACC	G	CCTCTCCC	CG	CGCGTTGG	CC	GATTCATT	AA	TGCAGCTGG	C	6060
6061	ACGACAGGT	T	TCCCGACT	GG	AAAGCGGG	CA	GTGAGCGCA	AA	CGCAATTA	AT	GTGAGTTAG	C	6120
6121	TCACTCATT	A	GGCACCCCA	G	GCTTTACAT	T	TTATGCTTC	C	GGCTCGTAT	G	TTGTGTGGA	A	6180
6181	TTGTGAGCG	G	ATAACAATT	T	CACACGCG	TC	ACTTGGCA	CT	GGCCGTCGT	T	TTACAACGC	T	6240
6241	GTGACTGGG	A	AAACCCTGG	C	GTTACCCA	AG	CTTTGTAC	AT	GGAGAAAAT	A	AAGTGAAAC	A	6300
6301	AAGCACTAT	T	GCACTGGCA	C	TCTTACC	GTT	ACTGTTTAC	C	CCTGTGGCA	AA	AAGCCTAT	G	6360
6361	GGGGTTCAT	G	CTTCTGAGG	C	ATCCGGGAG	C	TGAAGGCG	AT	GACCCTGCT	A	AGGCTGCAT	T	6420
6421	CAATAGTTT	A	CAGGCAAGT	G	CTACTGAG	TA	CATTGGCT	AC	GCTTGGGCT	A	TGGTAGTAG	T	6480
6481	TATAGTTGG	T	GCTACCAT	AG	GGATTAAAT	T	ATTCAAAA	AG	TTTACGAG	CA	AGGCTTCT	TA	6540
6541	AGCAATAGC	G	AAGAGGCCC	G	CACCGATCG	C	CCTTCCCA	AC	AGTTGCGCA	G	CCTGAATGG	C	6600
6601	GAATGGCG	CT	TTGCCTGG	T	TCCGGCACC	A	GAAGCGGT	GC	CGGAAAGCT	G	GCTGGAGTG	C	6660
6661	GATCTTCCT	G	AGGCCGAT	AC	GGTCGTCG	T	CCCTCAA	ACT	GGCAGATGC	A	CGGTTACG	AT	6720
6721	GCGCCCAT	CT	ACACCAAC	G	AACCTATCC	C	ATTACGGT	CA	ATCCGCCGT	T	TGTTCCAC	G	6780
6781	GAGAATCCG	A	CGGGTTGTT	A	CTCGCTCAC	A	TTTAATGTT	G	ATGAAAGCT	G	GCTACAGGA	A	6840
6841	GGCCAGAG	CT	GAATTATTT	T	TGATGGCG	TT	CCTATTGG	TT	AAAAAATG	AG	CTGATTTAA	C	6900
6901	AAAAATTTA	A	CGCGAATTT	A	AACAAAAT	AT	TAACGTTT	AC	AATTTAAAT	A	TTTGCTTAT	A	6960
6961	CAATCTTCT	C	GTTTTTGGG	G	CTTTTCTG	AT	TATCAACCG	G	GGTACATAT	G	ATTGACATG	C	7020
7021	TAGTTTTAC	G	ATTACCGTT	C	ATCGATTCT	C	TTGTTTGCT	C	CAGACTCTC	A	GGCAATGAC	C	7080
7081	TGATAGCCT	T	TGTAGATCT	C	TCAAAAAT	AG	CTACCCTCT	C	CGGCATTAT	T	TTATCAGCT	A	7140
7141	GAACGGTTG	A	ATATCATAT	T	GATGGTGAT	T	TGACTGTCT	C	CGGCCTTCT	T	CACCCTTTG	T	7200
7201	AATCTTTAC	C	TACACATT	AC	TCAGGCATT	G	CATTTAAAT	T	ATATGAGGG	T	TCTAAAAAT	T	7260
7261	TTTATCCTT	G	CGTTGAAAT	A	AAGGCTTCT	C	CCGCAAAAG	T	ATTACAGGG	T	CATAATGTT	T	7320
7321	TTGGTACA	AC	CGATTTAG	CT	TTATGCTCT	G	AGGCTTTAT	T	GCTTAATTT	T	GCTAATCTT	T	7380
7381	TGCCTTGC	CT	GTATGATT	TA	TTGGACGT	T							7409

FIG. 8-2

13 / 16

	10	20	30	40	50	60
1	AATGCTACTA	CTATTAGTAG	AATTGATGCC	ACCTTTTCAG	CTCGCGCCCC	AAATGAAAAT
61	ATAGCTAAAC	AGGTTATTGA	CCATTTGCGA	AATGTATCTA	ATGGTCAAAC	TAAATCTACT
121	CGTTGCGAGA	ATTGGGAATC	AACTGTTACA	TGGAATGAAA	CTTCCAGACA	CCGTACTTTA
181	GTTGCATATT	TAAAACATGT	TGAGCTACAG	CACCAGATTG	AGCAATTAAG	CTCTAAGCCA
241	TCTGCAAAAA	TGACCTCTTA	TCAAAAGGAG	CAATTAAGAG	TACTCTCTAA	TCCTGACCTG
301	TTGGAGTTTG	CTTCCGGTCT	GGTTTCGCTT	GAAGCTCGAA	TTAAAACGCG	ATATTGGAAG
361	TCTTTCGGGC	TTCTCTTTAA	TCTTTTTGAT	GCAATCCGCT	TTGCTTCTGA	CTATAATAGT
421	CAGGGTAAAG	ACCTGATTTT	TGATTTATGG	TCATTCTCGT	TTTCTGAACT	GTTTAAAGCA
481	TTTGAGGGGG	ATTCAATGAA	TATTTATGAC	GATTCCGCAG	TATTGGACGC	TATCCAGTCT
541	AAACATTTTA	CTATTACCCC	CTCTGGCAAA	ACTTCTTTTG	CAAAAGCCTC	TCGCTATTTT
601	GGTTTTTATC	GTCTGTGGT	AAACGAGGGT	TATGATAGTG	TTGCTCTTAC	TATGCCTCGT
661	AATTCCTTTT	GGCGTTATGT	ATCTGCATTA	GTTGAATGTG	GTATTCCTAA	ATCTCAAGT
721	ATGAATCTTT	CTACCTGTAA	TAATGTTGTT	CCGTTAGTTC	GTTTTATTAA	CGTAGATTTT
781	TCTTCCCAAC	GTCTGACTG	GTATAATGAG	CCAGTCTTAA	AAATCGCATA	AGGTAATTCA
841	CAATGATTAA	AGTTGAAATT	AAACCATCTC	AAGCCCAATT	TACTACTCGT	TCTGGTGTTC
901	CTCGTCAGGG	CAAGCCTTAT	TCACTGAATG	AGCAGCTTTG	TTACGTTGAT	TTGGGTAATG
961	AATATCCGGT	TCTTGTCAAG	ATTACTCTTG	ATGAAGGTCA	GCCAGCCTAT	GCGCCTGGTC
1021	TGTACACCGT	TCATCTGTCC	TCTTTCAAAG	TTGGTCAAGT	CGGTTCCCTT	ATGATTGACC
1081	GTCTGCGCCT	CGTTCGGGCT	AAGTAACATG	GAGCAGGTCT	CGGATTTCTG	CACAATTTAT
1141	CAGGCGATGA	TACAAATCTC	CGTTGTACTT	TGTTTCGCGC	TTGGTATAAT	CGCTGGGGGT
1201	CAAAGATGAG	TGTTTTAGTG	TATTCCTTCG	CCTCTTTCTG	TTTAGGTTGG	TGCCTTCGTA
1261	GTGGCATTAC	GTATTTTACC	CGTTTAAATG	AAACTTCCTC	ATGAAAAAGT	CTTTAGTCCT
1321	CAAAAGCCTCT	GTAGCCGTG	CTACCCTCGT	TCCGATGCTG	TCTTTTCGCT	CTGAGGGTGA
1381	CGATCCCGCA	AAAGCGGCTT	TAACTCCCTT	GCAAGCCCTA	GCGACCGAAT	ATATCGGTTA
1441	TGCGTGGGCG	ATGGTTGTTG	TCATTGTCTG	CGCAACTATC	GGTATCAAGC	TGTTTAAGAA
1501	ATTACCTCTG	AAAGCAAGCT	GATAAACCAG	TACAATTAAA	GGCTCCTTTT	GGAGCCTTTT
1561	TTTTTGGAGA	TTTTCAACGT	GAAAAAATTA	TTATTCGCAA	TTCTTTTAGT	TGTTCTTTTC
1621	TATTTCTACT	CCGCTGAAAC	TGTTGAAAGT	TGTTTAGCAA	AACCCCATAC	AGAAAAATTCA
1681	TTTACTAACG	TCTGGAAAGA	CGACAAAACT	TTAGATCGTT	ACGCTAATA	TGAGGGTTGT
1741	CTGTGGAATG	CTACAGGCGT	TGTAGTTTGT	ACTGGTGACG	AAACTCGATG	TTACGGTTACA
1801	TGGGTTCTTA	TTGGGCTTGC	TATCCCTGAA	AATGAGGGTG	GTGGCTCTGA	GGGTGGCGGT
1861	TCTGAGGGTG	GCGGTTCTGA	GGGTGGCGGT	ACTAAACCTC	CTGAGTACGG	TGATACACCT
1921	ATTCCGGGCT	ATACTTATAT	CAACCCTCTC	GACGGCACTT	ATCCGCCTGG	TACTGAGCAA
1981	AACCCCGCTA	ATCCTAATCC	TTCTCTTGAG	GAGTCTCAGC	CTCTTAATAC	TTTCATGTTT
2041	CAGAATAATA	GGTTCCGAAA	TAGGCAAGGG	GCATTAACCT	TTTATACGGG	CACTGTTACT
2101	CAAGGCACTG	ACCCCGTTAA	AACCTTATTAC	CAGTACACTC	CTGTATCACT	AAAAGCAATG
2161	TATGACGCTT	ACTGGAACGG	TAAATTCAGA	GACTGCGCTT	TCCATTCTGG	CTTTAATGAA
2221	GATCCATTCT	TTTGTGAATA	TCAAGGCCAA	TCGTCTGACC	TGCCTCAACC	TCCTGTCAAT
2281	GCTGGCGGCG	GCTCTGGTGG	TGTTTCTGGT	GGCGGCTCTG	AGGGTGGTGG	CTCTGAGGGT
2341	GGCGGTTCTG	AGGGTGGCGG	CTCTGAGGGA	GGCGGTTCCG	GTGDTGGCTC	TGGTTCGGGT
2401	GATTTTGATT	ATGAAAAGAT	GGCAAAACGT	AATAAGGGGG	CTATGACCGA	AAATGCCGAT
2461	GAAAACGCGC	TACAGTCTGA	CGCTAAAGGC	AAACTTGTAT	CTGTGCTGAC	TGATTACGGT
2521	GCTGCTATCG	ATGGTTTCAT	TGGTGACGTT	TCCGGCCTTG	CTAATGGTAA	TGGTGCTACT
2581	GGTGATTTTG	CTGGCTCTAA	TTCCCAAATG	GCTCAAGTCG	GTGACGGTGA	TAATTCACCT
2641	TTAATGAATA	ATTTCCGTCA	ATATTTACCT	TCCCTCCCTC	AATCGGTTGA	ATGTCGCCCT
2701	TTTGTCTTTA	GCGCTGGTAA	ACCATATGAA	TTTTCTATTG	ATTGTGACAA	AATAAACTTA
2761	TTCCGTGGTG	TCTTTGCGTT	TCTTTTATAT	GTTGCCACCT	TTATGTATGT	ATTTTCTACG
2821	TTTGCTAACA	TACTGCGTAA	TAAGGAGTCT	TAACTATGCC	AGTTCTTTTG	GGTATTCCGT
2881	TATTATTGCG	TTTCCTCGGT	TTCTTCTGGT	TAACTTTGTT	CGGCTATCTG	CTTACTTTTC
2941	TTAAAAAGGG	CTTCGGTAAG	ATAGCTATTG	CTATTTTCAT	GTTTCTTGCT	CTTATTATTG
3001	GGCTTAACTC	AATTCTTGTT	GGTTATCTCT	CTGATATTAG	CGCTCAATTA	CCCTCTGACT
3061	TTGTTCAAGG	TGTTCAAGTT	ATTCTCCCGT	CTAATGCGCT	TCCCTGTTTT	TATGTTATTC
3121	TCTCTGTAAA	GGCTGCTATT	TTCAATTTTT	ACGTTAAACA	AAAAATCGTT	TCTTATTTGG
3181	ATTGGGATAA	ATAATATGGC	TGTTTATTTT	GTAATGGCA	AATTAGGCTC	TGGAAGAACG
3241	CTCGTTAGCG	TTGGTAAGAT	TCAGGATAAA	ATTGTAGCTG	GGTGCAAAAT	AGCAAACTAT
3301	CTTGATTTAA	GGCTTCAAAA	CCTCCCGCAA	GTCGGGAGGT	TCGCTAAAAC	GCCTCGCGTT
3361	CTTAGAATAC	CGGATAAGCC	TTCTATATCT	GATTTGCTTG	CTATTGGGCG	CGGTAATGAT
3421	TCCTACGATG	AAAAATAAAA	CGGCTTGCTT	GTTCTCGATG	AGTGCGGTAC	TTGGTTTAAAT
3481	ACCCGTTCTT	GGAATGATAA	GGAAAGACAG	CCGATTATTG	ATTGGTTTCT	ACATGCTCGT
3541	AAATTAGGAT	GGGATATTAT	CTTCTTTGTT	CAGGACTTAT	CTATTGTTGA	TAAACAGGCG
3601	CGTTCTGCAT	TAGCTGAACA	TGTTGTTTAT	TGTCGTCGTC	TGGACAGAAT	TACTTTACCT
3661	TTTGTGCGTA	CTTTATATTC	TCTTATTACT	GGCTCGAAAA	TGCCTCTGCC	TAAATTACAT
3721	GTTGGCGTTG	TTAAATATGG	CGATTCTCAA	TTAAGCCCTA	CTGTTGAGCG	TTGGCTTTAT
3781	ACTGGTAAGA	ATTTGTATAA	CGCATATGAT	ACTAAACAGG	CTTTTCTAG	TAATTATGAT

FIG. 9-1

SUBSTITUTE SHEET

14 / 16

3841	TCCGGTGT	TTTCTTAT	AACGCCTT	TTATCACAC	GTCGGTAT	CAAACCAT	3900
3901	AATTTAGG	AGAAGATG	GCTTACTA	ATATATTT	AAAAGTTT	ACGCGTTC	3960
3961	TGTCTTGC	TTGGATTT	ATCAGCAT	ACATATAG	ATATAACCC	ACCTAAGCC	4020
4021	GAGGTTAA	AGGTAGTC	TCAGACCT	GATTTTGAT	AATTCACAT	TGACTCTTC	4080
4081	CAGCGTCT	ATCTAAGC	TCGCTATG	TTCAAGGAT	CTAAGGGAA	ATTAATTA	4140
4141	AGCGACGA	TACAGAAG	AGGTTATT	CTCACATAT	TTGATTTAT	TACTGTTTC	4200
4201	ATTAATAA	TAATTCAA	GAAATGTT	AATGTAATT	ATTTTGTTT	CTTGATGTT	4260
4261	GTTTCATC	CTTCTTTG	TCAGGTAAT	GAAATGAAT	ATTGCGCT	GCGCGATTT	4320
4321	GTAACCTG	ATTCAAAG	ATCAGGCG	TCCGTTATT	TTTCTCCCG	TGTAAAAGG	4380
4381	ACTGTTAC	TATATTCAT	TGACGTTAA	CCTGAAATC	TACGCAATT	CTTTATTTCT	4440
4441	GTTTTACG	CTAATAATT	TGATATGGT	CCTTCAATTC	CTTCCATTAT	TTAGAAGTAT	4500
4501	AATCCAAAC	ATCAGGATTA	TATTGATGAA	TTGCCATCAT	CTGATAATCA	GGAAATATGAT	4560
4561	GATAATCCG	CTCCTTCTG	TGGTTTCTT	GTTCCGCAAA	ATGATAATGT	TACTCAAAC	4620
4621	TTTAAAAAT	ATAACGTTG	GGCAAAGGAT	TTAATACGAG	TTGTCGAATT	GTTTGTAAG	4680
4681	TCTAATACT	CTAAATCCTC	AAATGTATTA	TCTATTGACG	GCTCTAATCT	ATTAGTTGTT	4740
4741	AGTGCACCTA	AAGATATTTT	AGATAACCTT	CCTCAATTCC	TTTCTACTGT	TGATTTGCCA	4800
4801	ACTGACCAGA	TATTGATTGA	GGGTTTGATA	TTTGAGGTTT	AGCAAGGTGA	TGCTTTAGAT	4860
4861	TTTTCATTTG	CTGCTGGCTC	TCAGCGTG	ACTGTTGCAG	GCGGTGTTAA	TACTGACCGC	4920
4921	CTCACCTCTG	TTTTATCTTC	TGCTGGTGGT	TCGTTCCGTA	TTTTTAATGG	CGATGTTTTA	4980
4981	GGGCTATCAG	TTTCGCGCATT	AAAGACTAAT	AGCCATTCAA	AAATATTGTC	TGTGCCACGT	5040
5041	ATTCTTACGC	TTTCAGGTTCA	GAAGGGTTCT	ATCTCTGTTG	GCCAGAATTG	CCCTTTTATT	5100
5101	ACTGGTCGTG	TGACTGGTGA	ATCTGCCAAT	GTAATAATC	CATTTTCAGAC	CATTGAGCGT	5160
5161	CAAAATGTAG	GTATTTCCAT	GAGCGTTTTT	CCTGTTGCAA	TGGCTGGCGG	TAATATTGTT	5220
5221	CTGGATATTA	CCAGCAAGGC	CGATAGTTTG	AGTTCTTCTA	CTCAGGCAAG	TGATGTTATT	5280
5281	ACTAATCAAA	GAAGTATTGC	TACAACGGTT	AATTTGCGTG	ATGGACAGAC	TCTTTTACTC	5340
5341	GGTGGCCTCA	CTGATTATAA	AAACACTTCT	CAAGATTCTG	GCGTACCCTT	CCTGTCTAAA	5400
5401	ATCCCTTTAA	TCGGCCTCCT	GTTTAGCTCC	CGCTCTGATT	CCAACGAGGA	AAGCAGCTTA	5460
5461	TACGTGCTCG	TCAAAGCAAC	CATAGTACGC	GCCCTGTAGC	GGCGCATTA	GCGCGGCGGG	5520
5521	TGTGGTGGTT	ACGCGCAGCG	TGACCCTAC	ACTTGCCAGC	GCCCTAGCGC	CCGCTCCTTT	5580
5581	CGCTTCTTTC	CCTTCCTTTC	TCGCCACGTT	CGCCGGCTTT	CCCCGTCAAG	CTCTAAATCG	5640
5641	GGGGCTCCCT	TTAGGGTTCC	GATTTAGTGC	TTTACGGCAC	CTCGACCCCA	AAAAACTTGA	5700
5701	TTTGGGTGAT	GGTTCACGTA	GTGGGCCATC	GCCCTGTATG	ACGGTTTTTC	GCCCTTTGAC	5760
5761	GTTGGAGTCC	ACGTTCTTTA	ATAGTGGACT	CTTGTTCCAA	ACTGGAACAA	CCTCAACCC	5820
5821	TATCTCGGGC	TATTCTTTTG	ATTTATAAGG	GATTTTGCCG	ATTTGGAAC	CACCATCAAA	5880
5881	CAGGATTTTC	GCCTGCTGGG	GCAAACACAG	GTGGACCGCT	TGCTGCAACT	CTCTCAGGGC	5940
5941	CAGGCGGTGA	AGGGCAATCA	GCTGTTGCC	GTCTCGCTGG	TGAAAAGAAA	AACCACCCTG	6000
6001	GCGCCCAATA	CGCAAACCGC	CTCTCCCCGC	GCGTTGGCCG	ATTCATTAAT	GCAGCTGGCA	6060
6061	CGACAGGTTT	CCCGACTGGA	AAGCGGGCAG	TGAGCGCAAC	GCAATTAATG	TGAGTTAGCT	6120
6121	CACCTCATTA	GCACCCAGG	CTTTACACTT	TATGCTTCCG	GCTCGTATGT	TGTGTGGAAT	6180
6181	TGTGAGCGGA	TAACAATTTT	ACACAGGAAA	CAGCTATGAC	CAGGATGTAC	GAATTCGCAG	6240
6241	GTAGGAGAGC	TCGGCGGATC	CGAGGCTGAA	GGCGATGACC	CTGCTAAGGC	TGCATTCAAT	6300
6301	AGTTTACAGG	CAAGTGCTAC	TGAGTACATT	GGCTACGCTT	GGGCTATGGT	AGTAGTTATA	6360
6361	GTTGGTGCTA	CCATAGGGAT	TAAATTATTC	AAAAAGTTTA	CGAGCAAGGC	TTCTTAACCA	6420
6421	GCTGGCGTAA	TAGCGAAGAG	GCCCGCACCG	ATCGCCCTTC	CCAACAGTTG	CGCAGCCTGA	6480
6481	ATGGCGAATG	GCGCTTTGCC	TGGTTTCCGG	CACCAAGAGC	GGTGCCGGAA	AGCTGGCTGG	6540
6541	AGTGCGATCT	TCCTGAGGCC	GATACGGTCG	TCGTCCCTC	AAACTGGCAG	ATGCACGGTT	6600
6601	ACGATGCGCC	CATCTACACC	AACGTAACCT	ATCCCATTA	GGTCAATCCG	CCGTTTGTTC	6660
6661	CCACGGAGAA	TCCGACGGGT	TGTTACTCGC	TCACATTTAA	TGTTGATGAA	AGCTGGCTAC	6720
6721	AGGAAGGCCA	GACGCGAATT	ATTTTGTATG	GCGTTCCTAT	TGGTTAAAAA	ATGAGCTGAT	6780
6781	TTAACAAAAA	TTTAACGCGA	ATTTTAAACA	AATATTAACG	TTTACAATTT	AAATATTTGC	6840
6841	TTATACAAAT	TTCTGTGTTT	TGGGGCTTTT	CTGATTATCA	ACCGGGGTAC	ATATGTTTGA	6900
6901	CATGCTAGTT	TTACGATTAC	CGTTTCATCGA	TTCTCTTGTT	TGCTCCAGAC	TCTCAGGCAA	6960
6961	TGACCTGATA	GCCTTTGTAG	ATCTCTCAAA	AATAGCTACC	CTCTCCGGCA	TTAATTTATC	7020
7021	AGCTAGAACG	GTTGAATATC	ATATTGATGG	TGATTTGACT	GTCTCCGGCC	TTTCTCACCC	7080
7081	TTTTGAATCT	TTACCTACAC	ATTACTCAGG	CATTGCATT	AAAAATATAG	AGGGTTCTAA	7140
7141	AAATTTTTAT	CCTTGCGTTG	AAATAAAGGC	TTCTCCCGCA	AAAGTATTAC	AGGGTCATAA	7200
7201	TGTTTTGAT	ACAACCGATT	TAGCTTTATG	CTCTGAGGCT	TTATTGCTTA	ATTTTGCTAA	7260
7261	TTCTTTGCCT	TGCCTGTATG	ATTTATTGGA	CGTT			7294

FIG. 9-2

15 / 16

	10	20	30	40	50	60
1	AATGCTACTA	CTATTAGTAG	AATTGATGCC	ACCTTTTTCAG	CTCGCGCCCC	AAATGAAAAT
61	ATAGCTAAAC	AGGTTATTGA	CCATTTGCGA	AATGTATCTA	ATGGTCAAAC	TAAATCTACT
121	CGTTTCGAGA	ATTGGGAATC	AACTGTTACA	TGGAATGAAA	CTTCCAGACA	CCGTACTTTA
181	GTTGCATATT	TAAAACATGT	TGAGCTACAG	CACCAGATTG	AGCAATTAAG	CTCTAAGCCA
241	TCTGCAAAAA	TGACCTCTTA	TCAAAAGGAG	CAATTAAGG	TACTCTCTAA	TCCTGACCTG
301	TTGGAGTTTG	CTTCCGGTCT	GGTTCGCTTT	GAAGCTCGAA	TTAAAACGCG	ATATTTGAAG
361	TCTTTCGGGC	TTCTCTTTAA	TCTTTTTGAT	TCATTTCCGT	TTCTGAACT	GTATAATAGT
421	CAGGGTAAAG	ACCTGATTTT	TGATTTATGG	GCAATCCGCT	TTCTGAACT	GTTTAAAGCA
481	TTTGAGGGGG	ATTCAATGAA	TATTTATGAC	GATTCCGCG	TATTGGACGC	TATCCAGTCT
541	AAACATTTTA	CTATTACCCC	CTCTGGCAAA	ACTTCTTTTG	CAAAAAGCCTC	TCGCTATTTT
601	GGTTTTTATC	GTCGTCTGGT	AAACGAGGGT	TATGATAGTG	TTGCTCTTAC	TATGCCTCGT
661	AATTCCTTTT	GGCGTTATGT	ATCTGCATTA	GTTGAATGTG	GTATTCCTAA	ATCTCAACTG
721	ATGAATCTTT	CTACCTGTAA	TAATGTTGTT	CCGTTAGTTC	GTATTCCTAA	CGTAGATTTT
781	TCTTCCCAAC	GTCCTGACTG	GTATAATGAG	CCAGTTCTTA	AAATCGCATA	AGGTAATTCA
841	CAATGATTAA	AGTTGAAATT	AAACCATCTC	AAGCCCAATT	TACTACTCGT	TCTGGTGTTC
901	CTCGTCAGGG	CAAGCCTTAT	TCACTGAATG	AGCAGCTTTG	TTACGTTGAT	TTGGGTAATG
961	AATATCCGGT	TCTTGTCAAG	ATTACTCTTG	ATGAAGGTCA	GCCAGCCTAT	GCGCCTGGTC
1021	TGTACACCGT	TCATCTGTCC	TCTTTCAAAG	TTGGTCAGTT	CGGTTCCCTT	ATGATTGACC
1081	GTCTGCGCCT	CGTTCCGGCT	AAGTAACATG	GAGCAGGTG	CGGATTTCTA	CACAAATTTT
1141	CAGGCGATGA	TACAAATCTC	CGTTGTACTT	TGTTTCGCGC	TTGGTATAAT	CGCTGGGGGT
1201	CAAAGATGAG	TGTTTTAGTG	TATTTCTTCG	CCTCTTTCTG	TTTAGGTTGG	TGCTTTCGTA
1261	GTGGCATTAC	GTATTTTACC	CGTTTAAATG	AAACTTCCTC	ATGAAAAAGT	CTTTAGTCCT
1321	CAAAGCCTCT	GTAGCCGTTG	CTACCCTCGT	TCCGATGCTG	TCTTTTCGCTG	CTGAGGGTGA
1381	CGATCCCGCA	AAAGCGGCCT	TTAACTCCCT	GCAAGCCTCA	GCGACCGAAT	ATATCGGTTA
1441	TGCGTGGGCG	ATGGTTGTTG	TCAATGTGCG	CGCAACTATC	GGTATCAAGC	TGTTTAAAGA
1501	ATTACACCTCG	AAAGCAAGCT	GATAAACCGA	TACAATTAAT	GGCTCCTTTT	GGAGCCTTTT
1561	TTTTTGGAGA	TTTTCAACGT	GAAAAAATTA	TTATTCGCAA	TTCTTTTAGT	TGTTCTTTTC
1621	TATTCTCACT	CCGCTGAAAC	TGTTGAAAGT	TGTTTAGCAA	AACCCCATAC	AGAAAATTCA
1681	TTTACTAACG	TCTGGAAAGA	GCACAAAAC	TTAGATCGTT	ACGCTAACTA	TGAGGGTTGT
1741	CTGTGGAATG	CTACAGGCGT	TGTAGTTTGT	ACTGGTGACG	AAACTCAGTG	TTACGGTACA
1801	TGGGTTCCTA	TTGGGCTTGC	TATCCCTGAA	AATGAGGGTG	GTGGCTCTGA	GGGTGGCGGT
1861	TCTGAGGGTG	GCGGTTCTGA	GGGTGGCGGT	ACTAAACCTC	CTGAGTACGG	TGATACACCT
1921	ATTCCGGGCT	ATACTTATAT	CAACCCTCTC	GACGGCACTT	ATCCGCCTGG	TACTGACCAA
1981	AACCCCGCTA	ATCCTAATCC	TTCTCTTGAG	GAGTCTCAGC	CTCTTAATAC	TTTCATGTTT
2041	CAGAATAATA	GGTTCCGAAA	TAGGCAGGGG	GCATTAACCT	TTTATACGGG	CACCTGTTACT
2101	CAAGGCATCG	ACCCCGTTAA	AACTTATTAC	CAGTACACTC	CTGTATCATC	AAAAGCCATG
2161	TATGACGCTT	ACTGGAACGG	TAAATTCAGA	GACTGCGCTT	TCCATTCTGG	CTTTAATGAA
2221	GATCCATTCT	TTTGTGAATA	TCAAGGCCAA	TGCTTGACC	TGCTCAACCT	CTCTGTTCAAT
2281	GCTGGCGGCG	GCTCTGGTGG	TGGTCTGGT	GGCGGCTCTG	AGGGTGGTGG	CTCTGAGGGT
2341	GGCGGTTCTG	AGGGTGGCGG	CTCTGAGGGA	GGCGGTTCCG	GTGGTGGCTC	TGTTTCCGGT
2401	GATTTTGATT	ATGAAAAGAT	GGCAAAACGCT	AATAAGGGGG	CTATGACCGA	AAATGCCGAT
2461	GAAAAACGCG	TACAGTCTGA	CGCTAAAGGC	AAACTTGATT	CTGTCGCTAC	TGATTACGGT
2521	GCTGCTATCG	ATGGTTTCAT	TGGTGACGTT	TCCGGCCTTG	CTAATGGTAA	TGGTGCTACT
2581	GGTGATTTTG	CTGGCTCTAA	TTCCCAAAATG	GCTCAAGTCG	GTGACGGTGA	TGAATCAACT
2641	TTAATGAATA	ATTTCCGTCA	ATATTTACCT	TCCCTCCCTC	AATCGGTTGA	ATGTCGCCCT
2701	TTTGTCTTTA	GCGCTGGTAA	ACCATATGAA	TTTTCTATTG	ATTGTGACAA	AATAAACTTA
2761	TTCCGTGGTG	TCTTTGCGTT	TCTTTTATAT	GTTGCCACCT	TTATGTATGT	ATTTTCTACG
2821	TTTGCTAACA	TACTGCGTAA	TAAGGAGTCT	TAATCATGCC	AGTTCTTTTG	GGTATTCCGT
2881	TATTATTGCG	TTTCCTCGGT	TTCTCTTCTG	TAACCTTTGT	CGGCTATCTG	CTTACTTTTC
2941	TTAAAAAGGG	CTTCGGTAAAG	ATAGCTATTG	CTATTTCATT	GTTCCTTGCT	CTTATTATTG
3001	GGCTTAACTC	AATTCTTGTG	GGTTATCTCT	CTGATATTAG	CGCTCAATTA	CCCTCTGACT
3061	TTGTTACAGG	TGTTACAGTTA	ATTCTCCCGT	CTAATGCGCT	TCCCTGTTTT	TATGTTATTC
3121	TCTCTGTAAA	GGCTGCTATT	TTCATTTTTG	ACGTTAAACA	AAAAATCGTT	TCTTATTTGG
3181	ATTGGGATAA	ATAATATGGC	TGTTTATTTT	GTAAGTGGCA	AATTAGGCTC	TGGAAAGACG
3241	CTCGTTAGCG	TTGGTAAGAT	TTAGGATAAA	ATTGTAGCTG	GGTGCAAAAT	AGCAACTAAT
3301	CTTGATTTAA	GGCTTCAAAA	CCTCCGCAAA	GTCGGGAGGT	TCGCTAAAAA	GCCTCGCGTT
3361	CTTAGAATAC	CGGATAAGCC	TCTATATCT	GATTTGCTTG	CTATTGGGCG	CGGTAATGAT
3421	TCCTACGATG	AAAATAAAAA	CGGCTTGCTT	GTTCTCGATG	AGTGCGGTAC	TTGGTTTAAAT
3481	ACCCGTTCTT	GGAATGATAA	GGAAAGACAG	CCGATTATTG	ATTGGTTTCT	ACATGCTCGT
3541	AAATTAGGAT	GGGATATTAT	TTTTCTTGTT	CAGGACTTAT	CTATTGTTGA	TAAACAGGCG
3601	CGTTCTGCAT	TAGCTGAACA	TGTTGTTTTT	TGTCGTCGTC	TGGACAGAAAT	TACTTTACCT
3661	TTTGTCCGTA	CTTTATATT	GTTTATTACT	GGCTCGAAAA	TGCCTCTGCC	TAAATTACAT
3721	GTTGGCGTTG	TTAAATATGG	CGATTCTCAA	TTAAGCCCTA	CTGTTGAGCG	TTGGCTTTAT
3781	ACTGGTAAGA	ATTTGTATAA	CGCATATGAT	ACTAAACAGG	CTTTTTCTAG	TAATTATGAT

FIG. 10-1

SUBSTITUTE SHEET

16 / 16

3841	TCCGGTGTTT	ATTCTTATTT	AACGCCTTAT	TTATCACACG	GTCGGTATTT	CAAACCATTA	3900
3901	AATTTAGGTC	AGAAGATGAA	GCTTACTAAA	ATATATTTGA	AAAAGTTTTT	ACGCGTTCTT	3960
3961	TGTCTTGCGA	TTGGATTTGC	ATCAGCATTT	ACATATAGTT	ATATAACCCA	ACCTAAGCCG	4020
4021	GAGGTTAAAA	AGGTAGTCTC	TCAGACCTAT	GATTTTGATA	AATTCACAT	TGACTCTTCT	4080
4081	CAGCGTCTTA	ATCTAAGCTA	TCGCTATGTT	TTCAAGGATT	CTAAGGGAAA	ATTAATTAAT	4140
4141	AGCGACGATT	TACAGAAGCA	AGGTTATTCA	CTCACATATA	TTGATTTATG	TACTGTTTCC	4200
4201	ATTAAAAAAG	GTAATTCAAA	TGAAATTGTT	AAATGTAATT	AATTTTGTTT	TCTTGATGTT	4260
4261	TGTTTCATCA	TCTTCTTTTG	CTCAGGTAAT	TGAAATGAAT	AATTCGCCTC	TGCGCGATTT	4320
4321	TGTAACCTGG	TATTCAAAGC	AATCAGGCGA	ATCCGTTATT	GTTTCTCCCG	ATGTAAAAGG	4380
4381	TACTGTTACT	GTATATTCAT	CTGACGTTAA	ACCTGAAAAT	CTACGCAATT	TCTTTATTTC	4440
4441	TGTTTTACGT	GCTAATAATT	TTGATATGGT	TGGTTCAATT	CCTTCCATAA	TTCAGAAAGTA	4500
4501	TAATCCAAAC	AATCAGGATT	ATATTGATGA	ATTGCCATCA	TCTGATAATC	AGGAATATGA	4560
4561	TGATAATTCC	GCTCCTTCTG	GTGGTTTCTT	TGTTCCGCAA	AATGATAATG	TTACTCAAAC	4620
4621	TTTTAAAAAT	AATAACGTTT	GGGCAAAGGA	TTTAATACGA	GTTGTGGAAT	TGTTTGTAAG	4680
4681	GTCTAAACT	TCTAAATCCT	CAAATGTATT	ATCTATTGAC	GGCTCTAATC	TATTAGTTGT	4740
4741	TAGTGCACCT	AAAGATATTT	TAGATAACCT	TCCTCAATTC	CTTTCTACTG	TTGATTTGCC	4800
4801	AACTGACCAG	ATATTGATTG	AGGGTTTGAT	ATTTGAGGTT	CAGCAAGGTG	ATGCTTTAGA	4860
4861	TTTTTCATTT	GCTGCTGGCT	CTCAGCGTGG	CACTGTTGCA	GGCGGTGTTA	ATACTGACCG	4920
4921	CCTCACCTCT	GTTTTATCTT	CTGCTGGTGG	TTCGTTCCGT	ATTTTTAATG	GCGATGTTTT	4980
4981	AGGGCTATCA	GTTCCGCGAT	TAAAGACTAA	TAGCCATTCA	AAAATATTGT	CTGTGCCACG	5040
5041	AATCTTACG	CTTTCAGGTC	AGAAGGGTTC	TATCTCTGTT	GGCCAGAATG	TCCCTTTTAT	5100
5101	TACTGGTCGT	GTGACTGGTG	AATCTGCCAA	TGTAATAAAT	CCATTTGAGA	CGATTGAGCG	5160
5161	TCAAAATGTA	GGTATTTCCA	TGAGCGTTTT	TCCTGTTGCA	ATGGCTGGCG	GTAATATTGT	5220
5221	TCTGGATATT	ACCAGCAAGG	CCGATAGTTT	GAGTTCCTCT	ACTCAGGCAA	GTGATGTTAT	5280
5281	TACTAATCAA	AGAAGTATTG	CTACAACGGT	TAATTTGCGT	GATGGACAGA	CTCTTTTACT	5340
5341	CGGTGGCCTC	ACTGATTATA	AAAACACTTC	TCAAGATTCT	GGCGTACCGT	TCCGTCTAA	5400
5401	AATCCCTTTA	ATCGGCCTCC	TGTTTAGCTC	CCGCTCTGAT	TCCAACGAGG	AAAGCAGGTT	5460
5461	ATACGTGCTC	GTCAAAGCAA	CCATAGTACG	CGCCCTGTAG	CGGCGCATTA	AGCGCGGCGG	5520
5521	GTGTGGTGGT	TACGCGCAGC	GTGACCGCTA	CACTTGCCAG	CGCCCTAGCG	CCCCTCCTT	5580
5581	TCGCTTTCTT	CCCTTCCTTT	CTCGCCACGT	TCGCCGCTT	TCCCCGTCAA	GCTCTAAATC	5640
5641	GGGGGCTCCC	TTTAGGGTTC	CGATTTAGTG	CTTTACGGCA	CCTCGACCCC	AAAAAACTTG	5700
5701	ATTTGGGTGA	TGGTTCACGT	AGTGGGCCAT	CGCCCTGATA	GACGGTTTTT	CGCCCTTTGA	5760
5761	CGTTGGAGTC	CACGTTCTTT	AATAGTGGAC	TCTTGTTCCA	AACTGGAACA	ACACTCAACC	5820
5821	CTATCTCGGG	CTATTCTTTT	GATTATATAAG	GGATTTTGCC	GATTTGCGAA	CCACCATCAA	5880
5881	ACAGGATTTT	CGCCTGCTGG	GGCAAACGAG	CGTGGACCGC	TTGCTGCAAC	TCTCTCAGGG	5940
5941	CCAGGCGGTG	AAGGGCAATC	AGCTGTTGCC	CGTCTCGCTG	GTGAAAAGAA	AAACCACCTT	6000
6001	GGCGCCCAAT	ACGCAAACCG	CCTCTCCCCG	CGCGTTGGCC	GATTCATTAA	TGCAGCTGGC	6060
6061	ACGACAGGTT	TCCCGACTGG	AAAGCGGGCA	GTGAGCGCAA	CGCAATTAAT	GTGAGTTAGC	6120
6121	TCACTCATTA	GGCACCCAG	GCTTTACACT	TTATGCTTCC	GGCTCGTATG	TTGTGTGGA	6180
6181	TTGTGAGCGG	ATAACAATTT	CACACGCGTC	ACTTGGCACT	GGCCGTCGTT	TTACAACGTC	6240
6241	GTGACTGGGA	AAACCCTGGC	GTTACCCAAG	CTTTGTACAT	GGAGAAAATA	AAGTGAAACA	6300
6301	AAGCACTATT	GCACTGGCAC	TCTTACCGTT	ACTGTTTACC	CCTGTGGCAA	AAGCCCTTCT	6360
6361	GAGGCATCCG	GGAGCTGAAG	GCGATGACCC	TGCTAAGGCT	GCATTCAATA	GTTTACAGGC	6420
6421	AAGTGCTACT	GAGTACATTG	GCTACGCTTG	GGCTATGGTA	GTAGTTATAG	TTGGTGCTAC	6480
6481	CATAGGGATT	AAATTATTCA	AAAAGTTTAC	GAGCAAGGCT	TCTTAAGCAA	TAGCGAAGAG	6540
6541	GCCCCGACCG	ATCGCCCTTC	CCAACAGTTG	CGCAGCCTGA	ATGGCGAATG	GCGCTTTGCC	6600
6601	TGGTTTCCGG	CACCAGAAGC	GGTGCCGGAA	AGCTGGCTGG	AGTGCGATCT	TCCTGAGGCC	6660
6661	GATACGGTCG	TCGTCCCCTC	AAACTGGCAG	ATGCACGGTT	ACGATGCGCC	CATCTACACC	6720
6721	AACGTAACCT	ATCCCATTTAC	GGTCAATCCG	CCGTTTGTTT	CCACGGAGAA	TCCGACGGGT	6780
6781	TGTTACTCGC	TCACATTTAA	TGTTGATGAA	AGCTGGCTAC	AGGAAGGCCA	GACGCGAATT	6840
6841	ATTTTGTATG	GCGTTCCTAT	TGTTTAAAAA	ATGAGCTGAT	TTAACAATAA	TTTAACGCGA	6900
6901	ATTTTAAACA	AATATTAAAC	TTTACAATTT	AAATATTTGC	TTATACAATC	TTCTGTTTTT	6960
6961	TGGGGCTTTT	CTGATTATCA	ACCGGGGTAC	ATATGATTGA	CATGCTAGTT	TTACGATTAC	7020
7021	CGTTCATCGA	TTCTCTTGTT	TGCTCCAGAC	TCTCAGGCAA	TGACCTGATA	GCCTTTGTAG	7080
7081	ATCTCTCAAA	AATAGCTACC	CTCTCCGGCA	TTAATTTATC	AGCTAGAACG	GTTGAATATC	7140
7141	ATATTGATGG	TGATTTGACT	GTCTCCGGCC	TTTCTCACCC	TTTTGAATCT	TTACCTACAC	7200
7201	ATTACTCAGG	CATTGCATTT	AAAATATATG	AGGGTCTAA	AAATTTTTAT	CCTTGCGTTG	7260
7261	AAATAAAGGC	TTCTCCCGCA	AAAGTATTAC	AGGGTCATAA	TGTTTTTGGT	ACAACCGATT	7320
7321	TAGCTTTATG	CTCTGAGGCT	TTATTGCTTA	ATTTTGCTAA	TTCTTTGCCT	TGCCTGTATG	7380
7381	ATTTATTGGA	CGTT					7394

FIG. 10-2

SUBSTITUTE SHEET

INTERNATIONAL SEARCH REPORT

International Application No. PCT/US91/07141

I. CLASSIFICATION OF SUBJECT MATTER (If several classification symbols apply, indicate all) ⁶		
According to International Patent Classification (IPC) or to both National Classification and IPC		
IPC(5): C12N 1/24, 15/00; C07H 21/00		
U.S. CL.: 435/252.33, 320.1, 172.3; 536/27		
II. FIELDS SEARCHED		
Minimum Documentation Searched ⁷		
Classification System	Classification Symbols	
U.S.	435/252.33, 320.1, 172.3, 69.1; 536/27	
Documentation Searched other than Minimum Documentation to the Extent that such Documents are Included in the Fields Searched ⁸		
APS, CAS: search terms: Codon bins, codon preference		
III. DOCUMENTS CONSIDERED TO BE RELEVANT ⁹		
Category ⁹	Citation of Document, ¹¹ with indication, where appropriate, of the relevant passages ¹²	Relevant to Claim No. ¹³
Y	EP. A. 0.383.620 (Cook) 22 August 1990. See entire document.	1-87
Y	US. A. 4,458.066 (Caruthers et al.) 03 July 1984. see entire document.	1-87
Y	US. A. 4.771.000 (Verrips et al.) 13 September 1988. see entire document.	8.9.24-26 32-34. 55-57. 64-66. 73-75. 81-87
Y	APPLIED MICROBIOLOGY AND BIOTECHNOLOGY. Volume 21. issued 1985. J.M. Jaynes et al., "Construction and expression of synthetic DNA fragments coding for polypeptides with elevated levels of essential amino acids". pages 200-205. see entire document.	1-87
<div style="display: flex; justify-content: space-between;"> <div style="width: 45%;"> <p>¹⁰ Special categories of cited documents:</p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> </div> <div style="width: 45%;"> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step</p> <p>"Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>"Z" document member of the same patent family</p> </div> </div>		
IV. CERTIFICATION		
Date of the Actual Completion of the International Search		Date of Mailing of this International Search Report
11 December 1991		22 JAN 1992
International Searching Authority		Signature of Authorized Officer
ISA/US		James Ketter ebw

III DOCUMENTS CONSIDERED TO BE RELEVANT (CONTINUED FROM THE SECOND SHEET)		
Category *	Citation of Document, with indication, where appropriate, of the relevant passages	Relevant to Claim No
Y	GENE, Volume 44, issued 1986. A.R. Oliphant, "Cloning of random-sequence oligodeoxynucleotides", pages 177-183, see entire document.	1-87
Y	PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCE, Volume 87, issued August 1990, Cwirla et al., "Peptides on phage: A vast library of peptides for identifying ligands", pages 6378-6382. see entire document.	1-87
Y	SCIENCE, Volume 249, issued 27 July 1990, J.J. Devlin, "Random Peptide Libraries: A Source of Specific Protein Binding Molecules", pages 404-406, see entire document.	1-87
Y	SCIENCE, Volume 249, issued 27 July 1990, J.K. Scott, "Searching for Peptide Ligands with an Epitope Library", pages 386-390, see entire document.	1-87
Y	E.-L. WINNACKER, "From Genes to Clones: Introduction to Gene Technology", published 1987 by VCH VmbH (Weinheim, Germany), See pages 276-279, especially Table 7-4.	1-87
Y	SCIENCE, Volume 228, issued 14 June 1985, G.P. Smith, "Filamentous Fusion Phage: Novel Expression Vectors That Display Cloned Antigens on the Virion Surface", pages 1315-1317. see entire document.	8.9.24-26. 32-34. 55-57. 64-66. 73-75. 81-87